ALBERT R. MANN
LIBRARY
CORNELL UNIV.

SHOUT
SHARE
IT

The original of this book is in
the Cornell University Library.

There are no known copyright restrictions in
the United States on the use of the text.

http://www.archive.org/details/cu31924000615397

M94

SYSTEMATIC CENSUS

OF

AUSTRALIAN PLANTS,

WITH

Chronologic, Literary and Geographic Annotations;

BY

BARON FERDINAND VON MUELLER,

K.C.M.G., M.D., PH.D., F.R.S.,

F.L.S., F.G.S., F.R.G.S., C.M.Z.S., H.F.R.H.S. &c.

GOVERNMENT BOTANIST FOR THE COLONY OF VICTORIA.

PART I.—VASCULARES.

"LEVATE OCULOS IN EXCELSUM ET VIDETE, QUIS CREAVIT HAEC OMNIA, QUIS EDUCIT MULTITUDINEM EORUM."—*Isaiah* xl. 26.

MELBOURNE:

PRINTED FOR THE VICTORIAN GOVERNMENT

BY M'CARRON, BIRD & CO., 37 FLINDERS LANE WEST.

1882.

QK431
M94.

To

GEORGE BENTHAM, Esq., F.R.S., V.P.L.S.,

TO

Sir JOSEPH HOOKER, K.C.S.I., C.B., F.R.S., V.P.L.S.

AND TO

M. ALPHONSE DE CANDOLLE, F.M.R.S., F.M.L.S.,

WHO AS HEIRS OF GREAT NAMES WORTHILY SUSTAIN WORLD-WIDE ANCESTRAL FAME,

AND WHO AS LEADERS IN PHYTOGRAPHY DURING MORE THAN A GENERATION'S TIME

WILL BE PRE-EMINENT IN BIOMORPHIC SCIENCE THROUGH ALL AGES,

THESE PAGES

ARE OFFERED AS AN APPRECIATIVE TRIBUTE

FROM A YOUNG COLONY IN ANTIPODAL REMOTENESS.

QK 431
M94

@ 38730

THIS work needs hardly to be introduced by prefatory observations to the notice of professed naturalists, who will recognise its scope and object at a glance. To those however, who may be less conversant in phytographic science, and who may use these pages, some explanatory preface would be needful, in order that they may fully understand the views, which guided the author through the details of this enumerative essay on Australian indigenous plants. Again, to all recipients it may be of some interest, to be informed of the causes, which gave rise to the present literary issue. Of the seven volumes of the "Flora Australiensis," elaborated by Mr. Bentham, mainly through the aid of the Botanic Departments of Kew and Melbourne, the earlier ones were written at a time, when a large portion of Australia remained yet untraversed by any explorers. Since then the territorial occupation as a whole has become nearly doubled, while new lines of geographic research have opened up vast additional tracts of lands for future settlements. Thus, the Flora of our island-continent became also more revealed, with the result of about 850 species of vascular plants having been added already to those described in the "Flora Australiensis"; but, irrespective of this, we have learned in later years much more concerning the regional distribution of the species formerly recorded. All these additional data, accumulated gradually during the last two decennia, though mainly kept together in the volumes of the "Fragmenta Phytographiae Australiae," became necessarily much disconnected and dispersed during the progress of discoveries and the early records connected therewith, so much so, that much difficulty was experienced of late, to command a clear and easy view over these scattered literary fragments. To collect and rearrange them into one or more volumes, conformous with those of the "Flora Australiensis," and supplementary to that work, proved beyond the range of operations of the venerable savant, who spent much of his precious time during fully sixteen years for grandly systematising throughout the native vegetation of these great parts of the British Dominions,—because his unrivalled experience and scientific valour were called more pressingly into channels of research, by which not only the realm of plants of Australia, but indeed that of the whole globe, became drawn both comprehensively and connectedly into the cycle of his observations. Perhaps it may fall yet to the share of the writer of the present pages, to furnish supplements to the "Flora Australiensis"; indeed, he has it under contemplation, to add one volume, solely to be devoted to a descriptive enumeration of all the additional species, and to be arranged in consonance with Bentham's views, whereas another volume would be rendered exclusively the vehicle of record of all new localities, from whence any species of the "Flora Australiensis" were obtained, when perhaps also such alterations, as largely augmented material may suggest, could be made to the descriptive text. But to complete the "Flora" in this manner, so far as our present state of knowledge will admit of it, would involve the thorough revision of the whole vast collections now stored in our local Botanical Museum; therefore, the connected re-examination of at least 80,000 specimens, from which the pages of the "Flora" largely, and those of the "Fragmenta" solely, have emanated. Such a task, to be performed with conscientious care, would need continued and close application throughout several years. Under these circumstances it seemed best, to pave the way for subsequent purposes through the issue of a preliminary publication, by which anyhow the use of both works, above referred to, would be facilitated. A postponement of the work for supplementary volumes, thus arising, would bring with it one advantage also, that of gaining time for carrying the augmentation of the Flora still further, as we would meanwhile profit from phytologic searches of travellers and settlers in wide regions of Australia, as yet in reference to vegetation but scantily investigated or altogether unexplored.

By deferring therefore the elaboration of supplemental volumes, we could aim more at future exhaustiveness, whereas the work as left by its illustrious author will meet mainly all present wants, and must remain the foundation for all future systematic study of the vegetation peculiar to this part of the globe. In offering now a separate statistic volume, the writer has availed himself of the opportunity, to place on record independently his views on preferable systematic sequences of orders, wherever occasionally his opinions differed from those enunciated by other phytographers. He furthermore could draw the limits of some genera and many species in accordance with original observations of his own; besides he was able to apply more rigorously the rules for systematic naming in strict application of chronologic data. But to avoid misapprehensions, the author at the very outset endeavours to anticipate objections, which may perhaps be raised against some portion of the nomenclature adopted in these pages. If, for instance, Haller is regarded as the earliest indicator of Cyperaceae, it must be conceded, that his wording of the order was "Cyperi" in 1742, as well shown in Pfeiffer's extensive work. If however in accordance with the more exact limitation of this ordinal group of plants we ascribe it under the name of

Cyperaceae to De Candolle, as defined in 1805, or to St. Hilaire as circumscribed in the same year, or to A. L. De Jussieu, who already in 1789 adopted the same order under the appellation Cyperoideae, then we evidently act unjustly in not recognising prior claims; indeed, so far as wording is concerned, we lose the substance for the shadow, especially as no necessity exists, to strive for etymologic uniformity at the sacrifice of fair priority. To overcome difficulties or ambiguities of these kinds, we could use the preposition "from" (to be rendered *nach* in German and *d'après* in French) for indicating the originator of an order, genus or species, after the quotation of any later accepted authority. Indeed this principle was adopted already for a list of Australian plant-genera, published by the Royal Society of New South Wales in 1881; and had the scope assigned to the present pages admitted of it, the same rule would have been extended to the species so far as applicable.

To accomplish the object of rendering this publication, as a commentary index, one for easy reference, the schematic form appeared the most eligible. But to bring it within the means available for issue, all specific synonymy and any secondary notations had to be abandoned. Indeed for these, in quoting the "Flora" and the "Fragmenta" throughout, all needful indications are afforded. The geographic notes had to be limited to the initial letters, indicative of our main colonial territories, in which respect a plan was followed, precisely the same as that adopted in 1866 for a geographic list of nearly one thousand species of trees, then already known to constitute the very varied arboreous vegetation of Australia (see volume of the Intercolonial Exhibition of 1866, and of the second Paris "Exposition," 1867.) On that occasion also the terms Choripetaleae and Synpetaleae were already used, to distinguish the chief divisions of the Dicotyledoneae, while the sequence of orders, effected then, was also mainly the same as that followed in the present pages, the Apetaleae of Jussieu or Monochlamydeae of De Candolle having been merged into the petaliferous divisions. But the system, built up by these two great masters in botanic science, is in its main features so genuinously natural, that no subsequent research could bring about very material changes, except in the one particular direction above indicated, a design more or less happily carried out from the time of the earlier writings of Brogniart to that of the latest essays of Jean Mueller, because it was felt, that so long as the Monochlamydeae remained isolated and associated with the Gymnospermeae, so long would we have an imperfect natural system. To proceed in chief alterations further than this, would give a less acceptable system. Even among the Thalamiflorae and Calyciflorae, represented in Australia, as placed together by Bentham, we had already not less than 58 genera, which are entirely apetalous or contain species, in which the corolla remains undeveloped, leaving out of consideration numerous absolutely extra-Australian genera wholly or partly apetalous. Again in Haloragis, Myriophyllum, Cotula, Soliva and some other genera the corolla is absent in the non-antheriferous flowers, while even the very commencement of the Candollean arrangement is made with the apetalous genera Clematis, Thalictrum and Anemone, soon followed by the equally well-known Caltha. Furthermore the Euphorbiaceae, which in the train of their alliances must carry with them always the Urticeae, count among their numerous genera more than one third provided with petals. Besides, in Proteaceae the floral envelope may be regarded as homologous to that of the closely allied Loranthaceae, with an absence of a calyx, comparable to the suppression of that organ in Diplolaena, Asterolasia and few other thus far exceptional genera. Again, a firm thread runs uninterruptedly through all the orders of Curvembryonatae or Amyliferae, whether placed in Thalamiflorae or Calyciflorae or Monochlamydeae or even as regards Plumbagineae into Corolliflorae; nor is there any real difficulty of finding for the rest of the Monochlamydeae proper places among naturally allied orders of supposed higher organisation. Nevertheless affinity is variously radial, not altogether uniserial, as beautifully demonstrated already by Linné in his map of ordinal alliances of plants, published by Fabricius and Giseke, or as lucidly exhibited as long ago as the middle of the last century by Bernard de Jussieu in the Royal Garden of Trianon, through arranging the plants in a class-ground, a method adopted in our Botanic Garden here also already in 1857. In using the term Calyceae for the three main-divisions of the Monocotyledoneae, the author wishes it to be understood, that it remains an open question, whether the flowers of two of these divisons are to be regarded as mono- or di-chlamydeous. But if in the latter case a petaleous development is attributed to the flowers, we are then bound by homology to carry the term of petals also to Junceae even and Restiaceae, and a change in the naming of the three great sections would be needful accordingly. In suppressing throughout the epigynous divisions for unison with the perigynous, into which they often pass so gradually, simplification is aimed at, without causing any disarrangement, or without attempting alterations in the great received systems except nominally. Furthermore we should always regard the apocarpic orders as those of the highest development, because the fruit—the final effort of all growth—when multiplying into separate and complete carpic elements, exhibits the most advanced vegetable organisation.

Much could it have been wished, to allot in this systematic census also columns of citation to those of the larger works, which are exclusively devoted to the Australian Flora. But Robert Brown's ever-memorable "Prodromus" records only vascular Acotyledoneae, Monocotyledoneae, some Apetaleae

and a portion of the Synpetaleae, as far as known in 1810. The two important volumes, issued by Lehmann and his coadjutors as "Plantae Preissianae," refer solely to South-West Australian species. Sir Joseph Hooker's grand "Flora Tasmaniae" became necessarily limited to the plants of that island, although prefaced by an extensive general essay.

A future re-edition of this census from augmented material, in ampler paginal form, might afford space also for noting these works. The "Flora Tasmaniae" will replace however the "Flora Australiensis" in the quotation-column of the second part of this census, in which all the "Evasculares" of Australia are to be enumerated, the total of them having risen already to about 3750 recorded species. For, be it understood, that the genial and learned Director of the Royal Botanic Garden of Kew has laid a firm foundation for the system of evascular Acotyledoneae of All-Australia in his Tasmanian Flora, that work being the first, in which any extensive and connectedly elaborated account of Mosses, Lichens, Fungs and Algs of Australia has been presented.

It was beyond the scope of these pages, to extend the statistics here given; otherwise the writer would have gladly assigned geographic columns also to Europe, Asia, Africa, North and South America for indicating the respective range of many Australian species of plants into other regions of the globe. Whoever may be interested in such phytogeographic subjects is referred to an able article, written by Dr. Engler, Director of the Botanic Garden of Kiel, for the volume of 1881 of his "Jahrbücher," or to the print of a discourse, delivered a few months ago before the School of Mines at Ballarat. To sum up some of the results of this statistic volume, tables are appended, setting forth the number of the species in each order as well as the grand totals arrived at, clearly immigrated plants being excluded from these pages throughout. Future phytological explorations are not likely, to add beyond several hundred specific forms, considered in conservative limitations, to those now adduced; but extensive additions are sure to be made yet to the records of regional distribution. Indeed even while this first part of the work went through the press, so many new localities became known, as to render an appendix now already necessary. For these addenda I am largely indebted either in notations or material to the Rev. Dr. Woolls, Professor Tate, the Rev. B. Scortechini and Mr. F. M. Bailey, while the supplemental notes concerning West Australian plants emanated chiefly from gatherings, which we owe to the Hon. J. Forrest, C.M.G. It should however be observed, that the whole of the vast additional collections, which accumulated in our Botanic Museum since the volumes of the "Flora Australiensis" successively appeared, could not receive in all instances timely close and critical attention for the elaboration of the present work, though its author had the advantage of departmental help from Mr. G. Luehmann in preliminary sorting and comparing of much of the supplemental material; and here it is an apt place, to recognise likewise the aid, afforded by Mr. Léon Henry, in writing out primarily the requisite notes from the "Flora" and the "Fragmenta." Nearly one hundred species more might have been taken into account already, for estimating the number of species in various orders; but the samples hitherto secured did not admit of obtaining accurate specific data, though the ordinal position could be recognised. Thus we know, that already some access to Anonaceae, Menispermaceae, Lauraceae, Euphorbiaceae, Rutaceae and a few other orders could be recorded.

The chronologic references for species, here more extensively carried out, than ever before, amount to nearly 10,000, including those for genera, in this first part of the present publication. The bibliographic works of Pritzel, Pfeiffer and Jackson have done excellent services in these particular inquiries. The position, given to several of the orders on this occasion, is to some extent assailable; thus—as well-known—reasons exist for moving Piperaceae to Nymphaeaceae, Aristolochieae and Nepenthaceae to Sarraceniaceae, Viniferae to Araliaceae, Thymeleae to Rhamnaceae, Balanophoreae and Podostemoneae to Halorageae, Droseraceae to Saxifrageae, Rubiaceae to Loganiaceae, Fluviales to Hydrocharideae, Casuarineae to Coniferae, although the two latter are neither anatomically nor morphologically approaching each other, notwithstanding some deceptive external resemblance, in which however Exocarpus also shares, not to speak of a few other genera. But in considerations like these we should not insist in a dogmatic spirit on the full acceptance of hitherto recognised systematic arrangements, particularly as not yet all the forms of the world's vegetation are known, and as at any moment the discovery of a new plant may turn the scale in weighing the affinity of its allies.

In incurring the responsibility of restoring some generic names, which had sunk almost into oblivion, the writer has been guided by the impartial rules of strict priority, feeling assured, that phytographers will become quite as quickly accustomed to changes from Stylidium to Candollea, or Ionidium to Hybanthus, or Chenolea to Bassia, as they became reconciled to the long forgotten appellations of Trema, Hevea, Centipeda, Galeola, Floriscopa, Nunnezharia and others, thereby discarding names, sanctioned by lengthened use and familiar to all of us, with a readiness as surprising as universal. To alterations or restitutions of specific names shall only be alluded here with a few words. Why Lamarck for an Eleusine selected designedly the appellation E. cruciata, or why Schleiden called a Wolffia purposely W. Michelii, can be as easily perceived as the advisability of withdrawing the specific name antarctica from a Vitis, Cymodocea or Dicksonia. Where authors bestowed several names

at the same time on a species, it should be free to those, who effect the reductions, to choose a collective designation for the consolidated species; cases in point are afforded by Euphrasia Brownii, Fimbristylis communis, Pappophorum commune, Danthonia penicillata.

As to the circumscription of species themselves, it will require yet much of assiduous research of future generations for arriving at a full understanding of the diagnostic value, which has to be attached respectively to the thousands of specific and therefore original forms in the Australian and any other vegetation. Of the plants, enumerated as species in the present pages, a considerable number may prove, that they owe their particularity to hybridism. In such cases they should occupy a subordinate position, and should also here, as elsewhere, receive their double parental name;— thus Lasiopetalum Tepperi would require to have its appellation changed into Lasiopetalum Baueri × discolor; but as the instances of ascertained hybridism among our native plants are as yet so very few, it was deemed best to admit at present the very limited number of known bastards under ordinary specific rank.

The geographical limitations in this work coincide with the political boundaries of the colonial territories, except that the tropic of Capricorn eastward to the 138th degree separates what is here called Northern Australia (N.A.), from the South- and West-Australian extratropic possessions. Such geographic segregations are necessarily quite arbitrary, though they serve our present purpose of assigning to each of the colonial divisions of Australia its number of specified plants; the limitation is the same as that adopted in the "Flora Australiensis," and as regards abbreviations also identical with the method of indications, chosen for the list of Australian trees in 1866. Here may it further be remarked, that the plants of the small and isolated insular spot, called Norfolk-Island, have been counted with those of N.S.W., to whose dominions this isle as well as Lord Howe's Island politically belongs. To draw the species into physiographic and regional complexes must be the work of future periods, when climatic and geologic circumstances throughout Australia shall have become more extensively known. The geographic columns in these pages indicate simply the occurrence of plants within any of the colonial areas, but have been extended even to such species, which merely may pass boundary-lines. In this way are noted for W.A. and S.A. such tropical plants, as reach not beyond Shark-Bay or the vicinity of the MacDonnell-Range; in the same manner Victoria is credited with those plants of New South Wales, which barely advance into East-Gippsland, while South-Australia obtains credit for several Tasmanian plants, which are confined to very limited stretches of country, chiefly in the extreme south-east of that province; and again a few of the New England plants are recorded for Queensland, although they pass merely on to the nearest adjacent ranges across the boundary. At the whole however it can be foreseen, that many a species, still accepted as genuine for these lists, will in the course of further and more facilitated research be eliminated from any future editions of this work, though still larger arrays, containing many a novelty, will have to be mustered for systematic rolls. The lines of demarkation between truly indigenous and more recently immigrated plants can no longer in all cases be drawn with precision; but whereas Alchemilla vulgaris and Veronica serpillifolia were found along with several European Carices in untrodden parts of the Australian Alps during the author's earliest explorations, Alchemilla arvensis and Veronica peregrina were at first only noticed near settlements. The occurrence of Arabis glabra, Geum urbanum, Agrimonia Eupatoria, Eupatorium cannabinum, Carpesium cernuum and some others may therefore readily be disputed as indigenous, and some questions concerning the nativity of various of our plants will probably remain for ever involved in doubts. While concluding these introductory remarks, it is incumbent on the writer, to acknowledge in grateful terms the consideration, shown him by the Hon. James Macpherson Grant, M.L.A., the Ministerial Chief of the Department, in allowing this work to be issued under the auspices of the Victorian Government and at departmental expenditure. Encouraged by generous support, the author will cheerfully continue through such time, as may still be allotted for his worldly career, to devote his strength and resources also in future mainly for the furtherance of Australian phytography, on which much of his efforts were concentrated through more than a third of a century; and as these enquiries are largely carried on without worldly gain, he feels free, to claim the co-operation of any educated and high-minded colonists in these great dominions of the British Crown for promoting studies, which cannot fail to be of great industrial advantage at all future time, which are calculated to advance continuously the cause of education in this part of the world also, which must far and wide contribute to intellectual enjoyments, and above all should lead to religious reverence of that Supreme Godly power, so gloriously revealed in nature's wondrous works.

DICOTYLEDONEAE.

Ray, Method. Plant. nova 2 (1682).

CHORIPETALEAE HYPOGYNAE.

F. v. Mueller, native plants of Victoria I, 1 (1879).

RANUNCULACEAE.

A. L. de Jussieu, Rec. de l'Acad. des sc. (1773) from B. de Jussieu (1759).

CLEMATIS, Linné, gen. pl. 163 (1737) from l'Ecluse (1576).

C. aristata, R. Brown in De Candolle, syst. veg. I, 147 (1818) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,6	M.fr.X,2.
C. pubescens, Huegel, enum. pl. Nov. Holl. austr. occ. I (1837)	W.A.	—	—	—	—	—	—	B.fl.I,6	
C. glycinoides, De Candolle, syst. veg. I, 145 (1818)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,7	M.fr.X,2.
C. microphylla, De Candolle, syst. veg. I, 147 (1818)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,7	M.fr.X,2; XI,20,27.
C. Fawcettii, F. v. Mueller, fragm. phytogr. Austr. X, 1 (1876)	—	—	—	—	N.S.W.	—	—	—	M.fr.X,1.

ANEMONE, Tournefort, inst. rei herb. 275, t. 147 (1700) from Hippocrates, Theophrastos and Dioscorides.

A. crassifolia, Hooker, icones pl. t. 257 (1840)	—	—	T.	—	—	—	—	B.fl.I,8	M.fr.X,1.

MYOSURUS, Dillenius, catal. pl. Gies. 106, t. 4 (1719) from Tabernaemontanus (1588). (Myosuros).

M. minimus, Linné, sp. pl. 284 (1753)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,8	M.fr.X,2; XI,27.

RANUNCULUS, Tournefort, inst. 285, t. 149 (1700) from Bock (1552).

R. aquatilis, Dodonæus stirp. hist. pempt. 387 (1583)	—	S.A.	T.	V.	—		—	B.fl.I,10	
R. Millani, F. v. Mueller in Hook. Kew misc. VII, 358 (1855)...	—	—	—	V.	N.S.W.	—	—	B.fl.I,10	M.fr.X,2.
R. anemoneus, F. v. Mueller in Trans. Phil. Soc. Vict. I, 97 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.I,11	
R. Gunnianus, Hooker, Journ. of Bot. I, 245, t. 233 (1834) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.I,11	
R. lappaceus, Smith in Rees's Cyclopædia, XXIX (1815) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,12	M.fr.X,2.
R. Muelleri, Bentham, Fl. Austr. I, 13 (1863)	—	—	—	V.	N.S.W.	—	—	B.fl.I,13	
R. rivularis, Bks. & Sol. in G. Forst. fl. ins. austr. prodr. 90 (1786)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,13	M.fr.X,2.
R. hirtus, Bks. & Sol. in G. Forst. fl. ins. austr. prodr. 90 (1786)	W.A.	—	—	V.	N.S.W.	Q.	—	B.fl.I,13	M.fr.X,2.
R. parviflorus, Linné, sp. pl. sec. ed. 780 (1762)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,14	M.fr.X,2; XI,27.

CALTHA, Ruppius, flora Jenensis 119 (1718) from Bock (1552). (Trollius).

C. introloba, F. v. Mueller in Trans. Phil. Soc. Vict. I, 98 (1855)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,15	M.fr.X,2; XI,136.

NYMPHAEACEAE.

Salisbury in Koenig & Sims's Ann. of Bot. II. 70 (1805).

CABOMBA, Aublet, Hist. pl. Guian. I, 321, t. 124 (1775). (Brasenia, Hydropeltis).

C. peltata, F. v. Mueller, native pl. of Vict. 15 (1874)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,60	M.fr.X,77.

NYMPHAEA, Tournefort, inst. 260, t. 137 8 (1700) from Theophrastos and Dioscorides.

N. stellata, Willdenow, spec. pl. II, 1153 (1799)	—	—	—	—	—	—	N.A.	—	M.fr.II,142; X,77.
N. gigantea, Hooker, Bot. Mag. t. 4647 (1852)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,61	M.fr.II,141; X,78.

NELUMBO, Tournefort, inst. 261 (1700). (Nelumbium).

N. nucifera, Gaertner, de fruct. I, 73 (1788)	—	—	—	—	—	Q.	N.A.	B.fl.I,62	M.fr.X,77.

DILLENIACEAE.

Salisbury, Parad. Lond. I, 73 (1801).

WORMIA, Rottboell, Nye Saml. Vid. Selsk. Skrivt. II, 532 (1783).

W. alata, R. Brown in De Cand. syst. veg. I, 434 (1818)... ...	—	—	—	—	—	Q.	—	B.fl.I,16	M.fr.VIII,124.

HIBBERTIA, Andrews, Bot. Reposit. t. 126 (1800). (Candollea, Pleurandra, Hemistemma, Adrastaea, Ochrolasia, Hemistephus, Huttia.)

H. Banksii, F. v. Mueller, pl. of Vict. I, 14, implied (1860) ...	—	—	—	—	—	Q.	—	B.fl.I,21	M.fr.II,1; XI,92.
H. Brownii, Bentham, Fl. Austr. I, 21 (1863)	—	—	—	—	—	Q.	—	B.fl.I,21	M.fr.II,1.
H. dealbata, F. v. Mueller, pl. of Vict. I, 14, implied (1860) ...	—	—	—	—	—	—	N.A.	B.fl.I,21	M.fr.II,1.
H. candicans, Bentham, Fl. Austr. I, 21 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.I,21	
H. Benthami, H. angustifolia, Benth. fl. Austr. I, 21 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.I,21	M.fr.II,1.
H. Muelleri, Bentham, Fl. Austr. I, 21 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,21	
H. ledifolia, Bentham, Fl. Austr. I, 22 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,22	
H. verrucosa, F. v. Mueller, pl. of Vict. I, 14, implied (1860) ...	W.A.	—	—	—	—	—	—	B.fl.I,22	M.fr.XI,92.
H. spicata, F. v. Mueller, fragm. II, 1 (1860)	W.A.	—	—	—			—	B.fl.I,22	M.fr.II,1; XI,20,92.
H. polystachya, Bentham, Fl. Austr. I, 22 (1863)...	W.A.	—	—	—	—	—	—	B.fl.I,22	M.fr.XI,92.
H. rhadinopoda, F. v. Mueller, fragm. XI, 91 (1880)	W.A.	—	—	—	—	—	—	—	M.fr.XI,91.
H. furfuracea, Bentham, Fl. Austr. I, 23 (1683)	W.A.	—	—	—	—	—	—	B.fl.I,23	M.fr.I.161; XI,92.
H. hypericoides, F. v. Mueller, pl. of Vict. I, 13, implied (1860)	W.A.	—	—	—	—	—	—	B.fl.I,23	
H. microphylla, Steudel in Lehmann pl. Preiss. I, 273 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.I,23	M.fr.I,217.
H. recurvifolia, Bentham, Fl. Austr. I, 24 (1863)...	W.A.	—	—	—	—	—	—	B.fl.I,24	
H. lineata, Steudel in Lehmann pl. Preiss. I, 272 (1844)... ...	W.A.	—	—	—	—	—	—	B.fl.I,24	M.fr.XI,92.
H. acerosa, Bentham, Fl. Austr. I, 24 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,24	M.fr.XI,92.
H. aurea, Steudel in Lehmann pl. Preiss. I, 272 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,25	M.fr.XI,92.
H. crassifolia, Bentham, Fl. Austr. I, 25 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,25	
H. nitida, F. v. Mueller, pl. of Vict. I, 15 (1860)...	—	—	—	—	N.S.W.	—	—	B.fl.I,25	
H. bracteata, Bentham, Fl. Austr. I, 26 (1863)	—	—	—	—	N.S.W.	-	—	B.fl.I,25	
H. densiflora, F. v. Mueller, pl. of Vict. I, 15 (1860)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,26	M.fr.VII,125; XI,92.
H. hirsuta, Bentham, Fl. Austr. 26 (1863)...	—	—	T.	—	—	—	—	B.fl.I,26	
H. synandra, F. v. Mueller, fragm. IV, 151 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,151,
H. stricta, R. Brown in De Candolle, syst. veg. I, 422 (1818) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,27	
H. humifusa, F. v. Mueller, pl. of Vict. I, 16, t. Suppl. I (1860)	—	—	—	V.	—	—	—	B.fl.I,27	

H. Billardieri, F. v. Mueller, pl. of Vict. I, 14 (1860)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,28	M.fr.IV,116.
H. gracilipes, Bentham, Fl. Austr. I, 28 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,28	M.fr.XI,92.
H. acicularis, F. v. Mueller, pl. of Vict. I, 17 (1860)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,29	M.fr.XI,93.
H. mucronata, F. v. Mueller, pl. of Vict. I, 17 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,29	M.fr.XI,92.
H. hermannifolia, De Candolle, syst. veg. I, 431 (1818)	—	—	—	—	N.S.W.	—	—	B.fl.I,30	
H. velutina, R. Brown in Bentham, Fl. Austr. I, 30 (1863) ...	—	—	—	—	—	Q.	—	B.fl.I.30	
H. oblongata, R. Brown in De Candolle, syst. veg. I, 431 (1818)	—	—	—	—	—	—	N.A.	B.fl.I,30	M.fr.XI,94.
H. melhanoides, F. v. Mueller, fragm. IV, 116 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,116.
H. tomentosa, R. Brown in De Candolle, syst. veg. I, 432 (1818)	—	—	—	—	—	—	N.A.	B.fl.I,30	
H. cistifolia, R. Brown in De Candolle, syst. veg. I, 431 (1818)	—	—	—	—	—	—	N.A.	B.fl.I,30	
H. echiifolia, R. Brown in Bentham, Fl. Austr. I. 31 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.I,31	
H. scabra, R. Brown in Bentham, Fl. Austr. I, 31 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.I,31	
H. lepidota, R. Brown in De Candolle, syst. veg. I, 432 (1818)...	—	—	—	—	—	—	N.A.	B.fl.I,31	
H. vestita, Cunningham in Bentham, Fl. Austr. I, 31 (1863) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,31	
H. serpyllifolia, R. Brown in De Cand. syst. veg. I, 430 (1818)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,32	
H. pedunculata, R. Brown in De Cand. syst. veg. I, 430 (1818)	—	—	—	V.	N.S.W.	—	—	B.fl.I,32	
H. ochrolasia, Bentham, Fl. Austr. I, 32 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,32	M.fr.VII,125;XI,93.
H. angustifolia, Salisbury, parad. Lond. 73 (1807)	—	—	T.	V.	—	—	—	B.fl.I,33	
H. fasciculata, R. Brown in De Candolle, syst. veg. I, 428 (1818)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,33	
H. virgata, R. Brown in De Candolle, syst. veg. I, 428 (1818) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,34	M.fr.XI,94.
H. inclusa, Bentham, Fl. Austr. I, 34 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,34	
H. rostellata, Turcz. in Bull. Soc. Mosc. XXII, part III, 8 (1849)	W.A.	—	—	—	—	—	—	B.fl.I,34	M.fr.XI,93.
H. glomerata, Bentham, Fl. Austr. I, 34 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,34	
H. argentea, Steudel in Lehmann, pl. Preiss. I. 268 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.I,35	
H. pilosa, Steudel in Lehmann, pl. Preiss. I, 272 (1844)... ...	W.A.	—	—	—	—	—	—	B.fl.I,35	
H. montana, Steudel in Lehmann, pl. Preiss. I, 270 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.I,35	M.fr.XI,94.
H. linearis, R. Brown in De Candolle, syst. veg. I, 428 (1818) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,36	
H. diffusa, R. Brown in De Candolle, syst. veg. I, 429 (1818) ...	—	—	—	V.	N.S.W.	—	—	B.fl.I,36	
H. saligna, R. Brown in De Candolle, syst. veg. I, 427 (1818) ...	—	—	—	—	N.S.W.	—	—	B.fl.I,37	M.fr.XI,94.
H. volubilis, Andrews in Bot. Reposit. t. 126 (1800)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,37	M.fr.IV,116;VII,125; XI,94.
H. grossularifolia, Salisbury, parad. Lond. t. 73 (1806)	W.A.	—	—	—	—	—	—	B.fl.I,37	M.fr.XI,94.
H. dentata, R. Brown in De Candolle, syst. veg. I, 426 (1818) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,38	M.fr.VII,125;XI,94.
H. Cunninghamii, Aiton in Hook. Bot. Mag. t. 318 (1832) ...	W.A.	—	—	—	—	—	—	B.fl.I,39	M.fr.XI,93.
H. perfoliata, Huegel, enum. pl. Nov. Holl. austr. occ. 3 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,38	M.fr.XI, 93.
H. bracteosa, Turczaninow, in Bull. Soc. Mosc. XXV, 140 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,38	M.fr.XI,94.
H. longifolia, F. v. Mueller, fragm. IV, 115 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,115.
H. oenotheroides, F. v. Mueller, fragm. VII, 37 (1870)	—	—	—	—	—	Q.	—	—	M.fr.VII,37.
H. glaberrima, F. v. Mueller, fragm. III, 1 (1862)	—	S.A.	—	—	—	—	—	B.fl.I,39	M.fr.III,1;XI,94.
H. Mylnei, Bentham, Fl. Austr. I, 39 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,39	
H. lasiopus, Bentham, Fl. Austr. I, 40 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,40	M.fr.XI,94.
H. potentilliflora, F. v. Mueller in Bentham, Fl. Austr. I, 40 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,40	M.fr.XI,94.
H. pungens, Bentham, Fl. Austr. I, 40 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,40	
H. nutans, Bentham, Fl. Austr. I, 40 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,40	
H. leptopus, Bentham, Fl. Austr. I, 41 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,41	
H. stellaris, Endlicher in Hueg. enum. 3 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,41	M.fr.XI,93.
H. obcuneata, Salisbury, parad. Lond. t. 73 (1807)	W.A.	—	—	—	—	—	—	B.fl.I,42	M.fr.XI,94.
H. glomerosa, F. v. M.; Candollea glomerosa, Benth. Fl. Aus. I, 43 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,43	
H. teretifolia, F. v. Mueller, fragm. IV, 117 (1864)	W.A.	—	—	—	—	—	—	B.fl.I,43	M.fr.IV,117;XI,95.
H. desmophylla, F. v. Mueller, fragm. XI, 95 (1880)	W.A.	—	—	—	—	—	—	B.fl.I,43	M.fr.XI,95.
H. helianthemoides, F. v. M.; Candollea helianthemoides, Turcz. in Bull. Mosc. XXII, part III, 8 (1849)	W.A.	—	—	—	—	—	—	B.fl.I,43	
H. depressa, Steudel in Lehm. pl. Preiss. I, 268 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,44	M.fr.XI,95.
H. Huegelii, F. v. Mueller, fragm. XI, 95 (1880)	W.A.	—	—	—	—	—	—	B.fl.I,44	M.fr.XI,95.
H. subvaginata, F. v. Mueller, fragm. XI, 95 (1880)	W.A.	—	—	—	—	—	—	B.fl.I,45	M.fr.XI,95.
H. vaginata, F. v. Mueller, fragm. XI, 96 (1880)...	W.A.	—	—	—	—	—	—	B.fl.I,45	M.fr.XI,96.
H. subexcisa, Steudel in Lehmann, pl. Preiss. I, 269 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.I,46	M.fr.XI,95.
H. squamosa, Turczaninow in Bull. Soc. Mosc. XXII, part III, 9 (1849)	W.A.	—	—	—	—	—	—	B.fl.I,46	
H. uncinata, F. v. M.; Candollea uncinata, Benth. Fl. Aus. I, 46 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,46	
H. salicifolia, F. v. Mueller, fragm. I, 161 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,46	M.fr.I,161;VII,124;XI,94.
H. Huttii, F. v. Mueller, fragm. VII, 123 (1870)	W.A.	—	—	—	—	—	—	B.fl.I,47	M.fr.VII,123;XI,20.
H. Goyderi, F. v. Mueller, fragm. VII, 123 (1870)	—	—	—	—	—	—	N.A.	—	M.fr.VII,123.
TETRACERA, Linné, gen. pl. 345 (1737).									
T. Nordtiana, F. v. Mueller, fragm. V, 1 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,1.
T. Daemeliana, F. v. Mueller, fragm. V, 191 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,191.
T. Wuthiana, F. v. Mueller, fragm. X, 49 (1876)	—	—	—	—	—	Q.	—	—	M.fr.X,49.
DILLENIA, Linné, gen. pl. 162 (1737).									
D. Andreana, F. v. Mueller, fragm. V, 175 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,175.
PACHYNEMA, R. Brown in De Candolle, syst. veg. I, 412 (1818).									
P. junceum, Bentham, Fl. Austr. I, 47 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,47	M.fr.X,124.
P. complanatum, R. Brown in De Cand., syst. veg. I, 412 (1818)	—	—	—	—	—	—	N.A.	B.fl.I,48	M.fr.X,124.
P. dilatatum, Bentham, Fl. Austr. I, 48 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,48	M.fr.X,124.

MAGNOLIACEAE.

J. de St. Hilaire, Expos. fam. nat. II, 74, t. 83-84 (1805).

DRIMYS, R. & G. Forster, char. gen. 83, t. 42 (1776). (Tasmania).									
D. aromatica, F. v. Mueller, pl. of Vict. I, 20 (1860)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,49	M.fr.VII,18.
D. membranea, F. v. Mueller, fragm. V, 175 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,175;VII 18.
D. dipetala, F. v. Mueller, pl. of Vict. I, 21 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,49	M.fr.VII,18.
D. Howeana, F. v. Mueller, fragm. VII, 17 (1870)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,17.

ANONACEAE.

A. L. de Jussieu, Gen. 283 (1789) from B. de Jussieu (1759).

UVARIA, Linné, Amoen. acad. 404 (1747).

U. membranacea, Bentham, Fl. Austr. I, 51 (1863) ... ... — — — — — Q. — B.fl.I,51
U. Goezeana, F. v. Mueller, fragm. VII, 125 (1871) ... ... — — — — — Q. — — M.fr.VII,125.

FITZALANIA, F. v. Mueller, fragm. IV, 33 (1863).

F. heteropetala, F. v. Mueller, fragm. IV, 33 (1863) ... ... — — — — — Q. — B.fl.I,51 M.fr.III,1;VII,126.

CANANGA, Rumphius, herb. Amboin. II, 195, t. 65 (1741).

C. odorata, J. Hooker and Thomson, fl. Indic. 130 (1855) ... — — — — — Q. — — M.fr.VI,1.

ANCANA, F. v. Mueller, fragm. V, 27 (1865).

A. stenopetala, F. v. Mueller, fragm. V, 27 (1865) ... ... — — — — N.S.W. Q. — — M.fr.V,27,189;VII,121.

POLYALTHIA, Blume, Fl. Jav. Anon. 68 (1829).

P. nitidissima, Bentham, Fl. Austr. I, 51 (1863) ... ... — — — — N.S.W. Q. — B.fl.I,51 M.fr.III,2;VII,126.

POPOWIA, Endlicher, Gen. 831 (1839).

P. australis, Bentham, Fl. Austr. I, 52 (1863) ... ... — — — — — — N.A. B.fl.I,52

MELODORUM, Loureiro, Fl. Cochinch. I, 351 (1790).

M. Leichhardtii, Bentham, Fl. Austr. I, 52 (1863) ... ... — — — — N.S.W. Q. — B.fl.I,52 M.fr.III,41,VI,2;VII,126.
M. Uhrii, F. v. Mueller, fragm. VI, 2 (1867) ... ... — — — — — Q. — — M.fr.VI,2.
M. Maccreai, F. v. Mueller, fragm. VI, 176 (1868) ... ... — — — — — Q. — — M.fr.VI,176,250.

MILIUSIA, Leschenault in Mém. Genèv. V, 213 (1832).

Species undetermined ... ... — — — — — Q. — —

SACCOPETALUM, Bennett in Horfs. pl. Jav. rar. 115, t. 35 (1838).

S. Bidwilli, Bentham, Fl. Austr. I, 53 (1863) ... ... — — — — — Q. — B.fl.I,53
S. Brahei, F. v. Mueller, fragm. VIII, 159 (1874) ... ... — — — — — Q. — — M.fr.VIII,159.

EUPOMATIA, R. Brown in App. Flind. voy. II, 597, t. 2 (1814).

E. Bennettii, F. v. Mueller, fragm. I, 45 (1859) ... ... — — — — N.S.W. Q. — B.fl.I,54 M.fr.I,45;VI,177.
E. laurina, R. Brown in App. Flind. voy. II, 597, t. 2 (1814) ... — — — V. N.S.W. Q. — B.fl.I,54 M.fr.I,45;VI,2;VII,126.

MONIMIEAE.

A. L. de Jussieu in Ann. du Mus. XIV, 30 (1809).

DORYPHORA, Endlicher, Gen. 3015 (1837) (Learosa).

D. sassafras, Endlicher, iconogr. gen. pl., t 10 (1838) ... ... — — — — N.S.W. — — B.fl.V,283

ATHEROSPERMA, Labillardière, Nov. Holl. pl. spec. II, 74, t. 224 (1806).

A. moschatum, Labillardière, Nov. Holl. pl. spec. II, 74, t. 224 (1806) — — T. V. N.S.W. — — B.fl.V,284 M.fr.VIII,142;X,106.

DAPHNANDRA, Bentham, Fl. Austr. V, 285 (1870).

D. micrantha, Bentham, Fl. Austr. V, 285 (1870)... ... ... — — — — N.S.W. Q. — B.fl.V,285
D. repandula, F. v. Mueller, fragm. X, 105 (1877) ... ... — — — — — Q. — — M.fr.X,105.

MOLLINEDIA, Ruiz & Pavon, prodr. 83, t. 15 (1794). (Kibara, Wilkiea).

M. Huegeliana, Tulasne in Ann. sc. nat. sér. quatr. III, 45 (1855) — — — — N.S.W. Q. — B.fl.V,286
M. Wardellii, F. v. Mueller, fragm. V, 155 (1866) ... ... — — — — — Q. — B.fl.V,287 M.fr.V,155;VI,252.
M. loxocarya, Bentham, Fl. Austr. V, 287 (1870)... ... ... — — — — — Q. — B.fl.V,287
M. acuminata, F. v. Mueller, fragm. V, 155 (1866) ... ... — — — — — Q. — B.fl.V,287 M.fr.V,155.
M. macrophylla, Tulasne in Ann. sc. nat. sér. quatr. III, 45 (1855) — — — — N.S.W. Q. — B.fl.V,288
M. longipes, Bentham, Fl. Austr. V, 289 (1870) ... ... — — — — — Q. — B.fl.V,289
M. laxiflora, Bentham, Fl. Austr. V, 289 (1870) ... ... — — — — — Q. — B.fl.V,289
M. pubescens, Bentham, Fl. Austr. V, 290 (1870) ... ... — — — — N.S.W. Q. — B.fl.V,290

HEDYCARYA, R. & G. Forster, char. gen. 127, t. 64 (1776).

H. Cunninghami, Tulasne in Archiv. du Mus. hist. nat. VIII, 408 (1855) — — — V. N.S.W. Q. — B.fl.V,291

PALMERIA, F. v. Mueller, fragm. IV, 151 (1864).

P. scandens, F. v. Mueller, fragm. IV, 152 (1864) ... ... — — — — N.S.W. Q. — B.fl.V,291 M.fr.IV,152.

PIPTOCALYX, Oliver in Bentham's Fl. Austr. V, 292 (1870).

P. Moorei, Oliver in Bentham's Fl. Austr. V, 292 (1870)... ... — — — — N.S.W. — — B.fl.V,292 M.fr.X,106.

MYRISTICEAE.

R. Brown, prodr. 399 (1810) from A. L. de Jussieu (1806).

MYRISTICA, Linné, gen. pl. ed. II, 524 (1742).

M. insipida, R. Brown, prodr. 400 (1810) ... ... — — — — — Q. N.A. B.fl.V,281

LAURACEAE.

Ventenat, Tabl. I, 245 (1799).

CRYPTOCARYA, R. Brown, prodr. 402 (1810). (Caryodaphne).

C. Murrayi, F. v. Mueller, fragm. V, 170 (1866) ... ... — — — — — Q. — B.fl.V,295 M.fr.V,170;X,253.
C. Mackinnoniana, F. v. Mueller, fragm. V, 169 (1866) ... ... — — — — — Q. — B.fl.V,296 M.fr.V,169.
C. patentinervis, F. v. Mueller in De Candolle, prodr. XV, 508 (1864) — — — — N.S.W. Q. — B.fl.V,296 M.fr.V,166.
C. obovata, R. Brown, prodr. 402 (1810) ... ... — — — — N.S.W. Q. — B.fl.V,296 M.fr.V,170.
C. glaucescens, R. Brown, prodr. 402 (1810) ... ... — — — — N.S.W. Q. N.A. B.fl.V,297 M.fr.V,165,166.
C. triplinervis, R. Brown, prodr. 402 (1810) ... ... — — — — N.S.W. Q. — B.fl.V,297 M.fr.V,166.
C. cinnamomifolia, Bentham, Fl. Austr. V, 298 (1870) ... ... — — — — — Q. — B.fl.V,298
C. Meissneri, F. v. Mueller, fragm. V, 170 (1866)... ... ... — — — — N.S.W. Q. — B.fl.V,298 M.fr.V,170.
C. australis, Bentham, Fl. Austr. V, 299 (1870) ... ... — — — — N.S.W. Q. — B.fl.V,299 M.fr.V,167.

BEILSCHMIEDIA, Nees in Wallich, pl. Asiat. rarior II, 69 (1831). (Nesodaphne).

B. obtusifolia, Bentham, in B. & H. gen. III, 1521 (1880) ... — — — — N.S.W. Q. — B.fl.V,299 M.fr.V,166.

ENDIANDRA, R. Brown, prodr. 402 (1810)

E. glauca, R. Brown, prodr. 402 (1810)	—	—	—	—	—	Q.	—	B.fl.V,300				
E. hypotephra, F. v. Mueller, fragm. V, 166 (1866)	—	—	—	—	—	Q.	—	B.fl.V,301	M.fr.V,166.			
E. discolor, Bentham, Fl. Austr. V, 301 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,301				
E. Sieberi, Nees, Syst. Laurin. 194 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.V,301	M.fr.V,166.			
E. virens, F. v. Mueller, fragm. II, 90 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,302	M.fr.V,166.			
E. Muelleri, Meissner in De Candolle, prodr. XV, 509 (1864) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,302				
E. pubens, Meissner in De Candolle, prodr. XV, 509 (1864) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,302				

CINNAMOMUM, Burman, fl. zeil. 62 (1737).

C. Tamala, C. G. & T. F. Nees, Syst. Laurin. 56 (1836)	—	—	—	—	—	Q.	—	B.fl.V,303	M.fr.V,165.			

LITSEA, Lamarck, Diction. III, 574 (1789). (Glabraria 1771, Tetranthera, Cylicodaphne, Litsaea).

L. chinensis, Lamarck, Diction. III, 574 (1789)	—	—	—	—	—	Q.	N.A.	B.fl.V,305				
L. Bindoniana, F. v. Mueller, Tetranthera, fragm. V, 167 (1866)	—	—	—	—	—	Q.	—	B.fl.V,305	M.fr.V,167.			
L. Hexanthus, A. L. de Jussieu in Annal. du Mus. VI, 212 (1805)	—	—	—	—	—	Q.	—	B.fl.V,305				
L. reticulata, Bentham in B. & J. H. gen. pl. III, part I, 161 (1880)	—	—	—	—	—	Q.	—	B.fl.V,306				
L. Zeylanica, C. G. & T. F. Nees, in Amoen. Bonn. I, 58t. 5 (1823)	—	—	—	—	—	Q.	—	B.fl.V,307				
L. dealbata, Nees, Syst. Laurin. 630 (1836)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,307				

CASSYTHA, Osbeck in Linné, sp. pl. 35 (1753).

C. nodiflora, Meissner in De Candolle, prodr. XV, 252 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.V,309				
C. glabella, R. Brown, prodr. 404 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,309	M.fr.V,167.			
C. flava, Nees in Lehmann, pl. Preiss. I, 620 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,310				
C. pubescens, R. Brown, prodr. 404 (1810)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,310	M.fr.V,167.			
C. phaeolasia, F. v. Mueller, fragm. V, 167 (1866)	—	—	—	V.	N.S.W.	—	—	B.fl.V,310				
C. paniculata, R. Brown, prodr. 404 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,311				
C. filiformis, Linné, sp. pl. 35 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.V,311	M.fr.V,167.			
C. melantha, R. Brown, prodr. 404 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,311	M.fr.V,167.			
C. micrantha, Meissner in De Candolle, prodr. XV, 256 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.V,312				
C. racemosa, Nees in Lehmann, pl. Preiss. I, 621 (1845)... ...	W.A.	—	—	—	—	Q.	—	B.fl.V,312				
C. pomiformis, Nees in Lehmann, pl. Preiss. I, 620 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,313				

HERNANDIA, Plumier, Gen. 6, t. 40 (1703).

H. peltata, Meissner in De Candolle, prodr. XV, 263 (1864) ...	—	—	—	—	—	Q.	—	B.fl.V,314				
H. bivalvis, Bentham, Fl. Austr. V, 314 (1870)	—	—	—	—	—	Q.	—	B.fl.V,314				

MENISPERMEAE.

A. L. de Jussieu, Gen. 284 (1789).

TINOSPORA, Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 35 (1851).

T. smilacina, Bentham in Journ. Linn. Soc. V, suppl. II, 2 (1861)	—	—	—	—	—	Q.	N.A.	B.fl.I,55	M.fr.IX,82.			
T. Walcottii, F. v. Mueller in Bentham's Fl. Austr. I, 56 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,56				

FAWCETTIA, F. v. Mueller, fragm. X, 93 (1877).

F. tinosporoides, F. v. Mueller, fragm. X, 93 (1877)	—	—	—	—	N.S.W.	—	—	—	M.fr.X,93;XI,136.			

COCCULUS, De Candolle, sys. veg. I, 515 (1818) from C. Bauhin (1623). (Pericampylus, Legnephora).

C. Moorei, F. v. Mueller, fragm. I, 162 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,56	M.fr.I,162;IX,81,197.			

HYPSERPA, Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 36 (1851). (Selwynia).

H. Selwyni, F. v. Mueller, fragm. IX, 82 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IV,153.			

TRISTICHOCALYX, F. v. Mueller, fragm. IV, 27 (1863).

T. pubescens, F. v. Mueller, fragm. IV, 27 (1863)	—	—	—	—	—	Q.	—	B.fl.I,58	M.fr.IV,27.			
T. diffusus, Miers, contrib. to Bot. III, 286 (1871)	—	—	—	—	—	Q.	—	—				

SARCOPETALUM, F. v. Mueller, pl. of Vict. I, 26 (1860).

S. Harveyanum, F. v. Mueller, pl. of Vict. 27 and 221 (1860) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,57	M.fr.IX,83.			

LEICHHARDTIA, F. v. Mueller, fragm. X, 67 (1876).

L. clamboides, F. v. Mueller, fragm. X, 68 (1876)	—	—	—	—	—	Q.	—	—	M.fr.X,68.			

STEPHANIA, Loureiro, Fl. Cochinch. II, 608 (1790).

S. hernandifolia, Walpers, report bot. syst. I, 96 (1842)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.I.57	M.fr.X,94.			

PACHYGONE, Miers in Ann. and Mag. nat. hist. sec. ser. VII, 37 (1851).

P. Hullsii, F. v. Mueller, fragm. IX, 81 (1875)	—	—	—	—	—	Q.	N.A.	—	M.fr.V,147.			

PLEOGYNE, Miers in Taylor's Ann. and Mag. of nat. hist. sec. ser. VII, 37 (1851). (Microclisia).

P. Cunninghami, Miers in Ann. & Mag. of nat. hist. sec. ser. VII, 37 (1851)	—	—	—	—	—	Q.	—	B.fl.I,59	M.fr.IX,82.			

CARRONIA, F. v. Mueller, fragm. IX, 84 (1875).

C. multisepalea, F. v. Mueller, fragm. IX, 171 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,84.			

ADELIOPSIS, Bentham & J. Hooker, Gen. pl. I, 436 (1862).

A. decumbens, Bentham, Fl. Austr. I, 59 (1863)	—	—	—	—	—	Q.	—	B.fl.I,59	M.fr.IX,83.			

PAPAVERACEAE.

A. L. de Jussieu, Gen. 235 (1789) from B. de Jussieu (1759).

PAPAVER, Tournefort, inst. 237, t. 119 (1700).

P. aculeatum, Thunberg, Fl. Capensis 431 (1813)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,63				

CAPPARIDEAE.

Ventenat, Tabl. III, 118 (1799).

CLEOME, Linné, syst. nat. 9 (1735); Linné, gen. 200 (1737). (Polanisia).

C. oxalidea, F. v. Mueller, fragm. I, 69 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.I,90	M.fr.I,69.			
C. tetrandra, Banks in De Candolle prodr. I, 240 (1824)... ...	—	—	—	—	—	Q.	N.A.	B.fl.I,90	M.fr.VII,40;IX,174.			
C. viscosa, Linné, sp. pl. 672 (1753)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,90	M.fr.IX,174.			

ROEPERLA, F. v. Mueller in Hooker's Kew Misc. IX, 15 (1857). (Tetratelia).

R. clemoides, F. v. Mueller in Hooker's Kew Misc. IX, 15 (1857)	—	—	—	—	—	Q.	N.A.	B.fl.I,91	M.fr.V,105;IX,174.

EMBLINGIA, F. v. Mueller, fragm. II, 2 (1860).

E. calceoliflora, F. v. Mueller, fragm. II, 3 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,92	M.fr.II,3;XI,20.

CADABA, Forskael, Fl. Aeg. Arab. 67 (1775).

C. capparoides, De Candolle, prodr. I, 244 (1824)...	—	—	—	—	—	—	—	N.A.	B.fl.I,92	

CAPPARIS, Tournefort, inst. 261, t. 139 (1700) from Theophrastos, Dioscorides and Plinius. (Busbeckea, Busbecquea).

C. umbellata, R. Brown in De Candolle, prodr. I, 247 (1824) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,93	M.fr.V,104;IX,172.
C. lasiantha, R. Brown in De Candolle, prodr. I, 247 (1824) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,94	M.fr.IX,173.
C. quiniflora, De Candolle, prodr. I, 247 (1824)	—	—	—	—	—	Q.	N.A.	B.fl.I,94	M.fr.IX,173.
C. spinosa, Linné, sp. pl. 503 (1753)	W.A.	—	—	—	—	—	N.A.	B.fl.I,94	M.fr.I,143;V,104;IX,173.
C. sarmentosa, Cunningham in Bentham's Fl. Austr. I, 95 (1863)	—	—	—	—	—	Q.	—	B.fl.I,95	M.fr.V,104;IX,173.
C. ornans, F. v. Mueller in Bentham's Fl. Austr. I, 95 (1863) ...	—	—	—	—	—	Q.	—	B.fl.I,95	M.fr.IX,173.
C. nobilis, F. v. Mueller in Bentham's Fl. Austr. I, 95 (1863) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,95	M.fr.I,163;V,104.
C. canescens, Banks in De Candolle, prodr. I, 246 (1824) ...	—	—	—	—	—	Q.	—	B.fl.I,96	M.fr.IX,173.
C. lucida, R. Brown in Bentham's Fl. Austr. I, 96 (1863) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,96	M.fr.I,163;V,104;IX,173.
C. Mitchelli, Lindley in Mitchell's Three Exped. I, 315 (1838)...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,96	M.fr.IX,173.
C. loranthifolia, Lindley in Mitchell's Trop. Austr. 220 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,97	
C. umbonata, Lindley in Mitchell's Trop. Austr. 257 (1848) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,97	M.fr.IX,173.
C. uberiflora, F. v. Mueller, fragm. IX, 172 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,172.
C. Shanesiana, F. v. Mueller, fragm. X, 94 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,94.
C. humistrata, F. v. Mueller, fragm. V, 156 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,156.
C. Thozetiana, F. v. Mueller, fragm. V. 104 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,156.

APOPHYLLUM, F. v. Mueller in Hooker's Kew Misc. IX, 307 (1857).

A. anomalum, F. v. Mueller in Hooker's Kew Misc. IX, 307 (1857)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,97	M.fr.IX,173;XI,27.

CRUCIFERAE.

A. L. de Jussieu, Gen. 237 (1789) from B. de Jussieu (1759).

NASTURTIUM, R. Brown in W. T. Aiton's hortus Kewensis IV, 110 (1812) from Linné, syst. nat. 9 (1735).

N. terrestre, R. Brown in Aiton's hort. Kewens. IV, 110 (1812)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,66	M.fr.XI,60.

BARBARAEA, Beckmann, Lex. botan. 33 (1801). (Barbarea).

B. vulgaris, R. Brown in De Candolle, prodr. I, 140 (1824) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.I,66	M.fr.VIII,142;XI,6,60.

ARABIS, Linné, gen. pl. 198 (1737) from Dalechamps (1554). (Turritis).

A. glabra, Crantz, stirp. Austriac. 36 (1767)	—	—	—	V.	N.S.W.	—	—	B.fl.I,67	

CARDAMINE, Tournefort, inst. 224, t. 109 (1700) from l'Ecluse (1576).

C. stylosa, De Candolle, regn. veg. syst. nat. II, 24 (1821) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,68	M.fr.VII,19;XI,60.
C. dictyosperma, Hooker, Journ. of Bot. I, 246 (1834)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,68	M.fr.VII,19;X,6,60.
C. radicata, J. Hooker in icones pl. t. 882 (1852)	—	—	T.	—	—	—	—	B.fl.I,69	M.fr.XI,6.
C. laciniata, F. v. Mueller in Transac. Phil. Soc. Vict. I, 34 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,69	M.fr.XI,27,60.
C. hirsuta, Linné, sp. pl. 655 (1753)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,70	M.fr.VII,19;XI,60.
C. eustylis, F. v. Mueller in Transact. Vict. Inst. I, 114 (1854)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,71	M.fr.VII,19;XI,60.

ALYSSUM, Tournefort, inst. 217 (1700) from l'Ecluse (1576). (Meniocus, Alysson).

A. minimum, Pallas, Reise Prov. Russ. Reich. III, 746 (1776)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,71	M.fr.XI,27,60.

WILKIA, Scopoli, introd. hist. nat. 317 (1777). (Malcolmia).

W. Africana, F. v. Mueller, native plants of Vict. I, 33 (1879)...	—	—	—	V.	N.S.W.	—	—	—	M.fr.VII,18.

SISYMBRIUM, Tournefort, inst. 225, t. 109 (1700). (Blennodia).

S. filifolium, F. v. Mueller in Transact. Phil. Soc. Vict. I, 34 (1854)	—	S.A.	—	—	N.S.W.	—	—	B.fl.I,73	
S. trisectum, F. v. Mueller in Transact. Vict. Inst. I, 114 (1854)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,74	M.fr.VII,20;XI,60.
S. nasturtioides, F. v. Mueller in Transact. Vict. Inst. I, 115 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,74	
S. Richardsii, F. v. Mueller, fragm. X, 105 (1877)	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,105.
S. eremigenum, F. v. Mueller, fragm. II, 143 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,74	M.fr.II,143.
S. Lucae, F. v. Mueller, fragm. XI, 59 (1879)	—	—	—	V.	N.S.W.	—	—	—	M.fr.XI,59.
S. cardaminoides, F. v. Mueller in Bentham, Fl. Austr. I, 75 (1863)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,75	M.fr.XI,27,59.

ERYSIMUM, Linné, gen. pl. 198 (1737). (Blennodia).

E. curvipes, F. v. Mueller in Linnaea XXV, 368 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,75	M.fr.VII,20.
E. brevipes, F. v. Mueller in Linnaea XXV, 367 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,75	M.VII,20;XI,6.
E. blennodioides, F. v. Mueller in Linnaea XXV, 367 (1852) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,76	M.fr.VII,20;XI,27,60.
E. Cunninghamii, Bentham, Fl. Austr. I, 76 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,76	
E. Blennodia, F. v. Mueller, fragm. X, 78 (1877)	—	S.A.	—	—	N.S.W.	—	—	B.fl.I,76	M.fr.VII,20;X,78.
E. capsellinum, F. v. Mueller, native pl. of Vict. I, 35 (1879) ...	—	—	—	V.	N.S.W.	—	—	B.fl.I,77	

STENOPETALUM, R. Brown in De Candolle, Mém. du Mus. VII, 239 (1821).

S. velutinum, F. v. Mueller, pl. Vict. I, 49 (1860)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,78	M.fr.VII,20;XI,27.
S. lineare, R. Brown in De Candolle, syst. veg. I, 513 (1821) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,78	M.fr.VII,20;XI,6.
S. sphaerocarpum, F. v. Mueller in Trans. Ph. Soc. Vict. I, 35 (1854)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,78	M.fr.VII,20.
S. nutans, F. v. Mueller, fragm. III, 27 (1862)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,79	M.fr.III,27;VII,20;XI,6.
S. robustum, Endlicher in Huegel enum. 4 (1837)	W.A.	S.A.	—	—	—	—	—	B.fl.I,79	M.fr.VII,20;XI,5,60.
S. pedicellare, F. v. Mueller in Benth. Fl. Austr. I, 79 (1863) ...	W.A.	—	—	—	—	—	—	B.fl.I,79	M.fr.VII,20.
S. croceum, Bunge in Lehmann, pl. Preiss. I, 258 (1844) ...	W.A.	—	—	—	—	—	—	—	M.fr.XI,6.

GEOCOCCUS, Drummond and Harvey in Hook. Kew Misc. VII, 52 (1855).

G. pusillus, Drumm. & Harv. in Hook. Kew Misc. VII, 52 (1855)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,80	M.fr.VII,19; XI,6,26.

MENKEA, Lehmann, Ind. sem. hort. bot. Hamb. 8 (1843).

M. australis, Lehmann in Ind. sem. hort. bot. Hamb. 8 (1843)...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,80	M.fr.II,142.
M. draboides, J. Hooker in Benth. Fl. Austr. I, 80 (1863) ...	W.A.	—	—	—	—	—	—	B.fl.I,80	M.fr.II,142;XII,19.
M. sphaerocarpa, F. v. Mueller, fragm. VIII, 223 (1874) ...	—	S.A.	—	—	—	—	—	—	M.fr.VIII,223.

CAPSELLA, Medicus, Pflanzen-Gattungen 85 (1792). (Hutchinsia partly, Thlaspi partly, Microlepidium).

C. elliptica, C. A. Meyer, Verz. der Pfl. v. Caucas. 194 (1831) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,81	M.fr.VII,19;XI.60.
C. antipoda, F. v. Mueller, pl. of Vict. I, 44 (1860)	—	—	T.	V.	—	—	—	B.fl.I,81	
C. pilosula, F. v. Mueller, pl. of Vict. I, 44 (1860)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,82	M.fr.XI,60.
C. Tasmanica, F. v. Mueller, fragm. XI, 26 (1878)	—	—	T.	—	—	—	—	B.fl.I,87	
C. cochlearina, F. v. Mueller, pl. of Vict. I, 51 (1860)	—	S.A.	—	—	N.S.W.	—	—	B.fl.I,88	M.fr.XI,26.
C. Drummondii, F. v. Mueller, fragm. XI, 26 (1878)	W.A.	S.A.	—	—	—	—	-	B.fl.I,88	M.fr.VII,90;XI,26.
C. humistrata, F. v. Mueller, fragm. XI, 25 (1878)	—	S.A.	—	—	N.S.W.		—	—	M.fr.XI,25.

SENEBIERA, De Candolle in Mém. soc. hist. nat. Par. I, 140 (1799).

S. integrifolia, De Candolle in Mém. soc. his. nat. Par. I, 144 (1799)	—	—	—	—	—	Q.	—	B.fl.I,82	

LEPIDIUM, Tournefort, inst. 215, t. 103 (1700) from Diosc. and Plinius. (Iberis partly, Lepia, Monoploca).

L. strongylophyllum, F. v. Mueller in Benth. Fl. Austr. I, 84 (1863)	—	S.A.	—	—	—	Q.	N.A.	B.fl.I,84	M.fr.XI,6.
L. linifolium, Bentham, Fl. Austr. I, 84 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,84	M.fr.VII,20;XI,6.
L. leptopetalum, F. v. Mueller, pl. of Vict. I, 48 (1860)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,84	
L. rotundum, De Candolle, syst. veg. II, 537 (1818)	W.A.	—	—	—	—	—	—	B.fl.I,85	M.fr.VII,20;XI,6.
L. phlebopetalum, F. v. Mueller, pl. of Vict. I, 47 (1860) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,85	M.fr.VII,19;XI,60.
L. pedicellosum, F. v. Mueller, fragm. XI, 27 (1878)	—	—	—	—	—	—	N.A.	—	M.fr.XI,27.
L. monoplocoides, F. v. Mueller in Trans. Phil. Soc. Vict. 35 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,85	M.fr.XI,27.
L. papillosum, F. v. Mueller in Linnaea XXV, 370 (1852) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,86	M.fr.VII,19;XI,6,28.
L. foliosum, Desvaux, Journ. de Bot. III, 164 et 180 (1813) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,86	M.fr.VII,19.
L. ruderale, Linné, sp. pl. 645 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,86	M.fr.VII,19;XI,27,61.

CAKILE, Tournefort, coroll. 49, t. 483 (1700).

C. maritima, Scopoli, Fl. Carniolica, 344 (1760)	—	S.A.	T.	V.	N.S.W.	—	—	—	M.fr.VII,19;XI,6,28,60.

VIOLACEAE.

De Candolle, Fl. Franc. IV, 801 (1805).

VIOLA, Tournefort, inst. 419, t. 236 (1700) from Plinius.

V. betonicifolia, Smith in Rees's Cyclopedia XXXVII (1817) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,99	M.fr.X,82.
V. hederacea, Labillardière, Nov. Holl. pl. spec. I, 66, t. 91 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,99	M.fr.X,82.
V. Cunninghamii, J. Hooker, Fl. N. Zel. I, 16 (1853)	—	—	T.	—	—	—	—	B.fl.I,100	
V. Caleyana, G. Don, gen. syst. dichlam. pl. I, 329 (1831) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.I,100	M.fr.X,82.

HYBANTHUS, Jacquin, Stirp. Amer. hist. 77, t. 75 (1763). (Ionidium, Pigea, Vlamingia).

H. floribundus, F. v. Mueller, nat. pl. of Vict. I, 45 (1879) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,112	M.fr.X,81.
H. Vernonii, F. v. Mueller, nat. pl. of Vict. I, 45 (1879) ...	—	—	—	V.	N.S.W.	—	—	B.fl.I,103	M.fr.X,81.
H. filiformis, F. v. Mueller, native pl. of Vict. I, 44 (1879) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,103	M.fr.X,81.
H. Tatei, F. v. Mueller in Trans. R. Socy. of S. Aust. IV, 102 (1882)	—	S.A.	—	—	—	—	—	—	
H. debilissimus. F. v. Mueller, fragm. XI, 4 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,4.
H. calycinus, F. v. Mueller, fragm. X, 81 (1877)	W.A.	—	—	—	—	—	—	B.fl.I,104	M.fr.XI.4.
H. enncaspermus, F. v. Mueller, fragm. X, 81 (1877)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,101	M.fr.X,81.

HYMENANTHERA, R. Brown in Tuck. Cong. 442 (1818).

H. Banksii, F. v. Mueller, first. gen. report 9 (1853)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,104	M.fr.X,82.

FLACOURTIEAE.

Cl. Richard in Mém. du Mus. I, 366 (1815).

COCHLOSPERMUM, Kunth, Malvac. 6 (1822).

C. Fraseri, Planchon in Hook. Lond. Journ. VI, 307 (1847) ...	—	—	—	—	—	—	N.A.	B.fl.I,106	
C. heteronemum, F. v. Mueller in Hook. Kew. Misc. IX, 15 (1857)	—	—	—	—	—	—	N.A.	B.fl.I,106	M.fr.IX,61.
C. Gillivraei, Bentham, Fl. Austr. I, 106 (1863)	—	—	—	—	—	Q.	—	B.fl.I,106	
C. Gregorii, F. v. Mueller, fragm. I, 71 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.I,106	M.fr.IX,61.

SCOLOPIA, Schreber, Gen. 335 (1789). (Phoberos).

S. Brownii, F. v. Mueller, fragm. III, 11 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,107	M.fr.III,11;IX,61.

XYLOSMA, G. Forster, prodr. 380 (1786).

X. ovata, Bentham, Fl. Austr. I, 108 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,108	M.fr.IX,60.

STREPTOTHAMNUS, F. v. Mueller, fragm. III, 28 (1862).

S. Moorei, F. v. Mueller, fragm. III, 28 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,108	M.fr.III,28.
S. Beckleri, F. v. Mueller, fragm. III, 28 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,108	M.fr.IX,61.

SAMYDACEAE.

J. Gaertner, de Fruct. III, 328 (1805).

CASEARIA, Jacquin, Stirp. Amer. hist. 132, t. 85 (1763).

C. esculenta, Roxburgh, Fl. Ind. II, 422 (1832)	—	—	—	—	—	Q.	—	B.fl.VI,309	
C. Dallachii, F. v. Mueller, fragm. V, 107 (1866)...	—	—	—	—	—	Q.	—	B.fl.III,309	M.fr.VI,252.

HOMALIUM, Jacquin, Stirp. Amer. hist. 170, t. 183 (1763). (Blackwellia).

H. Vitiense, Bentham in Journ. Linn. Soc. IV, 36 (1860) ...	—	—	—	—	—	Q.	—	B.fl.III,310	
H. brachybotrys, F. v. Mueller, fragm. II, 127 (1861)	—	—	—	—	—	Q.	—	B.fl.III,310	M.fr.II,127.

PITTOSPOREAE.

R. Brown in Flinders' voy. 542 (1814).

PITTOSPORUM, Banks in Gaertner, de Fruct. I, 286, t. 59 (1788).

P. rhombifolium, Cunningham in Hook. icon. pl. t. 621 (1844)...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,110	M.fr.XII,4.
P. melanospermum, F. v. Mueller, fragm. I, 70 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,111	M.fr.VI,167;XII,4.
P. undulatum, Andrews, Bot. reposit. 383 (1800)...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,111	M.fr.VI,167;XII,4.
P. revolutum, Aiton, Hort. Kew, sec. ed. II, 27 (1811)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,111	M.fr.VI,167&186;VII,140.
P. venulosum, F. v. Mueller, fragm. VI, 186 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VII,140;XII,4.
P. ferrugineum, Aiton, Hort. Kew, sec. ed. II, 27 (1811) ...	—	—	—	—	—	Q.	—	B.fl.I.112	M.fr.II,78;III,165;VI,167

P. rubiginosum, Cunningham in Ann. of Nat. Hist. IV, 108 (1842) — — — — — Q. — B.fl.I,112 M.fr.VI,167;VII,140;IX,
P. phillyroides, De Candolle, prodr. I, 347 (1824) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.I,112 M.fr.VI,167. [190.
P. bicolor, Hooker, Journ. Bot. I, 249 (1834) ... ... — — T. V. N.S.W. — — B.fl.I,113 M.fr.VIII,142;XII,4.
P. erioloma, C. Moore et F. v. Mueller, fragm. VII, 139 (1871) — — — — N.S.W. — — — M.fr.VII,139.

HYMENOSPORUM, R. Brown in F. v. Mueller, fragm. II, 77 (1860).]140.
H. flavum, F. v. Mueller, fragm. II, 77 (1860) ... ... — — — — N.S.W. Q. — B.fl.I,114 M.fr.II,77;VI,168;VII,

BURSARIA, Cavanilles, icon. et descr. pl. IV, 30, t. 350 (1797).
B. spinosa, Cavanilles, Icon. pl. IV, 30, t. 350 (1797) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.I,115 M.fr,XII,3.

MARIANTHUS, Huegel, enum. pl. Nov. Holl. 8 (1837). (Oncosporum, Rhytidosporum, Calopetalon).
M. procumbens, Bentham, Fl. Austr. I, 117 (1863) ... ... — — T. V. N.S.W. Q. — B.fl.I,117 M.fr.XII,2.
M. microphyllus, Bentham, Fl. Austr. I, 117 (1863) ... ... W.A. — — — — — — B.fl.I,117 M.fr.II,145.
M. villosus, Bentham, Fl. Austr. I, 117 (1863) ... ... W.A. — — — — — — B.fl.I,117
M. parviflorus, F. v. Mueller, fragm. II, 144 (1861) ... ... W.A. — — — — — — B.fl.I,118 M.fr.II,144;XII,2.
M. bignoniaceus, F. v. Mueller in Trans. Phil. Soc. Vict. 6 (1854) — S.A. — V. - — — B.fl.I,118 M.fr.XII,2.
M. Drummondianus, Bentham, Fl. Austr. I, 119 (1863) ... ... W.A. — — — — — — B.fl.I,119 M.fr.XII,2.
M. coeruleo-punctatus, Klotzsch in Link and Otto, icon. pl. 28, t. 12 (1841) ... ... W.A. — — — — — — B.fl.I,119 M.fr.XII,2.
M. candidus, Huegel, enum. pl. Nov. Holl. 8 (1837) ... ... W.A. — — — — — — B.fl.I,120 M.fr.XII,2.
M. erubescens, Putterlick in Endl. nov. stirp. dec. 60 (1839) ... W.A. — — — — — — B.fl.I,120 M.fr.XII,2.
M. ringens, F. v. Mueller, fragm. I, 218 (1859) ... ... W.A. — — — — — — B.fl.I,120 M.fr.I,218;XII,2.
M. pictus, Lindley in Bot. Regist. XXV, App. XXV (1839) — W.A. — — — — — — B.fl.I,121 M.fr.I,217;II,182;XII,2.

CITRIOBATUS, Cunningham, in Loudon's hort. Brit. supp. I, 585 (1832). (Ixiosporum).
C. multiflorus, Cunningham in Loud. hort. Brit. I, 585 (1832) ... — — — — N.S.W. Q. — B.fl.I,121 M.fr.II,77;VI,168;XII,3.
C. pauciflorus, Cunningham in Loud. hort. Brit. I, 585 (1832) ... — — — — — Q. N.A. B.fl.I,122 M.fr.II,76;VI 168;VII,180.

BILLARDIERA, Smith, Sp. Bot. New Holl. I, t. 1 (1793). (Pronaya, Campylanthera).
B. longiflora, Labillardière, Nov. Holl. plant. spec. I, 64, t. 89 (1804) — — T. V. N.S.W. — — B.fl.I,123
B. scandens, Smith, Spec. bot. New Holl. I, t. 1 (1793) ... ... — S.A. T. V. N.S.W. Q. — B.fl.I,123 M.fr.XI,123.
B. coriacea, Bentham, Fl. Austr. I, 124 (1863) ... ... W.A. — — — — — — B.fl.I,124 M.fr.XII,1.
B. cymosa, F. v. Mueller in Transact. Vict. Inst. I, 29 (1855) ... — S.A. — V. N.S.W. — — B.fl.I,124 M.fr.VI,168;VII,140.
B. elegans, F. v. Mueller, pl. of Vict. I, 78 (1860) ... ... W.A. — — — — — — B.fl.I,125 M.fr.XII,2.
B. variifolia, De Candolle, prodr. I, 346 (1824) ... ... W.A. — — — — — — B.fl.I,125 M.fr.
B. Lehmanniana, F. v. Mueller, pl. of Vict. I, 78 (1860) ... W.A. — — — — — — B.fl.I,125 M.fr.XII,2.
B. floribunda, F. v. Mueller in South Sc. Record II, 1 (1882) ... W.A. — — — — — — B.fl.I,120 M.fr.XII,1.

SOLLYA, Lindley, Bot. Regist. XVII, t. 1460 (1831). (Xerosollya.)
S. heterophylla, Lindley, Bot. Regist. XVII, t. 1466 (1831) ... W.A. — — — — — — B.fl.I,126 M.fr.XII,3.
S. parviflora, Turcz. in Bull. Soc. Mosc. XXVII, 361 (1854) ... W.A. — — — — — — B.fl.I,126 M.fr.XII,3.

CHEIRANTHERA, Cunningham, in Brongn. Bot. voy. Coquille, t. 77 (1829).
C. linearis, Cunningham in Bot. Regist. 1719 (1834) ... ... — S.A. — V. N.S.W. Q. — B.fl.I,127 M.fr.I,97;II,180;IX,168.
C. filifolia, Turczaninow in Bull. Soc. Mosc. XXVII, 364 (1854) W.A. — — — — — — B.fl.I,127 M.fr.I,97;II,79;IX,168.
C. parviflora, Bentham, Fl. Austr. I, 128 (1863) ... ... W.A. — — — — — — B.fl.I,128 M.fr.XII,3.

DROSERACEAE.

Salisbury, Parad. Lond. 95 (1809).

ALDROVANDA, Monti in Act. Bonon. II, 404, t. 12 (1747).
A. vesiculosa, Linné, sp. pl. 281 (1753) ... ... — — — — — Q. — — M.fr.VI,104;X,79.

DROSERA, Linné, gen. pl. 89 (1737). (Londera).
D. Indica, Linné, sp. pl. 282 (1753) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.II,456 M.fr.X,79.
D. Adelae, F. v. Mueller, fragm. IV, 154 (1864) ... ... — — — — — Q. — — M.fr.X,80.
D. Arcturi, Hooker, Journ. of Bot. I, 247 (1834) ... ... — — T. V. N.S.W. — — B.fl.II,456
D. glanduligera, Lehmann, pugill. VIII, 37 (1844) ... ... W.A. S.A. — V. N.S.W. — — B.fl.II,457 M.fr.X,80.
D. pygmaea, De Candolle, prodr. I, 317 (1824) ... ... — S.A. T. V. N.S.W. — — B.fl.II,457 M.fr.X,80.
D. platystigma, Lehmann, pugill. VIII, 37 (1844)... ... ... W.A. — — — — — — B.fl.II,457 M.fr.X,80.
D. pulchella, Lehmann, pugill. VIII, 38 (1844) ... ... W.A. — — — — — — B.fl.II,458 M.fr.X,80.
D. leucoblasta, Bentham, Fl. Austr. II, 458 (1864) ... ... W.A. — — — — - — B.fl.II,458 M.fr.X,80.
D. nitidula, Planchon in Ann. des sc. nat. IX, 285 (1848) ... W.A. — — — — — — B.fl.II,458
D. paleacea, R. Brown in De Candolle, prodr. I, 318 (1824) ... W.A. — — — - — — B.fl.II,458
D. parvula, Planchon in Ann. des sc. nat. IX, 287 (1848) ... W.A. — — — — — — B.fl.II,459
D. Burmanni, Vahl, Symbol. III, 50 (1794) ... ... — S.A. — — — Q. N.A. B.fl.II,459 M.fr.X,80.
D. spathulata, Labillardière, Nov. Holl. pl. spec. I, 79, t. 106 (1804) — S.A. T. V. N.S.W. Q. — B.fl.II,459
D. Drummondii, Lehmann, pl. Preiss. II, 235 (1847) ... ... W.A. — — — — — — B.fl.II,460
D. scorpioides, Planchon in Ann. des sc. nat. IX, 288 (1848) ... W.A. — — — — — — B.fl.II,460 M.fr.VIII,185;X,85.
D. petiolaris, R. Brown in De Candolle, prodr. I, 318 (1824) ... — — — — — Q. N.A. B.fl.II,460 M.fr.X,80.
D. binata, Labillardière, Nov. Holl. pl. spec. I, 78, t. 105 (1804) — S.A. T. V. N.S.W. Q. — B.fl.II,461 M.fr.X,80.
D. zonaria, Planchon in Ann. des sc. nat. IX, 303 (1848) ... W.A. — — — — — — B.fl.II,462
D. bulbosa, Hooker, icon. pl. t. 375 (1841)... ... ... W.A. — — — — — — B.fl.II,462
D. rosulata, Lehmann, pugill. VIII, 36 (1844) ... ... W.A. — — — — — — B.fl.II,462 M.fr.X,80.
D. Whittakerii, Planchon in Ann. des sc. nat. IX, 302 (1848) ... — S.A. — V. — — — B.fl.II,462
D. macrophylla, Lindley, Bot. Regist. XXV, App. XX, (1839)... W.A. — — — — — — B.fl.II,463
D. squamosa, Bentham, Fl. Austr. II, 463 (1864) ... ... W.A. — — — — — — B.fl.II,463
D. erythrorrhiza, Lindley, Bot. Regist. XXV, App. XX (1839)... W.A. — — — — — — B.fl.II,463
D. stolonifera, Endlicher in Hueg. enum. 5 (1837) ... ... W.A. — — — — — — B.fl.II,463
D. humilis, Planchon in Ann. des sc. nat. IX, 300 (1848) ... W.A. — — — — — — B.fl.II,464
D. ramellosa, Lehmann, pugill. VIII, 40 (1844) ... ... W.A. — — — - — — B.fl.II,464
D. platypoda, Turczaninow in Bull. del'Ac. de St. Petersb. 343 (1852) W.A. — — — — — — B.fl.II,464 M.fr.VIII,185;X,80.
D. auriculata, Backhouse in Ann. des sc. nat. IX, 295 (1848) ... — S.A. T. V. N.S.W. — — B.fl.II,465 M.fr.X,80.
D. lunata, Buchanan in De Cand. prodr. I, 319 (1824) ... ... — — — — — Q. — — —
D. peltata, Smith in Willd. spec. pl. I, 1546 (1797) ... ... — S.A. T. V. N.S.W. Q. — B.fl.II,465 M.fr.X,80.

D. Neesii, Lehmann, pugill. VIII, 42 (1844)	W.A.	—	—	—				—	B.fl.II,466	
D. gigantea, Lindley, Bot. Regist. XXV, App. XX (1839) ...	W.A.	—	—	—	—	—	—	—	B.fl.II,466	
D. myriantha, Planchon in Ann. des sc. nat. IX, 291 (1848) ...	W.A.	—	—	-	—	—	—	—	B.fl.II,466	
D. pallida, Lindley, Bot. Regist. XXV, App. XX (1839) ...	W.A.	—	-	—	—	—	—	—	B.fl.II,467	
D. penicillaris, Bentham, Fl. Austr. II, 467 (1864)	W.A.	—	—	—	—	—	—	—	B.fl.II,467	
D. filicaulis, Endlicher in Hueg. enum. 6 (1837)	W.A.	—	—	—	—	-	—	—	B.fl.II,467	
D. Huegelii, Endlicher in Hueg. enum. 6 (1837)	W.A.	—	—	-	—	—	—	—	B.fl.II,467	M.fr.VIII,185;X,81.
D. macrantha, Endlicher in Hueg. enum. 6 (1837)	W.A.	—	—	—	—	—	—	—	B.fl.II,468	
D. Menziesii, R. Brown in De Candolle, prodr. I, 319 (1824) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	—	B.fl.II,468	
D. calycina, Planchon in Ann. des sc. nat. IX, 299 (1848) ...	W.A.	—	—	-	-	—	—	—	B.fl.II,468	
D. heterophylla, Lindley, Bot. Regist. XXV, App. XX (1839)...	W.A.			-	—	—	—	—	B.fl.II,469	
D. Banksii, R. Brown in De Candolle, prodr. I, 319 (1824) ...	—	—	—	—	—	Q.	—	B.fl.II,469	
BYBLIS, Salisbury, Parad. Lond. t. 95 (1808).										
B. liniflora, Salisbury, Parad. Lond. t. 95 (1808)	—	—	—	—	-	Q.	N.A.	B.fl.II,470	M.fr.X,81.	
B. gigantea, Lindley Bot. Regist. XXV, App. XXI (1839) ...	W.A.	—	—	-	--	—	—	B.fl.II,470	

ELATINEAE.

Cambessèdes in Mém. du Mus. XVIII, 225 (1829).

ELATINE, Linné, gen. pl. 118 (1737).									
E. Americana, Arnott in Edinb. journ. nat. sc. I, 431 (1830) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,178	M.fr.XI,27.
BERGIA, Linné, mantiss. II, 152 (1771).									
B. ammannioides, Roth, nov. pl. spec. 219 (1821)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.I,180	M.fr.II,147.
B. pedicellaris, F. v. Mueller in Benth. Fl. Austr. I, 180 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,180	M.fr.II,145.
B. perennis, F. v. Mueller in Benth. Fl. Austr. I, 180 (1863) ...	—	—	—	—	—	-	N.A.	B.fl.I,181	M.fr.II,146.

HYPERICINAE.

J. de St. Hilaire, Expos. fam. II, 23, t. 75 (1805).

HYPERICUM, Tournefort, inst. 254, t. 131 (1700), from Dioscorides and Plinius.									
H. Japonicum, Thunberg, Fl. Jap. 295, t. 31 (1784)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,182	

GUTTIFERAE.

A. L. de Jussieu, Gen. 243 (1789).

CALOPHYLLUM, Linné, gen. 154 (1737).									
C. inophyllum, Linné, sp. pl. 513 (1753)	—	—	—	—	—	Q.	—	B.fl.I,183	M.fr.IX,175.
C. tomentosum, Wight, Illustrat. Pl. Ind. Orient. I, 128 (1840)	—	—	—	—	—	Q.	—	—	M.fr.IX,174.

POLYGALEAE.

A. L. de Jussieu, in Ann. du Mus. XIV, 386 (1809).

SALOMONIA, Loureiro, Fl. Cochinch. I, 14 (1790).									
S. oblongifolia, De Candolle, prodr. I, 334 (1824)...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,138	M.fr.XI,4.
POLYGALA, Tournefort, inst. 174, t. 79 (1700).									
P. Sibirica, Linné, sp. pl. 702 (1753)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,139	M.fr.XI,4.
P. leptalea, De Candolle, prodr. I, 325 (1824)	-	—	—	—	—	Q.	N.A.	B.fl.I,139	M.fr.XI,4.
P. eriocephala, F. v. Mueller in Benth. Fl. Austr. I, 139 (1863)	—	—	-	—	—	—	N.A.	B.fl.I,139	
P. persicarifolia, De Candolle, prodr. I, 326 (1818)	—	—	—	—	—	—	N.A.	—	M.fr.XI,4.
P. Chinensis, Linné, sp. pl. 704 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,140	M.fr.XI,4.
P. stenoclada, Bentham, Fl. Austr. I, 141 (1863)...	—	—	—	—	—	—	N.A.	B.fl.I,141	
COMESPERMA, Labillardière, Nov. Holl. pl. spec. II, 21 (1806).									
C. sphaerocarpum, Steetz in Lehm. pl. Preiss. II, 314 (1847) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,143	M.fr.XI,3.
C. scoparium, Steetz in Lehm. pl. Preiss. II, 309 (1847)... ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,143	
C. aphyllum, R. Brown in Benth. Fl. Austr. I, 143 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.I,143	
C. spinosum, F. v. Mueller, fragm. I, 144 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,144	M.fr.I,144.
C. volubile, Labillardière, Nov. Holl. pl. spec. II, 24, t. 163 (1806)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,144	M.fr.XI,3.
C. ciliatum, Steetz in Lehm. pl. Preiss. II, 304 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,144	M.fr.X,3.
C. integerimum, Endlicher in Hueg. enum. (1837)	W.A.	—	—	—	—	—	—	B.fl.I,145	M.fr.XI,3.
C. secundum, Banks in De Candolle, prodr. I, 334 (1824) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,145	
C. Drummondii, Steetz in Lehm. pl. Preiss. II, 301 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.I,145	M.fr.XI,2.
C. praecelsum, F. v. Mueller, fragm. XI, 2 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,2.
C. retusum, Labillardière, Nov. Holl. pl. spec. II, 22, t.160 (1806)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,145	M.fr.XI,2.
C, sylvestre, Lindley in Mitch. Trop. Austr. 342 (1848)	—	—	—	—	—	Q.	—	B.fl.I,146	M.fr.I,49;XI,2.
C. acerosum, Steetz in Lehm. pl. Preiss. II, 299 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,146	M.fr.XI,3.
C. ericinum, De Candolle, prodr. I, 334 (1824)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,146	M fr.I,48;XI,2.
C. viscidulum, F. v. Mueller, fragm. X, 4 (1876)	W.A.	—	—	—	—	—	—	—	M,fr.X,4;XI,2.
C. confertum, Labillardière, Nov. Holl. pl. spec. II, 23, t. 161 (1806)	W.A.	—	—	—	—	—	—	B.fl.I,147	M.fr.XI,3.
C. flavum, De Candolle, prodr. I, 334 (1824)	W.A.	—	—	—	—	—	—	B.fl.I,147	M.fr.XI,3.
C. calymega, Labillardière, Nov. Holl. pl. spec. II, 23, t. 162 (1806)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,147	M.fr.XI,3.
C. rhadinocarpum, F. v. Mueller, fragm. XI, 1 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,1.
C. lanceolatum, R. Brown in Benth. Fl. Austr. I, 148 (1863) ...	W.A.	—	—	—	—	—	—	B.fl.I,148	M.fr.XI,1.
C. defoliatum, F. v. Mueller, pl. of Vict. I, 189 (1862)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,148	M.fr.XI,3.
C. nudiusculum, De Candolle, prodr. I, 334 (1824)	W.A.	—	—	—	—	—	—	B.fl.I,148	M.fr.XI,4.
C. virgatum, Labillardière, Nov. Holl. pl. spec. II, 21 t. 159 (1806)	W.A.	—	—	—	—	—	—	B.fl.I,149	M.fr.XI,2.
C. polygaloides, F. v. Mueller in Trans. Phil. Soc. Vict. I, 7 (1854)	W.A.	S.A.	—	V.	—	—	—	B.fl.I,149	M.fr.XI,4.
XANTHOPHYLLUM, Roxburgh, Pl. Corom. III, 81, t. 284 (1819).									
X. Macintyrii, F. v. Mueller, fragm. V, 57 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,57;XI,4.

TREMANDREAE.

R. Brown in Flind. voy. II, 554 App. (1814).

PLATYTHECA, Steetz in Lehmann, pl. Preiss. I, 220 (1845).

P. galioides, Steetz in Lehm. pl. Preiss. I, 220 (1845)	W.A.	—	—	—	—	—	—	B.fl.I,136	M.fr.XII.

TETRATHECA, Smith, Spec. Bot. New Holl. I, t. 2 (1793).

T. Harperi, F. v. Mueller, fragm. V, 49 (1865)	W.A.	—	—	—	—	—	—	—	M.fr.V,189.
T. ciliata, Lindley in Mitch. three Exped. II, 206 (1838) ...	—	S.A.	T.	V.	—	—	—	B.fl.I,130	M.fr.XII.
T. efoliata, F. v. Mueller, fragm. X, 3 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,3.
T. aphylla, F. v. M. in South. Science Record, vol. II, May(1882)	W.A.	—	—	—	—	—	—	—	M.fr.XII.
T. ericifolia, Smith, Exot. Bot. I, 37, t. 20 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,131	M.fr.XII.
T. juncea, Smith, specim. of Bot. of New Holl. 5, t. 2 (1793) ...	—	—	—	—	N.S.W.	—	—	B.fl.I,132	M.fr.XII.
T. affinis, Endlicher in Hueg. enum. 7 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,132	M.fr.XII.
T. nuda, Lindley, Bot. Regist. XXV, App. XXXVIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.I,133	M.fr.XII.
T. virgata, Steetz in Lehm. pl. Preiss. I, 212 (1845)	W.A.	—	—	—	—	—	—	B.fl.I,133	M.fr.XII.
T. confertifolia, Steetz in Lehm. pl. Preiss. I, 214 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.I,133	M.fr.XII.
T. setigera, Endlicher in Hueg. enum. 8 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,133	M.fr.XII.
T. hirsuta, Lindley, Bot. Regist. XXV, App. XXXVIII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,134	M.fr.XII.
T. pilifera, Lindley, Bot. Regist. XXV, App. XXXVIII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,135	M.fr.XII.
T. filiformis, Bentham, Fl. Austr. I, 135 (1863)	W.A.	—	—	-	—	—	—	B.fl.I,135	M.fr.XII.

TREMANDRA, R. Brown in App. Flind. voy. II, 544 (1814).

T. stelligera, R. Brown in De Candolle, prodr. I, 344 (1824) ...	W.A.	—	—	—	—	—	—	B.fl.I,136	M.fr.XII.
T. diffusa, R. Brown in De Candolle, prodr. I, 344 (1824) ...	W.A.	—	—	—	—	—	—	B.fl.I,137	M.fr.XII.

MELIACEAE.

Ventenat, Tabl. III, 159 (1799).

HEDRAIANTHERA, F. v. Mueller, fragm. V, 58 (1865).

H. porphyropetala, F. v. Mueller, fragm. V, 58 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,58.

TURRAEA, Linné, mantiss. II, 150 (1771).

T. pubescens, Hellenius in K. Sv. Vent. Acad. Handl. 26, t. 10 (1788)—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,379	M.fr.V,144.

MELIA, Linné, gen. pl. 127 (1737).

M. Azedarach, Linné, sp. pl. 384 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,380	M.fr.V,144;IX,134;XI,135.

DYSOXYLUM, Blume, Bijdrag. 172 (1825). (Dysoxylon, Epicharis, Hartigshea).

D. oppositifolium, F. v. Mueller, fragm. V, 144 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,144;IX,61.
D. latifolium, Bentham, Fl. Austr. I, 381 (1863)	—	—	—	—	—	Q.	—	B.fl.I.381	M.fr.IX,61.
D. Klanderi, F. v. Mueller, fragm. V, 176 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,176;IX,61,134.
D. Fraserianum, Bentham, Fl. Austr. I, 381 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,381	M.fr.IX,61.
D. arborescens, Miquel in Annal. Mus. Bot. Lugd. IV, 24 (1868)	—	—	—	—	—	Q.	—	—	M.fr.IX,134.
D. Muelleri, Bentham, Fl. Austr. I, 381 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,381	M.fr.V,145.
D. Schiffneri, F. v. Mueller in Melb. Chemist, 53 (1881) ...	—	—	—	—	—	Q.	—	—	M.fr.XI,138.
D. Becklerianum, C. de Candolle in Monogr. phan. I, 509 (1878)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.X,138.
D. Lessertianum, Bentham, Fl. Austr. I, 382 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,382	M.fr.V,145;IX,61.
D. Schultzii, C. de Candolle in Monogr. phanerogr. I, 509 (1878)	—	—	—	—	—	—	N.A.	—	M.fr.XI,138.
D. rufum, Bentham. Fl. Austr. I, 382 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,382	M.fr.V,145.
D. Nernstii, F. v. Mueller, fragm. V, 176 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,176.

AMOORA, Roxburgh, pl. Corom. III, 54, t. 258 (1819).

A. nitidula, Bentham, Fl. Austr. I, 383 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,383	

SYNOUM, Adr. de Jussieu in Mém. du Mus. XIX, 226, t. 15 (1830).

A. glandulosum, de Jussieu in Mém. du Mus. XIX, 227, t. 15 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,384	M.fr.V,145;IX,134.
A. Muelleri, C. de Candolle in Monogr. phanerogr. I, 593 (1878)	—	—	—	—	—	Q.	—	—	M.fr.V,145;XI,138.

OWENIA, F. v. Mueller in Hook. Kew Misc. IX, 303 (1857).

O. acidula, F. v. Mueller in Hook. Kew Misc. IX, 304 (1857) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,385	M.fr.III,14;IX,134.
O. vernicosa, F. v. Mueller, fragm. III, 15 (1862)	—	—	—	—	—	—	N.A.	B.fl.I,385	M.fr.III,15;IX,134.
O. venosa, F. v. Mueller in Hook. Kew Misc. IX, 304 (1857) ...	—	—	—	—	—	Q.	—	B.fl.I,386	M.fr.III,14;IX,134.
O. cepiodora, F. v. Mueller, fragm. XI, 82 (1880)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,82.
O. reticulata, F. v. Mueller in Hook. Kew Misc. IX, 305 (1857)	W.A.	—	—	—	—	—	N.A.	B.fl.I,386	M.fr.III,13;IX,134.

AGLAIA, Loureiro, Fl. Cochinch. I, 173 (1790), non Allemand (1770). (Milnea, Nemedra).

A. elaeagnoides, Bentham, Fl. Austr. I, 383 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,383	M.fr.V,145;IX,134.

HEARNIA, F. v. Mueller, fragm. V, 55 (1865). (Aglaiopsis).

H. sapindina, F. v. Mueller, fragm. V, 55 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,55.

CARAPA, Aublet, Hist. pl. Guian. II, Suppl. 32, t. 357 (1775). (Xylocarpus).

C. Moluccensis, Lamarck, Encycl. meth. Bot. I, 621 (1783) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,387	M.fr.V,145;IX,134.

CEDRELA, P. Browne, Nat. hist. of Jamaica 158 (1756).

C. australis, F. v. Mueller, fragm. I, 4 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,387	M.fr.IX,133;XI,82.

FLINDERSIA, R. Brown in Flind. voy. II, 595 (1814). (Oxleya, Strzeleckia).

F. australis, R. Brown in Flind. voy. II, 595, t. 1 (1814) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,388	M.fr.I,65;II,174;IX,133.
F. Bourjotiana, F. v. Mueller, fragm. IX, 133 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,133.
F. Schottiana, F. v. Mueller, fragm. III, 25 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,388	M.fr.III,25;VI,143;IX,133.
F. Oxleyana, F. v. Mueller, fragm. I, 65 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,389	M.fr.I,65;III,25;IX,133.
F. Pimenteliana, F. v. Mueller, fragm. IX, 132 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,132.
F. Bennettiana, F. v. Mueller in Benth. Fl. Austr. I, 389 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,389	M.fr.IX.131;XI,135.
F. Ifflaiana, F. v. Mueller, fragm. X, 94 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,94.
F. Brayleyana, F. v. Mueller, fragm. V, 143 (1866)	—	—	—	—	—	Q.	—	—	M.fr.I,143;VI,252.
F. Strzeleckiana, F. v. Mueller, fragm. I, 65 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,389	M.fr.IX,133.

OCHNACEAE.

De Candolle in Ann. du Mus. XVII, 398 (1811).

BRACKENRIDGEA, Asa Gray, Bot. Wilk. Expl. Exped. 361, t. 42 (1854).

B. Australiana, F. v. Mueller, fragm. V, 29 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,29.

RUTACEAE.

A. L. de Jussieu, Gen. 296 (1789).

ZIERIA, Smith in Transact. Linn. Soc. IV, 216 (1798).

Z. laevigata, Smith in Rees's Cyclopaedia, XXXIX (1819) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,304	M.fr.IX,116.
Z. pilosa, Rudge in Transact. Linn. Soc. X, 293, t. 17 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.I,305	M.fr.I,100;IX,116.
Z. obcordata, Cunningham in Field. N.S.Wales, 330 (1625) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,305	M.fr.I,101;IX,116.
Z. cytisoides, Smith in Rees's Cyclopaedia, XXXIX (1819) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,306	M.fr.IX,116.
Z. Smithii, Andrews' Bot. Rep. t. 606 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,306	M.fr.I,100;IX,116.
Z. granulata, C. Moore in Benth. Fl. Austr. I, 307 (1863) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,307	M.fr.IX,116.
Z. veronicea, F. v. Mueller in Trans. phil. Soc. Vict. I, 11 (1854)	—	S.A.	T.	V.	—	—	—	B.fl.I,305	M.fr.IX,116.

BORONIA, Smith, Tracts relat. nat. hist. 285 (1798). (Cyanothamnus).

B. grandisepala, F. v. Mueller, fragm. I, 66 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,311	M.fr.I,66;II,179.
B. artemisifolia, F. v. Mueller, fragm. I. 66 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,311	M.fr.I,66;IX,111.
B. affinis, R. Brown in Benth. Fl. Austr. I, 311 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.I,311	
B. filicifolia, Cunningham in Benth. Fl. Austr. I, 311 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.I,311	
B. alata, Smith in Transact. Linn. Soc. VIII, 283 (1807) ...	W.A.	—	—	—	—	—	—	B.fl.I,312	M.fr.IX,111.
B. algida, F. v. Mueller in Transact. phil. Soc. Vict. I, 100 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.I,312	M.fr.IX,111.
B. Edwardsii, Bentham, Fl. Austr. I, 312 (1863)	—	S.A.	—	—	—	—	—	B.fl.I,312	
B. ternata, Endlicher, nov. stirp. dec. 6 (1839)	W.A.	—	—	—	—	—	—	B.fl.I,312	M.fr.IX,111.
B. ericifolia, Bentham, Fl. Austr. I, 313 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,313	M.fr.IX,111.
B. inconspicua, Bentham, Fl. Austr. I, 313 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,313	
B. eriantha, Lindley in Mitch. Trop. Austr. 298 (1848)	—	—	—	—	—	Q.	—	B.fl.I,313	
B. alulata, Solander in Benth. Fl. Austr. I, 313 (1863)	—	—	—	—	—	Q.	—	B.fl.I,313	
B. ledifolia, J. Gay, monogr. des Lasiopetalées 20 (1821) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,314	M.fr.IX,111.
B. lanceolata, F. v. Mueller, fragm. I, 66 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.I,314	M.fr.I,66;VII,38;IX,112;XI,
B. Bowmanii, F. v. Mueller, fragm. IV, 135 (1864)	—	—	—	—	—	Q.	—	—	M.fr.VII,38. [97.
B. Fraseri, Hooker, Bot. Mag. t. 4052 (1844)	—	—	—	—	N.S.W.	—	—	B.fl.I,315	M.fr.IX,111.
B. mollis, Cunningham in Lindl. Bot. Reg. 47 (1841)	—	—	—	—	N.S.W.	—	—	B.fl.I,315	M.fr.IX,112.
B. megastigma, Nees in Lehm. pl. Preiss. II, 227 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.I,315	M.fr.II,97;IV,112;IX,112.
B. heterophylla, F. v. Mueller, fragm. II, 98 (1861)	W.A.	—	—	—	—	—	—	B.fl.I,315	M.fr.II,98;IX,112.
B. elatior, Bartling in Lehm. pl. Preiss. I, 170 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,316	M.fr.II,98;IX,122;XI,97.
B. tetrandra, Labillardière, Nov. Holl. pl. spec. I, 98, t. 125 (1804)	W.A.	—	—	—	—	—	—	B.fl.I,316	
B. crassifolia, Bartling in Lehm. pl. Preiss. I, 169 (1844) ...	W.A.	—	—	—	-	—	—	B.fl.I,316	M.fr.IX,112.
B. albiflora, R. Brown in Benth. Fl. Austr. I, 317 (1863) ...	W.A.	—	—	-	-	—	—	B.fl.I,317	M.fr.IX,114.
B. lanuginosa, Endlicher in Hueg. enum. 16 (1837)	W.A.	—	—	-	—	—	—	B.fl.I,317	M.fr.IX,114.
B. pulchella, Turczaninow in Bull. Soc. Mosc. XXV, 162 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,318	M.fr.IX,114.
B. gracilipes, F. v. Mueller, fragm. II, 99 (1861)	W.A.	—	—	—	—	—	—	B.fl.I,318	M.fr.II,99;IX,114.
B. microphylla, Sieber in Spreng. syst. cur. post. 148 (1827) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,318	
B. pinnata, Smith, Tracts relat. nat. hist. 290, t. 4 (1798) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,318	M.fr.IX,114.
B. pilosa, Labillardière, Nov. Holl. pl. spec. I, 97, t. 124 (1804)	—	—	T.	V.	—	—	—	B.fl.I,319	
B. spinescens, Bentham, Fl. Austr. I, 319 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,319	
B. baeckeacea, F. v. Mueller, fragm. IX, 112 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,112.
B. coerulescens, F. v. Mueller in Trans. phil. Soc. Vict. 11 (1854)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,320	M.fr.IX,112.
B. defoliata, F. v. Mueller, fragm. IX, 113 (1875)	W.A.	—	—	—	-	—	—	—	M.fr.IX,113.
B. Busselliana, F. v. Mueller, fragm. IX, 113 (1875)	W.A.	—	—	—	-	—	—	—	M.fr.IX,113.
B. subcoerulea, F. v. Mueller, fragm. II, 100 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,320	M.fr.II,100;IX,112&198.
B. polygalifolia, Smith, Tracts relat. nat. hist. 297, t. 7 (1798)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,320	M.fr.IX,114.
B. falcifolia, Cunningham in Lindl. Bot. Reg. 47 (1841) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,322	M.fr.IX,114.
B. penicillata, Bentham, Fl. Austr. I, 322 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,322	
B. crassipes, Bartling in Lehm. pl. Preiss. I, 168 (1844) ...	W.A.	—	-	—	—	—	—	B.fl.I,322	M.fr.IX,116.
B. subsessilis, Bentham, Fl. Austr. I, 322 (1863)	W.A.	—	—	—	.	—	—	B.fl.I,322	
B. capitata, Bentham, Fl. Austr. I, 323 (1863)	W.A.	—	—	—	—	-	—	B.fl.I,323	M.fr.IX,116.
B. nematophylla, F. v. Mueller, fragm. II, 100 (1861)	W.A.	—	—	—	—	—	—	B.fl.I,323	M.fr.II,100.
B. crenulata, Smith in Transact. Linn. Soc. VIII, 284 (1807) ...	W.A.	—	—	—		-	—	B.fl.I,323	M.fr.XI,115&198.
B. serrulata, Smith, Tracts relat. nat. hist. 292 (1798)	—	—	—	—	N.S.W.	-	-	B.fl.I,323	M.fr.IX,115.
B. rhomboidea, Hooker icon. pl. t. 722 (1848)	—	—	T.	—	—	—	—	B.fl.I,324	
B. parviflora, Smith, Tracts relat. nat. hist. 295, t. 5 (1798) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,324	M.fr.IX,115.
B. viminea, Lindley, Bot. Regist. XXV, App. XVII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.I,324	
B. filifolia, F. v. Mueller, fragm. I, 3 (1858)	—	S.A.	—	V.	—	—	-	B.fl.I,325	M.fr.I,31;IX,114.
B. clavellifolia, F. v. Mueller in Trans. phil. Soc. Vict. I. 12 (1854)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,325	M.fr.I,99;IX,114&198.
B. scabra, Lindley, Bot. Regist. XXV, App. XVII (1839) ...	W.A.	—	—	—	—	.	—	B.fl.I,326	
B. thymifolia, Turcz. in Bull. Soc. Mosc. XXV, part II, 165 (1852)	W.A.	—	—	-	—		—	B.fl.I,326	M.fr.I,99;II,99,IX,116.
B. ovata, Lindley, Bot. Regist. 47 (1841)	W.A.	—	—	—	—	—	—	B.fl.I,326	M.fr.IX,116.
B. Barkeriana, F. v. Mueller, fragm. XI, 96 (1880)	—	—	—	—	N.S.W.	-	—	—	M.fr.XI,96.
B. denticulata, Smith in Transact. Linn. Soc. VIII, 284 (1807)	W.A.	—	—	—	—	—	—	B.fl.I,327	M.fr.IX,115&198.
B. Machardiana, F. v. Mueller, fragm. IX, 115 (1875)	W.A.	—	—	—	—	.	—	—	M.fr.IX,115.
B. spathulata, Lindley, Bot. Regist. XXV, App. XVII (1839)...	W.A.	—	—	—	—	—	—	B.fl.I,327	M.fr.IX,115.
B. juncea, Bartling in Lehm. pl. Preiss. I, 166 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,327	M.fr.IX,115.
B. cymosa, Endlicher in Hueg. enum. 16 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,328	M.fr.II,101;IX,116.

ERIOSTEMON, Smith in Transact. Linn. Soc. IX, 221 (1798). (Crowea, Phebalium, Asterolasia, Microcybe, Pleurandropsia, Geleznowia, Urocarpus, Sandfordia, Actinostigma).

E. virgatus, Cunningham in Hooker's Journ. of Bot. II, 417 (1840)	—	—	T.	—	—	—	—	B.fl.I,332	M.fr.IX,111.
E. nodiflorus, Lindley, Bot. Regist. XXV, App. XVII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.I,336	M.fr.IX,110.

E. spicatus, A. Richard, sert. Astrolab. 76, t. 27 (1833) ...	W.A.	—	—	—	—	—	—	B.fl.I,336	M.fr.IX,110.
E. pungens, Lindley in Mitch. Three Exped. II, 156 (1838) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,338	M.fr.IX,110.
E. montanus, F. v. M., pl. of Vict. I, 129 (1860)	—	—	T.	—	—	—	—	B.fl.I,338	M.fr.IX,109.
E. phylicifolius, F. v. M., fragm. I, 105 (1859)	—	—	—	V.	—	—	—	B.fl.I,339	M.fr.I,105.
E. umbellatus, Turcz. in Bull. Soc. Mosc. XXII, part III, 15 (1849)	—	—	—	V.	N.S.W.	—	—	B.fl.I,339	M.fr.I,104;IX,109.
E. Ralstoni, F. v. M., fragm. II, 101, t. 14 (1861)	—	—	—	V.	N.S.W.	—	—	B.fl.I,339	M.fr.III,165;IX,109.
E. Hillebrandi, F. v. M. in Transact. phil. Soc. Vict. I, 10 (1854) ...	—	S.A.	T.	V.	—	—	—	B.fl.I,340	M.fr.I,4;IX,109.
E. lamprophyllus, F. v. M. in Journ. pharm. Soc. Vict. II, 43 (1858)	—	—	—	V.	—	—	—	B.fl.I,340	M.fr.IX,109.
E. elatior, F. v. M., fragm. I, 181 (1859)	—	—	—	—	N.S.W.	—	—	B.fl.I,340	M.fr.IX,109.
E. ambiens, F. v. M., fragm. VI, 166 (1868)	—	—	—	—	N.S.W.	—	—	—	M.fr.VI,166;IX,110.
E. Oldfieldii, F. v. M., fragm. I, 3 (1858)	—	—	T.	—	—	—	—	B.fl.I,340	M.fr.IX,109.
E. rotundifolius, Cunningham in Hueg. enum. 15 (1837)... ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,341	
E. microphyllus, F. v. M. in Transact. phil. Soc. Vict. I, 99 (1854)	—	S.A.	—	—	—	—	—	B.fl.I,341	M.fr.IX,109.
E. phylicoides, F. v. M., fragm. I, 107 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,341	M.fr.IX,107.
E. ozothamnoides, F. v. M., fragm. I, 103 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,342	M.fr.IX,107.
E. Mortoni, F. v. M., fragm. IX, 108 (1875)	—	—	—	—	N.S.W.	—	—	B.fl.I,342	M.fr.IX,108.
E. sediflorus, F. v. M., fragm. I, 102 (1859)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,342	M.fr.I,102.
E. Nottii, F. v. M., fragm. VI, 22 (1867)	—	—	—	—	—	Q.	—	—	M.fr.VI,22.
E. lepidotus, Sprengel, syst. II, 322 (1825)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,343	M.fr.I,104;IX,107.
E. alpinus, F. v. M., fragm. I, 103 (1859)	—	—	—	V.	N.S.W.	—	—	—	M.fr.I,103.
E. tuberculosus, F. v. M., pl. of Vict. I, 130 (1862)	W.A.	—	—	—	—	—	—	B.fl.I,343	M.fr.IX,108.
E. Benthami, F. v. M., fragm. IX, 108 (1875)	W.A.	—	—	—	—	—	—	B.fl.I,343	M.fr.IX,108.
E. Maxwelli, F. v. M., fragm. IX, 108 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,103.
E. filifolius, F. v. M., fragm. IX, 108 (1875)	W.A.	—	—	—	—	—	—	B.fl.I,344	M.fr.IX,108.
E. squameus, Labillardière, Nov. Holl. pl. spec. I, 111, t. 141 (1804)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,344	M.fr.I,104;IX,107.
E. anceps, Sprengel, syst. II, 322 (1825)	W.A.	—	—	—	—	—	—	B.fl.I,345	M.fr.I,103;IX,108.
E. ovatifolius, F. v. M., fragm. I, 103 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,345	M.fr.I,103.
E. bilobus, F. v. M., fragm. I, 102 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,345	M.fr.IX,108.
E. Beckleri, F. v. M., fragm. IX, 109 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,108.
E. capitatus, F. v. M., fragm. I, 116 (1859)	W.A.	S.A.	—	—	—	—	—	B.fl.I,346	M.fr.I,106;IX,107.
E. Gelcznowii, F. v. M., fragm. I, 107 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,347	M.fr.IX,107.
E. correifolius, F. v. M., fragm. I, 105 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,350	M.fr.IX,107.
E. Cunninghami, F. v. M., fragm. IX, 107 (1875)	—	—	—	—	N.S.W.	—	—	B.fl.I,351	M.fr.IX,107.
E. mollis, Asterolasia mollis, Benth. Fl. Austr. I, 351 (1863) ...	—	—	—	—	N.S.W.	—	—	B.fl.I,349	
E. pleurandroides, F. v. M., fragm. I, 106 (1859)	—	—	—	V.	—	—	—	B.fl.I,349	M.fr.IX,107.
E. trymalioides, F. v. M., fragm. I, 106 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,351	M.fr.I,106.
E. Hookeri, F. v. M., fragm. I, 104 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,352	M.fr.VII,22;IX,107.
E. pallidus, Asterolasia pallida, Benth. Fl. Austr. I, 352 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,352	
E. Drummondii, F. v. M., fragm. I, 105 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,352	M.fr.VII,22.
E. grandiflorus, F. v. M., fragm. I, 105 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,353	M.fr.VI,21.
E. Crowei, F. v. Mueller, pl. of Vict. I, 119 (1860)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,329	M.fr.IX,106.
E. Turczaninowii, F. v. Mueller, pl. of Vict. I, 120 (1860) ...	W.A.	—	—	—	—	—	—	B.fl.I,330	M.fr.IX,106.
E. lanceolatus, K. F. Gaertner, de fruct. III, 154, t. 210 (1807)	—	—	—	V.	N.S.W.	—	—	B.fl.I,331	M.fr.IX,110.
E. Banksii, Cunningham in Hueg. enum. 15 (1837)	—	—	—	—	—	Q.	—	B.fl.I,332	
E. trachyphyllus, F. v. M. in Transact. phil. Soc. Vict. I, 99 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.I,332	M.fr.IX,110.
E. myoporoides, De Candolle, prodr. I, 720 (1824)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,333	M.fr.IX,110.
E. hispidulus, Sieber in Spreng. syst. cur. post. 164 (1827) ...	—	—	—	—	N.S.W.	—	—	B.fl.I,333	M.fr.IX,110.
E. buxifolius, Smith in Rees's Cyclop. XIII, 2 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.I,333	M.fr.IX,111.
E. obovalis, Cunningham in Field's N. S. Wales, 331 (1825) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,334	M.fr.IX,110.
E. scaber, Paxton, Mag. of Bot. XIII, 127 (1846)...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,334	M.fr.IX,111.
E. Brucei, F. v. M., fragm. VII, 38 (1869)	W.A.	—	—	—	—	—	—	—	M.fr.IX,110.
E. linearis, Cunningham in Hueg. enum. 16 (1837)	—	S.A.	—	—	N.S.W.	—	—	B.fl.I,334	
E. difformis, Cunningham in Hueg. enum. 15 (1837)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,335	M.fr.IX,110.
E. parvifolius, R. Brown in Benth. Fl. Austr. I, 335 (1863) ...	—	—	—	—	—	Q.	—	B.fl.I,335	
E. ericifolius, Cunningham in Benth. Fl. Austr. I, 335 (1863) ...	—	—	—	—	N.S.W.	—	—	B.fl.I,335	
PHILOTHECA, Rudge in Transact. Linn. Soc. XI, 298, t. 21 (1815). (Drummondita).									
P. australis, Rudge in Transact. Linn. Soc. XI, 298, t. 21 (1815)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,348	M.fr.VII,20,141.
P. ericoides, F. v. M., fragm. I, 107 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,349	M.fr.VII,21.
P. calida, F. v. M., fragm. VII, 21 (1869)	—	—	—	—	—	Q.	—	—	M.fr.VII,38.
CORREA, Smith in Transact. Linn. Soc. IV, 219 (1798). (Didymeria).									
C. aemula, F. v. M., fragm. I, 3 (1859)	—	S.A.	—	V.	—	—	—	B.fl.I,353	M.fr.IX,117.
C. alba, Andrews, Bot. Reposit. t. 18 (1797)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,354	M.fr.IX,117.
C. speciosa, Andrews, Bot. Reposit. t. 653 (1811)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,354	M.fr.IX,117.
C. Lawrenciana, Hooker, Journ. of Bot. I, 254 (1834)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,355	M.fr.VIII,142;IX,117.
C. decumbens, F. v. M. in Transact. phil. Soc. Vict. I, 30 (1854) ...	—	S.A.	—	—	—	—	—	B.fl.I,356	
NEMATOLEPIS, Turczaninow in Bull. Soc. Mosc. XXV, part III, 158 (1852). (Symphyopetalum).									
N. phlebalioides, Turczaninow in Bull. Soc. Mosc. part III, 158 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,356	M.fr.IX,106.
N. Euphemiae, F. v. M., fragm. III, 149 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,345	M.fr.IX,106.
CHORILAENA, Endlicher in Hueg. enum. 17 (1837).									
C. quercifolia, Endlicher in Hueg. enum. 17 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,357	M.fr.IX,106.
C. hirsuta, Bentham, Fl. Austr. I, 357 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,357	M.fr.IX,106.
DIPLOLAENA, R. Brown in Append. Flind. voy. II, 546 (1814).									
D. Dampieri, Desfontaines in Mém. du Mus. 452, t. 20 (1817) ...	W.A.	—	—	—	—	—	—	B.fl.I,358	M.fr.IX,106.
BOSISTOA, F. v. M. in Benth. Fl. Austr. I, 359 (1863).									
B. sapindiformis, F. v. M. in Benth. Fl. Austr. I, 359 (1863) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,359	M.fr.IX,103.
B. euodiformis, F. v. M., fragm. IX, 174 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,174.

ACRADENIA, Kippist in Transact. Linn. Soc. XXI, 207 (1855).

A. Frankliniae, Kippist in Transact Linn. Soc. XXI, 207 (1855)	—	—	T.	—	—	—	—			B.fl.I,323	M.fr.VI,167;IX,103.

EUODIA, R. et G. Forster, char. gen. 13, t. 7 (1776). (Melicope, Medicosma).

E. pentacocca, F. v. M., fragm. III, 41 (1862)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.III,168;IV,175.
E. Cunninghami, F. v. M., fragm. III, 2 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,362	M.fr.I,27;IX,102.
E. Fareana, F. v. M., fragm. IX, 101 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,101.
E. erythrococca, F. v. M., fragm. I, 26 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,360	M.fr.II,103,178;IX,102.
E. contermina, Moore & F. v. M., fragm. VII, 144 (1871) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,144.
E. octandra, F. v. M., fragm. II, 102 (1861)	—	—	—	—	N.S.W.	—	—	B.fl.I,360	M.fr.II,102;IX,102.
E. micrococca, F. v. M., fragm. I, 144 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,361	M.fr.II,103,180;IX,102.
E. xanthoxyloides, F. v. M., fragm. IV, 155 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,155.
E. Bonwickii, F. v. M., fragm. V, 56 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,179.
E. alata, F. v. M., fragm. VII, 142 (1871)	—	—	—	—	—	Q.	—	—	M.fr.VII,142.
E. accedens, Blume, Bijdr. 246 (1825)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.V,4,56;VII,22;IX,102.
E. littoralis, Endlicher, prodr. Fl. Norfolk. 86 (1833)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,102.
E. polybotrya, Moore & F. v. M., fragm. VII, 143 (1871) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,193.
E. vitiflora, F. v. M., fragm. VII, 144 (1871)	—	—	—	—	—	Q.	—	—	M.fr.VII,144.

BROMBYA, F. v. M., fragm. V, 4 (1865).

B. platynema, F. v. M., fragm. V, 4 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,56;IX,102.

PAGETIA, F. v. M., fragm. V, 178 (1866).

P. medicinalis, F. v. M., fragm. V, 178 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,178;IX,103.

BOUCHARDATIA, Baillon in Adans. VII, 350, t. 10 (1867).

B. neurococca, Baillon in Adans. IX, 110 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,360	M.fr.I,28;II,163,178;IX,103.

XANTHOXYLUM, Catesby in Linné, hort. Cliff. 487 (1737). (Zanthoxylum, Blackburnia).

X. brachyacanthum, F. v. M., pl. of Vict. I, 108 (1862) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,363	M.fr.VII,141;IX,104.
X. Blackburnia, Bentham, Fl. Austr. I, 363 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.I,363	
X. parviflorum, Bentham, Fl. Austr. I, 363 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,363	M.fr.VII,141;IX,104.
X. torvum, F. v. M., fragm. VII, 140 (1871)	—	—	—	—	—	Q.	—	—	M.fr.VII,140;IX,104.

GEIJERA, Schott, Rutac. 4, t. 7 (1834). (Coatesia).

G. Muelleri, Bentham, Fl. Austr. I, 364 (1863)	—	—	—	—	—	Q.	—	B.fl.I,364	M.fr.I,26;IX,104.
G. salicifolia, Schott, Rutac. t. 4 (1834)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,364	M.fr.VII,22;IX,105;XI,134.
G. parviflora, Lindley in Mitch. Trop. Austr. 102 (1848) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,364	M.fr.VII,22;XI,105;XI,139.

PLEIOCOCCA, F. v. Mueller, fragm. IX, 117 (1875).

P. Wilcoxiana, F. v. M., fragm. IX, 117 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,117.

ACRONYCHIA, R. & G. Forster, char. gen. 53, t. 27 (1776). (Cyminosma).

A. Baueri, Schott, Rutac. t. 3 (1834)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,366	M.fr.I,26;IX,103.
A. laevis, R. & G. Forster, char. gen. 53, t. 27 (1776)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,366	M.fr.I,27;IX,104.
A. imperforata, F. v. M., fragm. I, 26 (1858)	—	—	—	—	—	Q.	—	B.fl.I,367	M.fr.I,26;IX,104.
A. melicopoides, F. v. M., fragm. V, 3 (1865)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IV,117;VII,145.
A. acidula, F. v. M., fragm. IV, 154 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,154.
A. vestita, F. v. M., fragm. IV, 155 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,155;IX,104.
A. Endlicheri, Schott, fragm. bot. III, t. 2	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,103.
A. tetrandra, F. v. M., fragm. IX, 104 (1875)	—	—	—	—	—	Q.	—	—	M.fr.V,179;IX,104.

HALFORDIA, F. v. Mueller, fragm. V, 43 (1865).

H. drupifera, F. v. M., fragm. V, 43 (1865)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.V,5,189;IX,103.
H. scleroxyla, F. v. M., fragm. VII, 142 (1871)	—	—	—	—	—	Q.	—	—	M.fr.VII,142;IX,103.

GLYCOSMIS, Correa in Ann. du Mus. VI, 384 (1805).

G. pentaphylla, Correa in Annal. du Mus. VI, 384 (1805) ...	—	—	—	—	—	Q.	—	B.fl.I,367	M.fr.IX,105.

MICROMELUM, Blume, Bijdrag. 137 (1825).

M. pubescens, Blume, Bijdrag. 137 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.I,368	M.fr.I,25;IX,105.

MURRAYA, Koenig in Linné, mant. alt. 563 (1771). (Chalcas, 1767).

M. exotica, Koenig in Linné, mant. alt. 563 (1771)	—	—	—	—	—	Q.	—	B.fl.I,369	M.fr.IX,106.
M. crenulata, Oliver in Journ. Linn. Soc. V, suppl. II, 29 (1861)	—	—	—	—	—	Q.	—	B.fl.I,369	

CLAUSENA, Burman, Fl. Indica index & t. 29 (1768).

C. brevistyla, Oliver in Journ. Linn. Soc. V, Suppl. II. 31 (1861)	—	—	—	—	—	Q.	—	B.fl.I,369	M.fr.IX,106.

ATALANTIA, Correa in Ann. du Mus. VI, 383 (1805).

A. glauca, J. Hooker in B. & H. Gen. pl. I, 305 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,370	M.fr.XI,105.
A. recurva, Bentham, Fl. Austr. I, 370 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,370	M.fr.VII,142.

CITRUS, Linné, gen. pl. 230 (1737), from Plinius.

C. Planchonii, F. v. M., Report Intercol. Exhib. 23 (1867) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,371	M.fr.IX,105. [IX,105.
C. Australasica, F. v. M., fragm. I, 26 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,371	M.fr.I,26;II,178;III,167;

PENTACERAS, J. Hooker in B. & H. Gen. pl. I, 298 (1862).

P. australis, J. Hooker in B. & H. Gen. pl. I, 298 (1862) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,365	M.fr.IX,104.

SIMARUBEAE.

De Candolle in Ann. du Mus. XVII, 323 (1811).

AILANTUS, Desfontaines, in Act. Acad. Paris, 265 (1786).

A. imberbiflora, F. v. M., fragm. III, 42 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,373	M.fr.III,42;VI,166.

BRUCEA, J. S. Miller, icon. plantar. t. 25 (1780).

B. Sumatrana, Roxburgh, Fl. Indic. I, 469 (1820)	—	—	—	—	—	—	N.A.	B.fl.I,373	M.fr.IX,166.

HYPTIANDRA, J. Hooker in B. & H. Gen. pl. 294 (1862).

H. Bidwilli, J. Hooker in B. & H. Gen. pl. 294 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,374	M.fr.VI,165.

CADELLIA, F. v. Mueller, fragm. II, 25 (1860).

C. pentastylis, F. v. M., fragm. II, 26, t. 12 (1860)	—	—	—	—	N.S.W.	—	—	B.fl.I,374	M.fr.II,25&171.
C. monostylis, Bentham, Fl. Austr. I, 375 (1863)...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,375	M.fr.IV,166;VIII,34.

SURIANA, Plumier, nov. plant. Amer. gen. 37, t. 40 (1703).

S. maritima, Linné, sp. pl. 284 (1753)	—	—	—	—	—	Q.	—	B.fl.I,375

HARRISONIA, R. Brown in Mém. du Mus. XII, 517 (1825). (Ebelingia).

H. Brownii, A. L. de Juss. in Mém. du Mus. Par. XII, 530, t. 28 (1825)	—	—	—	—	—	—	N.A.	B.fl.I,376

ZYGOPHYLLEAE.

R. Brown in App. Flind. voy. II, 545 (1814).

NITRARIA, Linné, syst. nat. X, 1044 (1759).

N. Schoberi, Linné, sp. pl. ed. sec. 638 (1762)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,291	M.fr.XI,30.

ZYGOPHYLLUM, Linné, syst. nat. 8 (1735), Linné, gen. 126 (1737). (Sarcozygium).

Z. apiculatum, F. v. M. in Linnaea XXV, 373 (1852)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,292	M.fr.XI,29.
Z. glaucescens, F. v. M. pl. of Vict. I, 228 (1862)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,293	M.fr.XI,29.
Z. iodocarpum, F. v. M. in Linnaea XXV, 372 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,293	M.fr.XI,29.
Z. prismatothecum, F. v. M. in Linnaea XXV, 375 (1852) ...	—	S.A.	—	—	—	—	—	B.fl.I,293	
Z. ammophilum, F. v. M. in Linnaea XXV, 376 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	—	M.fr.XI,28.
Z. Billardieri, de Candolle, prodr. I, 705 (1824)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,293	M.fr.XI,29.
Z. fruticulosum, de Candolle, prodr. I, 705 (1824)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,294	M.fr.XI,29.
Z. Howittii, F. v. M., fragm. III, 150 (1863)	—	S.A.	—	—	—	—	—	—	M.fr.III,150;XI,29.

TRIBULUS, L'Obel, Icon. II, 84 (1581), from Theophr. Diosc. & Plinius. (Tribulopsis).

T. terrestris, L'Obel, stirp. icones II, 84 (1581)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,288	M.fr.XI,30.
T. ranunculiflorus, F. v. M., fragm. I, 48 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,288	M.fr.I,48.
T. hystrix, R. Brown in App. Sturt. Exped. 6 (1849)	—	S.A.	—	—	—	—	N.A.	B.fl.I,289	M.fr.XI,30.
T. macrocarpus, F. v. M. in Benth. Fl. Austr. I, 289 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.I,289	
T. platypterus, Bentham, Fl. Austr. I, 289 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,289	
T. hirsutus, Bentham, Fl. Austr. I, 289 (1863) ... — ...	—	—	—	—	—	—	N.A.	B.fl.I,289	
T. minutus, Leichhardt in Benth. Fl. Austr. I, 291 (1863) ...	—	—	—	—	—	Q.	—	B.fl.I,291	
T. Brownii, F. v. M., pl. of Vict. I, 99 (1860)	—	—	—	—	—	Q.	N.A.	B.fl.I,290	M.fr.I,48.
T. bicolor, F. v. M., pl. of Vict. I, 99 (1860)	—	—	—	—	—	—	N.A.	B.fl.I,290	M.fr.I,47.
T. Solandri, F. v. M., pl. of Vict. I, 99 (1860)	—	—	—	—	—	Q.	N.A.	B.fl.I,290	M.fr.I,47; XI,30.

LINEAE.

De Candolle. Théor. élém. bot. 214 (1813).

HUGONIA, Linné, gen. pl. 134 (1737).

H. Jenkinsii, F. v. M., fragm. V, 7 (1865)	—	—	—	—	—	Q.	—	—	M.fr.IX,190.

ERYTHROXYLUM, P. Brown, Hist. pl. of Jamaica 278 (1756), from Theophr. Diosc. & Plinius. (Erythroxylon).

E. australe, F. v. M. in Transact. Vict. Inst. III, 22 (1855) ...	—	—	—	—	—	Q.	—	B.fl.I,284	
E. ellipticum, R. Brown in Benth. Fl. Austr. I, 284 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.I,284	

LINUM, Tournefort, inst. 339, t. 176, (1700), from Theophr. Diosc. & Plinius.

L. marginale, Cunningh. in Hook. Lond. Jour. Bot. VII, 169 (1848)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,283

GERANIACEAE.

A. L. de Jussieu, Gen. 268 (1789), from B. de Jussieu (1759).

GERANIUM, Tournefort, inst. 266, t. 142 (1700), from Diosc. & Plinius.

G. Carolinianum, Linné, sp. pl. 682 (1753)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,296
G. sessiliflorum, Cavanilles, Diss. 198, t. 77 (1790)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,297

ERODIUM, L. Héritier, Geraniologia, t. 1 (1787).

E. cygnorum, Nees in Lehm. pl. Preiss. I, 162 (1844)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,297	M.fr.XI,20.

PELARGONIUM, L'Héritier, Geraniologia, t. 7 (1787).

P. australe, Willdenow, sp. pl. III, 675 (1800)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,298
P. Rodneyanum, Mitchell, three Exped. II, 144 (1838)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,299

OXALIS, Linné, gen. pl. 134 (1737), from Plinius.

O. Magellanica, G. Forster in Comm. Goetting. IX, 33 (1789) ...	—	—	T.	V.	—	—	—	B.fl.I,300
O. corniculata, Linné, sp. pl. 435 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,301

MALVACEAE.

Adanson in Mém. Acad. fr. 224 (1761).

LAVATERA, Tournefort in Act. Acad. Par. 86, t. 3 (1706).

L. plebeja, Sims in Bot. Mag. t. 2269 (1821)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,185	M.fr.IX,130.

MALVASTRUM, Asa Gray in Mém. Amer. Acad. IV, 21 (1849). (Malva partly).

M. spicatum, A. Gray in Mém. Amer. Acad. IV, 21 (1849) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,187	M.fr.VI,170;IX,130.
M. tricuspidatum, A. Gray in Mém. Amer. Acad. IV, 21 (1849)	—	-	—	—	N.S.W.	Q.	—	B.fl.I,187	

PLAGIANTHUS, R. & G. Forster, char. gen. 85, t. 43 (1776). (Asterotrichon, Blepharanthemum, Lawrencia, Halothamnus).

P. sidoides, Hooker, Bot. Mag. t. 3396 (1835)	—	—	T.	—	—	—	—	B.fl.I,188	M.fr.VI,169.
P. pulchellus, A. Gray, Bot. Amer. Expl. Exped. I, 181 (1854)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,189	M.fr.IX,130.
P. spicatus, Bentham in proceed. Linn. Soc. VI, 103 (1862) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,189	M.fr.VI,170;IX,130.
P. glomeratus, Bentham in proceed. Linn. Soc. VI, 103 (1862)...	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,190	M.fr.IX,130.
P. diffusus, Bentham, Fl. Austr. I, 190 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.I,190	
P. microphyllus, F. v. M., fragm. I, 29 (1858)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,190	M.fr.VII,170;IX,13.
P. Berthae, F. v. M., fragm. V, 103 (1866)	W.A.	S.A.	—	—	—	—	—	—	M.fr.V,103.

SIDA, Linné, gen. pl. 205 (1737), from Theophrastos.

S. corrugata, Lindley in Mitch. Three Exped. II, 13 (1838)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,192		
S. intricata, F. v. M. in Transact. phil. Soc. Vict. I, 19 (1860)	W.A.	S.A.	—	—	N.S.W.	—	N.A.	B.fl.I,193		
S. macropoda, F. v. M. in Benth. Fl. Austr. I, 193 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,193		
S. virgata, Hooker in Mitch. Trop. Austr. 361 (1848)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,194		
S. echinocarpa, F. v. M., fragm. XI, 62 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,62.	
S. cryphiopetala, F. v. M., fragm. III, 4 (1862)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.I,194	M.fr.III,4.	
S. petrophila, F. v. M. in Linnaea XXV, 381 (1852)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,194		
S. calychymenia, J. Gay in de Candolle, prodr. I, 462 (1824)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,194		
S. physocalyx, F. v. M., fragm. III, 3 (1862)	—	—	—	—	—	—	N.A.	B.fl.I,195	M.fr.III,3.	
S. subspicata, F. v. M. in Benth. Fl. Austr. I, 195 (1863)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,195		
S. pleiantha, F. v. M. in Benth. Fl. Austr. I, 195 (1863)	—	—	—	—	—	Q.	—	B.fl.I,195	M.fr.VI,169.	
S. spinosa, Linné, sp. pl. 683 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,196		
S. rhombifolia, Linné, sp. pl. 684 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,196		
S. cordifolia, Linné, sp. pl. 684 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,196		
S. platycalyx, F. v. M. in Benth. Fl. Austr. I, 197 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,197		
S. cleisocalyx, F. v. M., fragm. X, 73 (1876)	—	—	—	—	—	—	N.A.	—	M.fr.X,73.	
S. inclusa, Bentham, Fl. Austr. I, 197 (1863)	—	S.A.	—	—	-	—	N.A.	B.fl.I,197	M.fr.IX,181;XI,32.	
S. cardiophylla, F. v. M., fragm. VIII, 242 (1874)	W.A.	S.A.	—	—	—	—	N.A.	—	M.fr.VIII,242.	
S. Hookeriana, Miquel in Lehm. pl. Preiss. I, 242 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,197		
S. lepida, F. v. M., fragm. VI, 168 (1868)	—	—	—	—	—	—	N.A.	—	M.fr.XI,32.	

ABUTILON, Tournefort, inst. 99, t. 25 (1700), from Camerarius (1586).

A. tubulosum, Hooker in Mitch. Trop. Austr. 390 (1848)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,200	M.fr.IX,131.	
A. amplum, Bentham, Fl. Austr. I, 200 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,200		
A. leucopetalum, F. v. M. in Benth. Fl. Austr. I, 200 (1863)	—	S.A.	—	—	N.S.W.	—	N.A.	B.fl.I,200	M.fr.II,12.	
A. Mitchelli, Bentham, Fl. Austr. I, 201 (1863)	—	S.A.	—	—	—	Q.	—	B.fl.I,201		
A. micropetalum, F. v. M. in Benth. Fl. Austr. I, 201 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,201	M.fr.VI,169 S.;IX,131;XI,	
A. cryptopetalum, F. v. M. in Benth. Fl. Austr. I, 201 (1863)	W.A.	S.A.	—	—	N.S.W.	—	N.A.	B.fl.I,201	M.fr.II,11;IX,13.	[32.
A. geranioides, Bentham, Fl. Austr. I, 202 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,202		
A. otocarpum, F. v. M. in Transact. phil. Soc. Vict. I, 13 (1854)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,202	M.fr.XI,32.	
A. longilobum, F. v. M., fragm. IX, 130 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,130.	
A. subviscosum, Bentham, Fl. Austr. I, 202 (1863)	—	—	—	—	—	Q.	—	B.fl.I,202		
A. Indicum, G. Don, gen. syst. I, 504 (1831)	—	—	—	—	—	Q.	N.A.	B.fl.I,202		
A. auritum, G. Don, gen. syst. I, 500 (1831)	—	—	—	—	—	Q.	—	B.fl.I,203	M.fr.XI,32.	
A. Avicennae, Gaertner, de fruct. II, 251, t. 135 (1791)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,203		
A. graveolens, Wight & Arnott, Prodr. Fl. Pen. Ind. or. I, 56 (1840)	—	—	—	-	—	Q.	—	B.fl.I,204		
A. oxycarpum, F. v. M. in Benth. Fl. Austr. I, 204 (1863)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,204	M.fr.II,12;IX,131;XI,32.	
A. muticum, G. Don, gen. syst. I, 502 (1831)	—	—	—	—	—	Q.	—	B.fl.I,204		
A. Cunninghamii, Bentham, Fl. Austr. I, 205 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.I,205		
A. exonemum, F. v. M., fragm. XI, 63 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,63.	
A. Fraseri, Hooker in Walpers, annal. II, 158 (1851)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,205		
A. halophilum, F. v. M. in Linnaea XXV, 381 (1852)	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.I,206	M.fr.XI,27.	
A. macrum, F. v. M., fragm. IX, 59 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,59.	
A. crispum, G. Don, gen. syst. I, 502 (1831)	—	—	—	—	—	—	N.A.	B.fl.I,206	M.fr.VI,169.	

URENA, Dillenius, Hort. Eltham. 430, t. 319 (1732).

U. lobata, Linné, sp. pl. 692 (1753)	—	—	—	—	—	Q.	—	B.fl.I,206	M.fr.VI,169;IX,130.	
U. Armitiana, F. v. M., fragm. X, 78 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,78.	

PAVONIA, Cavanilles, monadelph. dissert. II, App. II (1786). (Greevesia).

P. hastata, Cavanilles, dissert. III, 138, t. 47 (1786)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,207	M.fr.IX,130.	

HOWITTIA, F. v. Mueller in Transact. Vict. Inst. I, 116 (1855).

H. trilocularis, F. v. M. in Transact. Vict. Inst. I, 116 (1855)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,198	M.fr.VI,169.	

HIBISCUS, Linné, syst. nat. (1735); Linné, gen. pl. 207 (1737). (Fugosia partly, Abelmoschus, Paritium).

H. ficulneus, Linné, sp. pl. 605 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,209	M.fr.I,67.	
H. Abelmoschus, Linné, sp. pl. 696 (1753)	—	—	—	—	—	—	N.A.	—	M.fr.VI,169.	
H. rhodopetalus, F. v. M. in Benth. Fl. Austr. I, 209 (1863)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,209	M.fr.IX,229;XI,31.	
H. Manihot, Linné, sp. pl. 696 (1753)	—	—	—	—	—	Q.	—	B.fl.I,210		
H. Notho-Manihot, F. v. M., fragm. V, 57 (1865)	—	—	—	—	—	Q.	—	—	M.fr.IX,130.	
H. Trionum, Linné, sp. pl. 697 (1753)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,210	M.fr.II,115;IX,129;XI,31.	
H. brachysiphonius, F. v. M., fragm. I, 67 (1858)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,210	M.fr.I,67;VI,169.	
H. Drummondii, Turczan. in Bull. Soc. Mosc. XXXI, 195 (1858)	W.A.	—	—	—	—	—	—	B.fl.I,211	M.fr.I,220;IX,129;XI,31.	
H. microlaenus, F. v. M., fragm. II, 116 (1861)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,211	M.fr.III,5;IX,129.	
H. Pinonianus, Gaudichaud in Freyc. voy. Bot. 476, t. 100 (1826)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,211	M.fr.II,116.	[IX,129;XI,31.
H. cannabinus, Linné, sp. pl. ed. sec., 979 (1763)	—	—	—	—	—	Q.	N.A.	B.fl.I,212	M.fr.II,117;III,166;VI,269;	
H. divaricatus, Graham in Edinb. phil. Journ. (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,212	M.fr.II,118;IX,129;XI,31.	
H. Fitzgeraldii, F. v. M., fragm. VIII, 242 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,242.	
H. Elsworthii, F. v. M., fragm. VIII, 241 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,241.	
H. heterophyllus, Ventenat, jard. de la Malm., t. 103 (1804)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,212	M.fr.VI,170;XI,31.	
H. diversifolius, N. Jacquin, icon. pl. rar. 551 (1789)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,213	M.fr.II,117;VI,169.	
H. splendens, Fraser in Edinb. phil. journ. (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,213	M.fr.VI,170;XI,31&123.	
H. zonatus, F. v. M., fragm. I, 221 (1859)	—	—	—	—	—	—	N.A.	B.fl.I,213	M.fr.VI,170.	
H. Coatesii, F. v. M., fragm. III, 5 (1862)	—	-	—	—	—	—	N.A.	B.fl.I,214	M.fr.III,5.	
H. leptocladus, Bentham, Fl. Austr. I, 214 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,214	M.fr.VI,169;IX,129.	
H. setulosus, F. v. M., fragm. I, 221 (1859)	—	—	—	—	—	—	N.A.	B.fl.I,214	M.fr.I,221;XI,31.	
H. Goldsworthii, F. v. M., fragm. XI, 30 (1878)	-	—	—	—	—	—	N.A.	—	M.fr.XI,30.	
H. pentaphyllus, F. v. M., fragm. II, 13 (1860)	—	—	-	—	—	—	N.A.	B.fl.I,214	M.fr.VI,169;XI,32.	
H. geranioides, Cunningham in Benth. Fl. Austr. I, 215 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,215		
H. vitifolius, Linné, sp. pl. 696 (1753)	—	—	—	—	—	Q.	—	B.fl.I,215	M.fr.II,114;VI,169.	
H. panduriformis, N. Burman, Fl. Ind. 151, t. 47 (1768)	—	—	—	—	—	—	N.A.	B.fl.I,215	M.fr.II,115;VI,169;IX,129.	
H. Normani, F. v. M., fragm. III, 4 (1862)	-	—	-	-	-	Q.	—	B.fl.I,216	M.fr.III,4;VI,169.	

H. Krichauffii, F. v. M., Rep. Babb. Exped. 7 (1858) ... ... — S.A. — — N.S.W. Q. — B.fl.I,216 M.fr.VI,169;IX,129.
H. Farragei, F. v. M., fragm. VIII, 241 (1874) ... ... W.A. S.A. — — N.S.W. — — — M.fr.IX,129&196;XI,32.
H. Sturtii, Hooker in Mitch. trop. Austr. 363 (1848) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.I,216 M.fr.II,13.
H. phyllochlaenus, F. v. M., fragm. IX, 128 (1875) ... ... — — — — — Q. — — M.fr.IX,128.
H. Huegelii, Endlicher in Hueg. enum. 10 (1837)... ... ... W.A. S.A. — — — — — B.fl.I,217 M.fr.IX,128.
H. Wrayae, Lindley, bot. Regist. t. 69 (1840) ... ... W.A. S.A. — — — — — — M.fr.IX,128.
H. tiliaceus, Linné, sp. pl. 694 (1753) ... ... — — — — N.S.W. Q. N.A. B.fl.I,218 M.fr.VI,170;IX,129;XI,32.
H. cuneiformis, de Candolle, prodr. I, 454 (1824)... ... ... W.A. — — — — — N.A. B.fl.I,219 M.fr.XI,32.
H. hakeaefolius, Giordano, Mém. di nuov. sp. d'Ibisco (1833) ... W.A. S.A. — — — — — B.fl.I,220 M.fr.IX,128.

LAGUNARIA, G. Don, gen. syst. dichlam. pl. I, 485 (1831).
L. Patersoni, G. Don, gen. syst. I, 485 (1831) ... ... — — — — N.S.W. Q. — B.fl.I,218 M.fr.IX,130;XI,32.

THESPESIA, Solander in Ann. du Mus. IX, 290 (1807).
T. populnea, Solander in Ann. du Mus. IX, 290, t. 8 (1807) ... — — — — — Q. N.A. B.fl.I,221 M.fr.VII,170;IX,130;XI,32.

GOSSYPIUM, Linné, gen. pl. 206 (1737), from Camerarius (1586). (Fugosia partly, Sturtia). [IX,127;XI,63.
G. Sturtii, F. v. M., fragm. III, 6 (1862) ... ... — S.A. — — N.S.W. — — B.fl.I,222 M.fr.III,174; VI,169 & 251;
G. australe, F. v. M., fragm. I, 46 (1858) ... ... — S.A. — — — Q. N.A. B.fl.I,220 M.fr.III,6;IX,127,XI,32&63
G. thespesioides, F. v. M., fragm. IX, 127 (1875) ... ... — — — — — — N.A. B.fl.I,220 M.fr.V.177;VI,169;XI,32&63
G. Cunninghamii, Todaro, Monogr. Gen. Gossyp. 7 (1878) ... — — — — — — N.A. B.fl.I,220 M.fr.VI,169;IX,129;XI,32&
G. costulatum, Todaro, Monogr. Gen. Gossyp. 7 (1878) ... ... — — — — — — N.A. B.fl.I,221 M.fr.IX,127;XI,32&63. [63.
G. flaviflorum, F. v. M., fragm. IX, 127 (1875) ... ... — — — — — — N.A. — M.fr.XI,63.
G. populifolium, F. v. M., fragm. IX, 127 (1875)... ... ... — — — — — — N.A. B.fl.I,221 M.fr.XI,32&63.
G. Robinsoni, F. v. M., fragm. IX, 126 (1875) ... ... W.A. — — — — — N.A. — M.fr.XI,63.

ADANSONIA, Linné, sp. pl. I, 1190 (1753).
A. Gregorii, F. v. M. in Hook. Kew misc. IX, 14 (1857) ... — — — — — — N.A. B.fl.I,223 M.fr.VII,40;XI,32.

CAMPTOSTEMON, Masters in J. Hook. icon. pl. XII, 18, t. 1119 (1876).
C. Schultzii, Masters in J. Hook. icon. pl. XII, 18, t. 1119 (1876) — — — — — — N.A. — M.fr.XI,32.

BOMBAX, Linné, sp. pl. I, 511 (1753).
B. Malabarica, de Candolle, prodr. I, 479 (1824) ... ... — — — — — Q. N.A. B.fl.I,223 M.fr.XI,32.

STERCULIACEAE.

Ventenat, Jard. de la Malm. II, 91 (1804).

STERCULIA, Linné, Fl. Zeyl. 166 (1747).
S. quadrifida, R. Brown in Benn. pl. Jav. rar. 233 (1852) ... — — — — N.S.W. Q. N.A. B.fl.I,227 M.fr.VI,173.
S. laurifolia, F. v. M. fragm. VI, 172 (1868) ... ... — — — — — Q. — — M.fr.VI,172.

BRACHYCHITON, Schott & Endlicher, Melat. bot. 34 (1832). (Trichosiphon, Poecilodermis, Delabechea).
B. paradoxum, Schott, Melatem. 34 (1832) ... ... — — — — — — N.A. B.fl.I,227
B. Bidwilli, Hooker, Bot. Mag. t. 5133 (1859) ... ... — — — — — Q. — B.fl.I,228
B. discolor, F. v. M., fragm. I, 1 (1858) ... ... — — — — N.S.W. Q. N.A. B.fl.I,228 M.fr.I,242;II,177.
B. incanum, R. Brown in Benn. pl. Jav. rar. 234 (1852)... ... — — — — — — N.A. B.fl.I,228
B. luridum, C. Moore in F. v. M., fragm. I, 1 (1858) ... — — — — N.S.W. Q. — B.fl.I,228 M.fr.I,1;II,177.
B. platanoides, R. Brown in Benn. pl. Jav. rar. 234 (1852) ... — — — — — Q. N.A. B.fl.I,229 M.fr.VI,173.
B. acerifolium, F. v. M., fragm. I, 1 (1858) ... ... — — — — N.S.W. Q. — B.fl.I,229 M.fr.I,1;VI,173.
B. diversifolium, R. Brown in Benn. pl. Jav. rar. 234 (1852) ... W.A. — — — — — N.A. B.fl.I,230
B. populneum, R. Brown in Benn. pl. Jav. rar. 234 (1852) ... — — — V. N.S.W. Q. — B.fl.I,229
B. Delabechii, F. v. M., pl. of Vict. I, 157 (1860)... ... ... — — — — — Q. — B.fl.I,230
B. Gregorii, F. v. M. in Hook. Kew misc. IX, 199 (1857) ... W.A. S.A. — — — — — — M.fr.IX,137.

TARRIETIA, Blume, Bijdr. 227 (1825). (Argyrodendron).
T. argyrodendron, Bentham, Fl. Austr. I, 230 (1863) ... ... — — — — N.S.W. Q. — B.fl.I,230 M.fr.VI,173;IX,42.
T. trifoliolata, F. v. M., fragm. IX, 43 (1875) ... ... — — — — N.S.W. Q. — — M.fr.IX,43.

HERITIERA, Dryander in Aiton, Hort. Kew III, 546 (1789).
H. littoralis, Dryander in Aiton, Hort. Kew III, 546 (1789) ... — — — — — Q. N.A. B.fl.I,231 M.fr.VI,173.

UNGERIA, Schott & Endlicher, Melat. 27—31 (1832).
U. floribunda, Schott & Endlicher, Melat. 27, t. 4 (1832) ... — — — — N.S.W. — — — M.fr.IX,169.

HELICTERES, Plukenet Phytogr. 181, t. 245 (1692).
H. Isora, Linné, sp. pl. 963 (1753) ... ... — — — — — — N.A. B.fl.I,232 M.fr.VI,173.

METHORIUM, Schott & Endlicher, Melat. 29—30 (1832).
M. spicatum, Helicteris, Colebrooke in G. Don, syst. I, 507 (1831) — — — — — Q. — — M.fr.VII,39.
M. canum, Schott, Melatem. 29, t. 5 (1832) ... ... — — — — — — N.A. B.fl.I,232
M. dentatum, F. v. M. in Benth. Fl. Austr. I, 232 (1863) ... — — — — — — N.A. B.fl.I,232
M. semiglabrum, F. v. M., fragm. V, 43 (1865) ... ... — — — — — Q. — — M.fr.V,43.

MELHANIA, Forskael, Fl. Aeg. Arab. 64 (1775).
M. incana, Heyne in Wight & Arn. prodr. 68 (1834) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.I,234 M.fr.I,69;VI,174.

MELOCHIA, Dillenius, Hort. Eltham. 221, t. 176 (1732). (Riedleya).
M. pyramidata, Linné, sp. pl. 674 (1753) ... ... — — — — — Q. N.A. B.fl.I,234 M.fr.VI,173.
M. corchorifolia, Linné, sp. pl. 675 (1753) ... ... — — — — — Q. N.A. B.fl.I,235 M.fr.VI,173.

HERMANNIA, Tournefort, inst. 656, t. 432 (1700). (Gilesia, Mahernia partly).
H. Gilesii, F. v. M., fragm. IX, 42 (1875) ... ... — S.A. — — — — — — M.fr.IX,42.

DICARPIDIUM, F. v. Mueller in Hook. Kew misc. IX, 302 (1857).
D. monoicum, F. v. M., in Hook. Kew misc. IX, 302 (1857) ... — — — — — — N.A. B.fl.I,235

WALTHERIA, Linné, gen. pl. 203 (1737).
W. Indica, Linné, sp. pl. 637 (1753) ... ... — S.A. — — — Q. N.A. B.fl.I,236 M.fr.VI,173.

ABROMA, N. Jacquin, Hort. Vindob. III, t. 1 (1776).
A. fastuosa, Salisbury, Parad. Lond. 102 (1807) ... ... — — — — — Q. — B.fl.I,236 M.fr.VI,173.

COMMERCONIA, R. & G. Forster, char. gen. 43, t. 22 (1776). (Ruelingia, Achilleopsis).

C. salvifolia, Bentham, Fl. Austr. I, 238 (1863)	—	—	—	—	—	Q.	—	B.fl.I,238	
C. dasyphylla, Andrews, Bot. Reposit. t. 603 (1804)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,238	M.fr.X,22.
C. rugosa, Rulingia, Steetz in pl. Preiss. II, 352 (1847) ...	—	—	—	—	N.S.W.	—	—	B.fl.I,238	
C. Preissii, Steudel in Lehm. pl. Preiss. I, 237 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,239	
C. cinerea, Steudel in Lehm. pl. Preiss. I, 238 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,239	
C. cygnorum, Steudel in Lehm. pl. Preiss. I, 237 (1844)... ...	W.A.	—	—	—	—	—	—	B.fl.I,239	
C. platycalyx, Rulingia, Benth. fl. Austr. I, 240 (1863)	W.A.	—	—	—	—		—	B.fl.I,240	
C. parviflora, Rulingia, Endlicher in Hueg. enum. 12 (1837) ...	W.A.	—	—	—	—	—	—	B.fl.I,240	
C. hermanniaefolia, Rulingia, Steetz in pl. Preiss. II, 353 (1847)	—	—	—	—	N.S.W.	—	—	B.fl.I,240	
C. Kempeana, F. v. M., fragm. XI, 113 (1881)	—	S.A.	—	—	—	—	—	—	M.fr.XI,113.
C. laxophylla, F. v. M., fragm. X, 22 (1876)	—	S.A.	—	—	—	—	N.A.	B.fl.I,240	M.fr.I,68;X,22.
C. magniflora, F. v. M., fragm. VIII, 223 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.X,22;XI,114.
C. cuneata, Rulingia, Turczan. in Bull. Mosc. XXV, 151 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,241	
C. rotundifolia, Rulingia, Turcz. in Bull. de Mosc. XXV, 151 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,241	
C. densiflora, F. v. M., fragm. X, 21 (1876)	W.A.	—	—	—	—	—	—	B.fl.I,241	M.fr.VI,175;XI,114.
C. Fraseri, J. Gay in Mém. du Mus. Par. X, 215, t. 15 (1823) ...	—	—	—	V.	N.S.W.	—	—	B.fl.I,242	
C. Leichhardtii, Bentham, Fl. Austr. I, 242 (1863)	—	—	—	—	—	Q.	—	B.fl.I,242	M.fr.VI,175;X,22;XI,114.
C. echinata, R. & G. Forster, char. gen. 43, t. 22 (1776)... ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,243	M.fr.VI,175;X,22.
C. Gaudichaudi, J. Gay in Mém. du Mus. Par. X, 213, t. 14 (1823)	W.A.	—	—	—	—	—	—	B.fl.I,243	
C. crispa, Turczaninow in Bull. Soc. Mosc. XIX, 501 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.I,243	
C. melanopetala, F. v. M., fragm. X, 21 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,21.
C. pulchella, Turczaninow in Bull. Soc. Mosc. XIX, 502 (1846)	W.A.	—	—	—	—	—	—	B.fl.I,244	M.fr.X,22.
C. microphylla, Bentham, Fl. Austr. I, 244 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,244	
C. craurophylla, F. v. M., fragm. IX, 59 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,59.

HANNAFORDIA, F. v. Mueller, fragm. II, 9 (1860).

H. quadrivalvis, F. v. M., fragm. II, 9 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,248	M.fr.XI,20&115.
H. Shanesii, F. v. M., fragm. VI, 175 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VI,175.
H. Bissillii, F. v. M., fragm. X, 95 (1877)	W.A.	S.A.	—	—	N.S.W.	—	—	—	M.fr.XI,115.

SERINGEA, Sprengel, Anleit. II, 649 (1818). (Keraudrenia).

S. platyphylla, J. Gay in Mém. du Mus. Par. VII, 443 (1821) ...	—	—	—	—	N.S.W.	—	—	B.fl.I,244	M.fr.II,5;VI,175;X,97
S. lanceolata, Steetz in Lehm. pl. Preiss. II, 349 (1847)	—	—	—	—	—	Q.	—	B.fl.I,245	M.fr.VI,174;X,97.
S. corollata, Steetz in Lehm. pl. Preiss. II, 330 (1847)	—	S.A.	—	—	—	Q.	N.A.	B.fl.I,246	M.fr.II,5;VI,174;X,96.
S. Hillii, Keraudrenia, F. v. M. in Benth. Fl. Austr. I, 246 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,246	M.fr.VI,174;X,97.
S. nephrosperma, F. v. M. in Hook. Kew misc. IX, 15 (1857) ...	—	S.A.	—	—	—	—	N.A.	B.fl.I,246	M.fr.II,5;X,97.
S. microphylla, F. v. M., fragm. II, 5 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,247	M.fr.II,5.
S. adenolasia, F. v. M., fragm. X, 96 (1877)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,96.
S. integrifolia, F. v. M., fragm. II, 5 (1860)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,247	M.fr.II,5;VI,175;X,97

THOMASIA, J. Gay in Mém. du Mus. VII, t. 6—7 (1821). (Leucothamnus, Rhychostemon).

T. macrocarpa, Huegel in Endlicher, nov. stirp. dec. 32 (1839)...	W.A.	—	—	—	—	—	—	B.fl.I,250	
T. rugosa, Turczaninow in Bull. Soc. Mosc. XIX, 501 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.I,250	
T. montana, Steudel in Lehm. pl. Preiss. I, 230 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,250	
T. tenuivestita, F. v. M., fragm. II, 7 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,251	M.fr.II,7.
T. solanacea, J. Gay in Mém. du Mus. VII, 456, t. 21 (1821) ...	W.A.	—	—	—	—	—	—	B.fl.I,251	
T. brachystachys, Turczan. in Bull. Soc. Mosc. XXV, 143 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,251	
T. discolor, Steudel in Lehm. pl. Preiss. I, 233 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,251	
T. quercifolia, J. Gay in Mém. du Mus. VII, 459, t. 21 (1821) ...	W.A.	—	—	—	—	—	—	B.fl.I,252	
T. foliosa, J. Gay in Mém. du Mus. VII, 454, t. 22 (1821) ...	W.A.	—	—	—	—	—	—	B.fl.I,252	
T. triloba, Turczaninow in Bull. Soc. Mosc. XIX, 500 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.I,252	
T. triphylla, J. Gay in Mém. du Mus. VII, 458 (1821)	W.A.	—	—	—	—	—	—	B.fl.I,252	
T. purpurea, J. Gay in Mém. du Mus. VII, 452, t. 21 (1821) ...	W.A.	—	—	—	—	—	—	B.fl.I,253	
T. macrocalyx, Steudel in Lehm. pl. Preiss. I, 230 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.I,253	M.fr.VI,175.
T. pauciflora, Lindley, Bot. Regist. XXV, App. XVIII (1839)...	W.A.	—	—	—	—	—	—	B.fl.I,253	
T. rhynchocarpa, Turcz. in Bull. Soc. Mosc. XXV, 142 (1852)...	W.A.	—	—	—	—	—	—	B.fl.I,254	M.fr.II,8;VI,175.
T. grandiflora, Lindley, Bot. Regist. XXV, App. XVIII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,254	
T. cognata, Steudel in Lehm. pl. Preiss. I, 232 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,254	
T. rulingioides, Steudel in Lehm. pl. Preiss. I, 232 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.I,255	
T. angustifolia, Steudel in Lehm. pl. Preiss. I, 232 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.I,255	
T. petalocalyx, F. v. M. in Transact. phil. Soc. Vict. I, 35 (1854)	W.A.	S.A.	—	V.	—	—	—	B.fl.I,255	
T. Sarotes, Turczaninow in Bull. Soc. Mosc. XXV, 145 (1852)...	W.A.	—	—	—	—	—	—	B.fl.I,255	

GUICHENOTIA, J. Gay in Mém. du Mus. VII, 448, t. 5 (1821). (Sarotes).

G. ledifolia, J. Gay in Mém. du Mus. VII, 449, t. 5 (1821) ...	W.A.	—	—	—	—	—	—	B.fl.I,258	M.fr.VII,174;XI,114.
G. macrantha, Turczaninow in Bull. Soc. Mosc. XIX, 500 (1846)	W.A.	—	—	—	—	—	—	B.fl.I,258	
G. semihastata, Bentham, Fl. Austr. I, 258 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,258	M.fr.II,4.
G. Sarotes, Bentham, Fl. Austr. I, 258 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,258	
G. micrantha, Bentham, Fl. Austr. I, 258 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,258	M.fr.II,7.

LASIOPETALUM, Smith in Transact. Linn. Soc. IV, 216 (1798). (Corethrostylis, Asterochiton).

L. laxiflorum, F. v. M., fragm. XI, 112 (1881)	W.A.	—	—	—	—	—	—	B.fl.I,256	M.fr.XI,112.
L. pymacum, Bentham, Austr. I, 256 (1863)	W.A.	—	—	—	—	—	—	—	M.fr.XI,113.
L. stelligerum, Bentham, Austr. I, 256 (1863)	W.A.	—	—	—	—	—	—	—	M.fr.XI,113.
L. glutinosum, F. v. M., fragm. XI, 113 (1881)	W.A.	—	—	—	—	—	—	—	M.fr.XI,113.
L. discolor, Hooker, Comp. to Bot. Mag. I, 276 (1835)	W.A.	S.A.	T.	—	—	—	—	B.fl.I,260	M.fr.XI,110.
L. dasyphyllum, Sieber in Hook. journ. of bot. II, 414 (1840) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.I,261	M.fr.VI,174;XI,110.
L. indutum, Steudel in Lehm. pl. Preiss. I, 235 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,261	M.fr.XI,110.
L. Maxwelli, F. v. M., fragm. XI, 107 (1881)	W.A.	—	—	—	—	—	—	—	M.fr.XI,107.
L. Ogilvieanum, F. v. M., fragm. XI, 107 (1881)...	W.A.	—	—	—	—	—	—	—	M.fr.XI,107.
L. Behrii, F. v. M., in Transact. phil. Soc. Vict. I, 36 (1854) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,261	
L. parviflorum, Rudge in Transact. Linn. Soc. X, 297, t. 19 (1811)	—	—	—	V.	N.S.W.	—	—	B.fl.I,262	M.fr.VI,174.
L. parvuliflorum, F. v. M., fragm. VI, 174 (1868)	W.A.	—	—	—	—	—	—	—	M.fr.VI,174.
L. Fitzgibbonii F. v. M. in South. Sc. Record II, Apr. (1882)...	W.A.	—	—	—	—	—	—	—	M.fr.XII.

L. oppositifolium, F. v. M., fragm. II, 5 (1860)	W.A	—	—	-	-	-	-	-	B.fl.I,262	M.fr.XI,112.
L. micranthum, J. Hooker, Fl. Tasman. I, 51 (1860)	—	—	T.	—	—	--	—	B.fl.I,262	
L. macrophyllum, Graham in Bot. Mag. t. 3098 (1844)	—	—	—	—	N.S.W.	-	-	B.fl.I,262	
L. Baueri, Steetz in Lehm. pl. Preiss. II, 339 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,263	M.fr.VI,174;XI,110.
L. Tepperi, F. v. M., fragm. XI, 109 (1881)	—	S.A.	—	—	—	-	-	—	M.fr.XI,109.
L. rufum, R. Brown in Benth. Fl. Austr. I, 263 (1863)	—	—	—	—	N.S.W.		--	B.fl.I,263	
L. ferrugineum, Smith in Bot. Reposit. t. 308 (1799)	—	—	—	V.	N.S.W.	-	—	B.fl.I,263	
L. acutiflorum, Turczaninow in Bull. Soc. Mosc. XXV, 145 (1852)	W.A.	—	—	—	—		-	B.fl.I,264	M.fr.XI,109.
L. Oldfieldii, F. v. M., fragm. II, 6 (1860)	W.A.	—	—	—	—	-	—	—	M.fr.II,6.
L. quinquenervium, Turcz. in Bull. Soc. Mosc. XXV, 146 (1852)	W.A.	—	—	—	—	-	--	—	M.fr.XI,109.
L. Drummondii, Bentham, Fl. Austr. I, 264 (1863)	W.A.	—	—	—	-	—	—	B.fl.I,264	M.fr.VI,174;XI,112.
L. rosmarinifolium, Bentham, Fl. Austr. I, 264 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,264	M.fr.XI,112.
L. cordifolium, Endlicher in Hueg. enum. 10 (1837)	W.A.	—	—	—		—	-	B.fl.I,265	M.fr.VI,174;XI,112.
L. Schulzenii, F. v. M., pl. of Vict. I, 145 implied (1861) ...	—	S.A.	—	V.	—	—	—	B.fl.I,265	
L. floribundum, Bentham, Fl. Austr. I, 265 (1863)	W.A.	—	--	—	—	—	—	B.fl.I,265	M.fr.XI,112.
L. molle, Bentham, Fl. Austr. I, 265 (1863)	W.A.	—	—	—	-	—	—	B.fl.I,265	M.fr.XI,111.
L. membranaceum, Bentham, Fl. Austr. I, 266 (1863)	W.A.	--	—	—	—	—	—	B.fl.I,266	M.fr.XI,112.
L. bracteatum, Bentham, Fl. Austr. I, 266 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,266	
LYSIOSEPALUM, F. v. Mueller, fragm. I, 143 (1859).										
L. Barryanum, F. v. M., fragm. I, 143 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,267	M.fr.VI,174.
L. rugosum, Bentham, Fl. Austr. I, 267 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,267	

TILIACEAE.

A. L. de Jussieu, Gen. 289 (1789).

BERRYA, Roxburgh, Pl. Coromand. III, 60, t. 264 (1819).									
B. ammonilla, Roxburgh, pl. Coromand. III, 60, t. 264 (1819) ...	—	—	—	—	—	Q.	—	B.fl.I,268	
GREWIA, Linné, syst. nat. 9 (1735); Linné, gen. pl. 276 (1737).									
G. orientalis, Linné, sp. pl. 964 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,270	
G. multiflora, A. L. de Jussieu in Ann. du Mus. IV, 89, t. 47 (1804)	—	—	—	—	—	Q.	N.A.	B.fl.I,270	M.fr.VI,172.
G. breviflora, Bentham, Fl. Austr. I, 270 (1863)	—	—	-	—	—	—	N.A.	B.fl.I,270	
G. latifolia, F. v. M. in Benth. Fl. Austr. I, 271 (1863)		—	—	—	N.S.W.	Q.	—	B.fl.I,271	M.fr.VI,172;VIII,4.
G. polygama, Roxburgh, Fl. Ind. II, 588 (1832)	--	—	—	—	—	Q.	N.A.	B.fl.I,271	
G. xanthopetala, F. v. M. in Benth. Fl. Austr. I, 271 (1863) ...	—	—	—	-	—	—	N.A.	B.fl.I,271	
G. scabrella, Bentham, Fl. Austr. I, 272 (1863)	—	—	—	—	—	Q.	—	B.fl.I,272	
G. orbifolia, F. v. M. in Benth. Fl. Austr. I, 272 (1863)... ...	—	—	—	—	—	—	N.A.	B.fl.I,272	
G. pleiostigma, F. v. M., fragm. VIII, 4 (1872)	—	—	—	-	—	Q.	—	—	M.fr.VIII,4.
TRIUMFETTA, Plumier, nov. gen. 40, t. 8 (1703).									
T. procumbens, G. Forster, prodr. fl. insul. Austr. 35 (1786) ...	—	—	—	—	—	Q.	—	B.fl.I,273	M.fr.III,8;VI,172.
T. appendiculata, F. v. M., fragm. III, 7 (1862)	—	—	—	—	—	—	N.A.	B.fl.I,273	M.fr.VIII,5.
T. chactocarpa, F. v. M., fragm. XI, 61 (1879)	—	—	—	—	-	—	N.A.	—	M.fr.XI,61.
T. glauscescens, R. Brown in Benth. Fl. Austr. I, 273 (1863) ...	—	—	—	--	—	—	N.A.	B.fl.I,273	
T. denticulata, R. Brown in Benth. Fl. Austr. I, 273 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.I,274	
T. micracantha, F. v. M., fragm. III, 7 (1862)	—	—	—	—	-	—	N.A.	B.fl.I,274	M.fr.VIII,5;XI,62.
T. leptacantha, F. v. M., fragm. XI, 62 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,62.
T. plumigera, F. v. M., fragm. I, 69 (1858)	—	-	—	—	—	—	N.A.	B.fl.I,274	M.fr.III,8;VI,172.
T. pilosa, Roth, nov. pl. spec. 223 (1821)	—	-	—	--	—	Q.	—	—	M.fr.IV,28;VIII,5.
T. parviflora, Bentham, Fl. Austr. I, 274 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,274	
NETTOA, Baillon in Adans. VI, 238, t. 7 (1866).									
N. chrozophorifolia, Baillon in Adans. VI, 238, 242 (1866) ...	—	—	—	—	—	—	N.A.	—	
CORCHORUS, Tournefort, inst. 259, t. 135 (1700).									
C. echinatus, Bentham, Fl. Austr. I, 276 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,276	M.fr.III,8;VIII,5.
C. hygrophilus, Cunningham in Benth. Fl. Austr. I, 276 (1863)	—	—	—	—	—	Q.	—	B.fl.I,276	M.fr.VIII,5.
C. Cunninghamii, F. v. M., fragm. III, 8 (1862)	—	—	-	—	N.S.W.	Q.	—	B.fl.I,276	M.fr.167;VI,172.
C. olitorius, Linné, sp. pl. 529 (1753)	—	—	—	—	—	—	N.A.	B.fl.I,276	M.fr.VIII,5.
C. tridens, Linné, mantiss. alter. 566 (1771)	—	—	—	—	—	—	N.A.	B.fl.I,276	
C. acutangulus, Lamarck, encycl. meth. II, 104 (1786)	—	—	—	--	—	Q.	N.A.	B.fl.I,277	M.fr.VIII,6.
C. fascicularis, Lamarck, encycl. meth. II, 104 (1786)	—	--	--	-	—	—	N.A.	B.fl.I,277	
C. pumilio, Bentham, Fl. Austr. I, 277 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,277	
C. vermicularis, F. v. M., fragm. III, 10 (1862)	—	—	—	—	—	—	N.A.	B.fl.I,277	M.fr.VIII,5.
C. tomentellus, F. v. M., fragm. III, 10 (1862)	—	-	—	—	—	Q.	—	B.fl.I,278	M.fr.III,10.
C. sidoides, F. v. M., fragm. III, 9 (1862)	W.A.	S.A.	--	—	—	Q.	N.A.	B.fl.I,278	M.fr.VIII,6;XI,62.
C. leptocarpus, Cunningham in Benth. Fl. Austr. I, 278 (1863)...	—	--	—	—	—	—	N.A.	B.fl.I,278	
C. elachocarpus, F. v. M., fragm. VIII, 6 (1872)	—	-	—	—	—	—	N.A.	—	M.fr.VIII,6. [62
C. Walcottii, F. v. M., fragm. III, 9 (1862)	—	—	—	—	—	—	N.A.	B.fl.I,278	M.fr.VI,172;VIII,6; XI.
C. trilocularis, Linné, mantiss. 77 (1771)	—	—	—	—	—	Q.	—	—	M.fr.VIII,5.
SLOANEA, Linné, hort. Cliffort. 210 (1737). (Echinocarpus.)									
S. Australis, F. v. M., fragm. IV, 91 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,279	M.fr.VI,172;VIII,2.
S. Langii, F. v. M., fragm. V, 28 (1865)	—	—	—	—	—	Q.	—	—	M.fr.VIII,2.
S. Macbrydei, F. v. M., fragm. VI, 170 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VI,170.
S. Woollsii, F. v. M., fragm. VI, 171 (1868)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VIII,2.
ARISTOTELIA, L'Héritier, stirp. II, 21, t. 16 (1784). (Friesia.)									
A. peduncularis, J. Hooker, Fl. Nov. Fel. I, 33 (1853)	—	—	T.	—	—	—	—	B.fl.I,280	M.fr.VIII,2.
A. Australasica, F. v. M., fragm. II, 79 (1860)	—	—	—	—	N.S.W.	—	—	B.fl.I,280	M.fr.VIII,2.
A. megalosperma, F. v. M., fragm. IX, 84 (1875)...	—	—	—	—	—	Q.	—	—	M.fr.IX,84.
ELAEOCARPUS, Burman, thesaur. Zeil. 39, t. 40 (1737). (Elaiocarpos, Dicera.)									[VIII.
E. holopetalus, F. v. M., fragm. II, 143 (1861)	—	—	—	V.	N.S.W.	—	—	B.fl.I,281	M.fr.IV,173;VI,172;
E. Arnhemicus, F. v. M., Docum. intercol. Exhib. 24 (1867) ...	—	—	—	—	—	—	N.A.	—	M.fr.X,4.

E. obovatus, G. Don, gen. syst. dichl. pl. I, 599 (1831)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,281	M.fr.II,80;VI,172.
E. cyaneus, Aiton, Epit. hort. Kew, addend. 367 (1814) ..	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,281	M.fr.VI,172;VIII,2. [12.
E. grandis, F. v. M., fragm. II, 81 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,281	M.fr.II.175;VI.172;VIII,
E. foveolatus, F. v. M., fragm. V, 157 (1866)	—	—	—	—	—	Q.	—	—	M.fr.VI,172;VIII,2.
E. ruminatus, F. v. M., fragm. VIII, 1 (1872)	—	—	—	—	—	Q.	—	—	M.fr.X,4.
E. Grahami, F. v. M., fragm. X, 3 (1876)	—	—	—	—	—	Q.	—	—	M.fr.X,3.
E. sericopetalus, F. v. M., fragm. VI. 171 (1868)...	—	—	—	—	—	Q.	—	—	M.fr.VI,171.

EUPHORBIACEAE.

A. L. de Jussieu, Gen. 384 (1789) from B. de Jussieu (1759).

CALYCOPEPLUS, Planchon in Bull. de la Soc. Bot. de Fr. VIII, 31 (1861).

C. ephedroides, Planchon in Bull. Soc. Bot. Fr. VIII, 31 (1861)	W.A.	—	—	—	—	—	—	B.fl.VI,53	
C. marginatus, Bentham, Fl. Austr. VI, 53 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,53	

EUPHORBIA, Linné, gen. pl. 152 (1737) from Plinius.

E. Atoto, G. Forster, florul. ins. Austr. prodr. 36 (1786) ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,46	
E. Sparmanni, Boissier, Cent. Euphorb. 5 (1860)	—	—	—	—	N.S.W.	—	—	B.fl.VI,46	
E. Mitchelliana, Boissier in De Candolle, prodr. XV, 25 (1866)	W.A.	—	—	—	—	Q.	N.A.	B.fl.VI,47	M.fr.VI,182.
E. schizolepis, F. v. M. in Benth. Fl. Austr. VI, 47 (1873) ...	—	—	—	—	—	—	N.A.	B.fl.VI,47	
E. Schultzii, Bentham in Fl. Austr. VI, 47 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,47	
E. Armstrongiana, Boissier in De Candolle, prodr. XV, 47 (1866)	—	—	—	—	—	—	N.A.	B.fl.VI,48	
E. erythrantha, F. v. M., fragm. II, 152 (1861)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VI,48	M.fr.XI,64.
E. Muelleri, Boissier in De Candolle, prodr. XV, 27 (1866) ...	—	—	—	—	—	—	N.A.	B.fl.VI,48	
E. Drummondii, Boissier, Cent. Euphorb. 14 (1860)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VI,49	M.fr.VI,182.
E. alsiniflora, Baillon, Adans. VI, 288 (1866)	—	—	—	—	—	—	N.A.	B.fl.VI,49	
E. Wheeleri, Baillon, Adans. VI, 286 (1866)	—	S.A.	—	—	—	—	N.A.	B.fl.VI,49	
E. myrtoides, Boissier in De Candolle, prodr. XV, 15 (1866) ...	W.A.	—	—	—	—	—	N.A.	B.fl.VI,50	
E. micradenia, Boissier in De Candolle, prodr. XV, 27 (1866) ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,50	
E. Macgillivrayi, Boissier in De Candolle, prodr. XV, 26 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,50	
E. serrulata, Reinwardt in Blum. Bijdrag. 635 (1826)	—	—	—	—	—	Q.	N.A.	B.fl.VI,51	
E. filipes, Bentham, Fl. Austr. VI, 51 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,51	
E. Careyi, F. v. M., fragm. XI, 64 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,64.
E. pilulifera, Linné, sp. pl. 454 (1753)	—	—	—	—	—	Q.	—	B.fl.VI,51	
E. Norfolkiana, Boissier in De Cand., prodr. XV, 206 (1866) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.
E. eremophila, Cunningham in Mitch. Trop. Austr. 348 (1848)...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,52	

MONOTAXIS, Brongniart in Duperr. Voy. Bot. 223, t. 49 (1829). (Hippocrepandra.)

M. macrophylla, Bentham, Fl. Austr. VI, 79 (1873)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,79	
M. linifolia, Brongniart in Duperr. Voy. Bot. 223, t. 49 (1829)...	—	—	—	—	N.S.W.	—	—	B.fl.VI,79	
M. occidentalis, Endlicher in Hueg. enum. 19 (1837)	W.A.	—	—	—	—	—	—	B.fl.VI,79	
M. lurida, Bentham, Fl. Austr. VI, 80 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,80	
M. megacarpa, F. v. M., fragm. IV, 143 (1865)	W.A.	—	—	—	—	—	—	B.fl.VI,80	M.fr.IV,143.
M. gracilis, Baillon, Adans. VI, 293 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,80	
M. grandiflora, Endlicher in Hueg. enum. 19 (1837)	W.A.	—	—	—	—	—	—	B.fl.VI,81	
M. luteiflora, F. v. M., fragm. X, 51 (1877)	W.A.	—	—	—	—	—	—	—	M.fr.X,51.

PORANTHERA, Rudge in Transact. Linn. Soc. X. 302, t. 22 (1811).

P. ericifolia, Rudge in transact. Linn. Soc. X, 302, t. 22 (1811)	—	—	—	—	N.S.W.	—	—	B.fl.VI,55	
P. ericoides, Klotzsch in Lehm. pl. Preiss. II, 239 (1847) ...	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,55	
P. Huegelii, Klotzsch in Lehm. pl. Preiss. II, 231 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.VI,55	
P. corymbosa, Brongniart in Duperr. Voy. Cog. Bot. 219, t. 56 (1826)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,56	M.fr.VI,183.
P. microphylla, Brongniart in Duperr. Voy. Cog. Bot. 218, t. 50 (1826)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VI,56	M.fr.XI,20.

MICRANTHEUM, Desfontaines in Mém. du Mus. hist. nat. IV, 253 (1818). (Caletia.)

M. ericoides, Desfontaines in Mém. du Mus. IV, 253, t. 14 (1818)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,57	
M. hexandrum, J. Hooker in Lond. Journ. VI, 283 (1847) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,57	

PSEUDANTHUS, Sieber in Sprengel, syst. veg. cur. post. 25 (1827). (Stachystemon.)

P. pimeloides, Sieber in Sprengel, syst. veg. cur. post. 25 (1827)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,59	
P. ovalifolius, F. v. M. in Transact. phil. Inst. Vict. II, 66 (1857)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,59	
P. micranthus, Bentham, Fl. Austr. VI, 59 (1873)	—	S.A.	—	—	—	—	—	B.fl.VI,59	
P. divaricatissimus, Bentham, Fl. Austr. VI, 60 (1873)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,60	
P. orientalis, F. v. M., fragm. II, 14 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,60	M.fr.II,14.
P. occidentalis, F, v. M., fragm. I, 107 (1859)	W.A.	—	—	—	—	—	—	B.fl.VI,60	M.fr.I,107.
P. nematophorus, F. v. M., fragm. II, 14 (1860)	W.A.	—	—	—	—	—	—	B.fl.VI,60	M.II,14.
P. polyandrus, F. v. M., fragm. II, 153 (1861)	W.A.	—	—	—	—	—	—	B.fl.VI,62	M.fr.II,153;IV,173
P. brachyphyllus, F. v. M. in Transact. R. Soc. N. S. Wales 11 (1881)	W.A.	—	—	—	—	—	—	B.fl.VI,62	
P. vermicularis, F. v. M. in Transact. R. Soc. N.S.W. 11 (1881)	W.A.	—	—	—	—	—	—	B.fl.VI,62	

BEYERIA, Miquel in Ann. sc. nat. sér. trois. I, 350 (1844). (Calyptrostigma, Beyeriopsis.)

B. viscosa, Miquel in Ann. des. sc. nat. sér. trois. I, 350, t. 15 (1844)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,64	M fr.I,230.
B. lasiocarpa, F. v. M. in Benth. Fl. Austr. VI, 65 (1873) ...	—	—	—	V.	N.S.W.	—	—	B.fl.VI,65	M.fr.VI,182.
B. opaca, F. v. M. in Transact. phil. Soc. Vict. I, 16 (1854) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,65	M.fr.VI,182.
B. uncinata, F. v. M. in Benth. Fl. Austr. VI, 65 (1873) ...	—	S.A.	—	—	—	—	—	B.fl.VI,65	
B. latifolia, Baillon, Adans. VI, 304 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,66	
B. cygnorum, Baillon, Adans. VI, 309 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,66	
B. cinerea, Baillon, Adans. VI, 309 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,66	
B. cyanescens, Bentham, Fl. Austr. VI, 66 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,66	
B. lepidopetala, F. v. M., fragm. I, 230 (1859)	W.A.	—	—	—	—	—	—	B.fl.VI,67	M.fr.I,230;III,164.
B. similis, Baillon, Adans. VI, 309 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,67	
B. brevifolia, Baillon, Adans. VI, 309 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,67	
B. Drummondii, J. Mueller in Linnaea XXXIV, 58 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.VI,68	
B. tristigma, F. v. M., fragm. VI, 181 (1868)	—	—	—	—	—	Q.	—	B.fl.VI,68	M.fr.VI,181.

RICINOCARPUS, Desfontaines in Mém. du Mus. III, 459, t. 22 (1817). (Roeperia.)

R. pinifolius, Desfontaines in Mém. du Mus. III, 459, t. 22 (1817)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,70	M.fr.VI,182.
R. tuberculatus, J. Mueller in Linnaea XXXIV, 60 (1865)	W.A.	—	—	—	—	—	—	B.fl.VI,70	
R. cyanescens, J. Mueller in Linnaea XXXIV, 60 (1865)	W.A.	—	—	—	—	—	—	B.fl.VI,70	
R. psilocladus, Bentham, Fl. Austr. VI, 71 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,70	
R. glaucus, Endlicher in Hueg. enum. 18 (1837)	W.A.	—	—	—	—	—	—	B.fl.VI,71	M.fr.VI,182.
R. major, J. Mueller in Linnaea XXXIV, 59 (1865)	—	—	T.	—	—	—	—	B.fl.VI,72	
R. Bowmanii, F. v. M., fragm. I, 181 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,72	M.fr.I,181.
R. ledifolius, F. v. M., fragm. I, 76 (1858)	—	—	—	—	—	Q.	—	B.fl.VI,72	M.fr.I,76.
R. rosmarinifolius, Bentham, Fl. Austr. VI, 72 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,72	
R. marginatus, Bentham, Fl. Austr. VI, 73 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,73	
R. speciosus, J. Mueller in De Candolle, prodr. XV, 204 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.VI,73	
R. trichophorus, J. Mueller in Linnaea XXXIV, 60 (1865)	W.A.	—	—	—	—	—	—	B.fl.VI,73	
R. muricatus, J. Mueller in Linnaea XXXIV, 61 (1865)	W.A.	—	—	—	—	—	—	B.fl.VI,73	
R. velutinus, F. v. M., fragm. IX, 2 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,2.

BERTYA, Planchon in Hook. Lond. Journ. Bot. IV, 472, t. 16 (1845).

B. gummifera, Planchon in Hook. Lond. Journ. IV, 473 (1845)	—	—	—	—	N.S.W.	—	—	B.fl.VI,75	
B. pinifolia, Planchon in Hook. Lond. Journ. IV, 473 (1845)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,75	
B. quadrisepala, F. v. M., fragm. X, 52 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,52.
B. Cunninghamii, Planchon in Hook. Lond. Journ. IV, 473 (1845)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,75	M.fr.X,52.
B. dimerostigma, F. v. M. in Wing's S. Science Journ. II (1882)	W.A.	—	—	—	—	—	—	—	M.fr.XII.
B. rosmarinifolia, Planchon in Hook. Lond. Journ. IV, 473 (1845)	—	—	T.	—	N.S.W.	—	—	B.fl.VI,76	
B. Mitchelli, J. Mueller in Linnaea XXXIV, 63 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,76	
B. oppositifolia, F. v. M. & O'Shanesy in S. Sc. Rec. II, May (1882)	—	—	—	—	—	Q.	—	—	M.fr.XII.
B. oleaefolia, Planchon in Hook. Lond. Journ. IV, 473 (1845)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,76	
B. rotundifolia, F. v. M., fragm. IV, 34 (1864)	—	S.A.	—	—	—	—	—	B.fl.VI,77	M.fr.IV,34.
B. pedicellata, F. v. M., fragm. IV, 143 (1865)	—	—	—	—	—	Q.	—	B.fl.VI,77	M.fr.IV,143.
B. pomaderrioides, F. v. M., fragm. IV, 34 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.VI,77	M.fr.IV,34.
B. Findlayi, F. v. M., fragm. VIII, 141 (1874)	—	—	—	V.	N.S.W.	—	—	—	M.fr.VIII,141.

AMPEREA, Adr. de Jussieu, Euphorb. gen. 35, t. 10 (1824).

A. protensa, Nees in Lehm. pl. Preiss. II, 229 (1847)	W.A.	—	—	—	—	—	—	B.fl.VI,82	
A. volubilis, F. v. M. in Benth. Fl. Austr. VI, 82 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,82	M.fr.VI,182.
A. micrantha, Bentham, Fl. Austr. VI, 83 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,83	
A. conferta, Bentham, Fl. Austr. VI, 83 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,83	
A. ericoides, Adr. de Jussieu, Tent. Euphorb. 112, t. 10 (1824)	W.A.	—	—	—	—	—	—	B.fl.VI,83	
A. spartioides, Brongniart in Duperr. Voy. Cog. 221, t. 49 (1826)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,84	

ANTIDESMA, Linné, Fl. Zeyl. 169 (1747).

A. Ghaesembilla, Gaertner, de Fruct. I, 189, t. 39 (1788)	—	—	—	—	—	—	N.A.	B.fl.VI,85	M.fr.VI,183.
A. Dallachyanum, Baillon, Adans. VI, 337 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.VI,85	M.fr.VI,183.
A. Bunius, Sprengel, syst. veg. I, 826 (1825)	—	—	—	—	—	Q.	—	B.fl.VI,86	
A. parviflorum, Thwaites & F. v. M., fragm. IV, 86 (1864)	—	—	—	—	—	Q.	—	B.fl.VI,86	M.fr.IV,86;VI,183.
A. Schultzii, Bentham, Fl. Austr. VI, 86 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,86	
A. erostre, F. v. M. in Benth. Fl. Austr. VI, 87 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,87	
A. sinuatum, Bentham, Fl. Austr. VI, 87 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,87	

ANDRACHNE, Linné, syst. nat. 9 (1735); Linné, gen. pl. 287 (1737).

A. Decaisnei, Bentham, Fl. Austr. VI, 88 (1873)	W.A.	—	—	—	—	Q.	N.A.	B.fl.VI,88	M.fr.VI,183.

ACTEPHILA, Blume, Bijdr. XII, 581 (1825).

A. grandifolia, Baillon, Adans. VI, 330 & 360, t. 10 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,89	
A. Mooreana, Baillon, Adans. VI, 330 & 336 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,89	
A. latifolia, Bentham, Fl. Austr. VI, 89 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,89	
A. petiolaris, Bentham, Fl. Austr. VI, 89 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,89	
A. sessilifolia, Bentham, Fl. Austr. VI, 90 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,90	

DISSILIARIA, F. v. M. in Baillon, Adans. VII, 356, pl. 1 (1867). (Choriceras.)

D. baloghioides, F. v. M. in Baillon, Adans. VII, 356 (1867)	—	—	—	—	—	Q.	—	B.fl.VI,90	
D. Muelleri, Baillon, Adans. VII, 359, t. 1 (1867)	—	—	—	—	—	Q.	—	B.fl.VI,91	
D. tricornis, Bentham, Fl. Austr. VI, 91 (1873)	—	—	—	—	—	Q.	N.A.	B.fl.VI,91	

PETALOSTIGMA, F. v. M. in Hook. Kew Misc. IX, 17 (1857). (Hylococcus.)

P. quadriloculare, F. v. M. in Hook. Kew Misc. IX, 17 (1857)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,92	M.fr.VI,182.

PHYLLANTHUS, Linné, gen. pl. 282 (1737), from J. Commelyn. (Glochidion, Synostemon, Bradleia.)

P. Ferdinandi, J. Mueller in Regensb. Flora, 379 (1865)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,96	M.fr.VI,183.
P. lobocarpus, Bentham, Fl. Austr. VI, 97 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,97	
P. ditassoides, J. Mueller in Regensb. Flora, 487 (1864)	—	—	—	—	—	—	N.A.	B.fl.VI,97	
P. Adami, J. Mueller in De Candolle, prodr. XV, 327 (1866)	—	—	—	—	—	—	N.A.	B.fl.VI,97	
P. thesioides, Bentham, Fl. Austr. VI, 98 (1873)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,98	
P. hirtellus, J. Mueller in De Candolle, prodr. XV, 326 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,98	M.fr.III,89.
P. rigens, J. Mueller in Regensb. Flora, 513 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,99	
P. ochrophyllus, Bentham, Fl. Austr. VI, 99 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,99	
P. rigidulus, F. v. M. in Linnaea XXXIV, 72 (1865)	—	—	—	—	—	—	N.A.	B.fl.VI,99	
P. ramosissimus, J. Mueller in Linnaea XXXIV, 70 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,100	
P. Tatei, F. v. M. in Wing's South. Science Record II, 55 (1882)	—	S.A.	—	—	—	—	—	—	
P. rhytidospermus, F. v. M. in Linnaea XXXIV, 70 (1865)	—	—	—	—	—	—	N.A.	B.fl.VI,100	
P. albiflorus, F. v. M. in Linnaea XXXIV, 70 (1865)	—	—	—	—	—	Q.	—	B.fl.VI,100	
P. crassifolius, J. Mueller in Regensb. Flora, 513 (1864)	W.A.	—	—	—	—	—	—	B.fl.VI,100	
P. elachophyllus, F. v. M. in Benth. Fl. Austr. VI, 101 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,101	
P. uberiflorus, F. v. M. in Adansonia, 343 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,101	
P. reticulatus, Poiret, encycl. meth. V, 298 (1804)	—	—	—	—	—	—	N.A.	B.fl.VI,101	
P. baccatus, F. v. M. in Benth. Fl. Austr. VI, 102 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,102	

P. Urinaria, Linné, sp. pl. 982 (1753)	—	—	—	—	—	—	N.A.	B.fl.VI,102				
P. trachygyne, Bentham, Fl. Austr. VI, 103 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,103				
P. Maderaspatanus, Linné, sp. pl. 982 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VI,103				
P. Mitchelli, Bentham, Fl. Austr. VI, 103 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,103				
P. Gastroemii, J. Mueller in De Cand., prodr. XV, 358 (1866)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,104				
P. Dallachyanus, Bentham, Fl. Austr. VI, 104 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,104				
P. subcrenulatus, F. v. M., fragm. I, 108 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,104	M.fr.I,108.			
P. calycinus, Labillardière, Nov. Holl. pl. spec. II, 75, t. 225 (1806)	W.A.	S.A.	—	—	—	—	—	B.fl.VI,105				
P. flagellaris, Bentham, Fl. Austr. VI, 106 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,106				
P. similis, J. Mueller in Linnaea XXXIV, 71 (1865)	—	—	—	—	—	Q.	—	B.fl.VI,106				
P. microcladus, J. Mueller in Linnaea XXXIV, 71 (1865) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,106				
P. grandisepalus, F. v. M. in Linnaea XXXIV, 72 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.VI,106				
P. Carpentariae, J. Mueller in Linnaea XXXIV, 72 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.VI,107				
P. Fuernrohrii, F. v. M. in Transact. phil. Soc. Vict. I, 15 (1854)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,107				
P. hebecarpus, Bentham, Fl. Austr. VI, 108 (1873)	—	—	—	—	—	Q.	N.A.	B.fl.VI,108				
P. lacunarius, F. v. M. in Transact. phil. Soc. Vict. I, 14 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,108				
P. trachyspermus, F. v. M. in Transact. phil. Soc. Vict. I, 14 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,108				
P. australis, J. Hooker in Lond. Journ. VI, 284 (1847)	—	—	T.	—	N.S.W.	—	—	B.fl.VI,108				
P. thymoides, Sieber in Linnaea XXVIII, 566 (1866)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,109				
P. scaber, Klotzsch in Lehm. pl. Preiss. I, 179 (1844)	W.A.	—	—	—	—	—	—	B.fl.VI,109				
P. indigoferoides, Bentham, Fl. Austr. VI, 110 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,110				
P. aridus, Bentham, Fl. Austr. VI, 110 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,110				
P. Gunnii, J. Hooker in Lond. Journ. VI, 284 (1847)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,110	M.fr.VI,183.			
P. simplex, Retzius, observ. V, 29 (1789)	—	—	—	—	—	Q.	—	B.fl.VI,111				
P. filicaulis, Bentham, Fl. Austr. VI, 111 (1873)	—	—	—	—	N.S.W.	—	—	B.fl.VI,111				
P. minutiflorus, F. v. M. in Linnaea XXXIV, 75 (1863)... ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,112				
P. Armstrongii, Bentham, Fl. Austr. VI, 112 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,112				
P. buxifolius, Reinwardt in Blume, catal. plant. Buitenz. 106 (1823)	—	—	—	—	—	Q.	—	—	M.fr.X,121.			
BREYNIA, R. & G. Forster, char. gen. 145, t. 73 (1776). (Melanthesa, Melanthesiopsis.)												
B. cernua, J. Mueller in De Candolle, prodr. XV, 439 (1866) ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,113				
B. oblongifolia, J. Mueller in De Candolle, prodr. XV, 110 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,114				
B. stipitata, J. Mueller in De Candolle, prodr. XV, 442 (1866) ..	—	—	—	—	—	Q.	N.A.	B.fl.VI,114				
B. rhynchocarpa, Bentham, Fl. Austr. VI, 114 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,114				
SECURINEGA, A. L. de Jussieu, Gen. 388 (1789). (Flueggea.)												
S. Abyssinica, A. Richard, tentam. fl. Abyss. II, 256 (1851) ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,116				
S. Leucopyrus, J. Mueller in De Cand., prodr. XV, 451 (1866)...	—	—	—	—	—	Q.	—	B.fl.VI,116				
NEOROEPERA, F. & J. Mueller in De Candolle, prodr. XV, 489 (1866).												
N. buxifolia, F. & J. Mueller in De Cand., prodr. XV, 489 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,116				
N. Banksii, Bentham, Fl. Austr. VI, 117 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,117				
BISCHOFFIA, Blume, Bijdr. 1168 (1825).												
B. Javanica, Blume, Bijdr. 1168 (1825)	—	—	—	—	—	Q.	—	—	M.fr.VIII,141.			
HEMICYCLIA, Wight & Arnott in Edinb. phil. Journ. XIV, 297 (1833).												
H. sepiaria, Wight & Arnott in Edinb. phil. Journ. XIV, 297 (1833)	—	—	—	—	—	—	N.A.	B.fl.VI,117	M.fr.IV,119;VI,182.			
H. Australasica, J. Mueller in De Candolle, prodr. XV, 487 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,118				
H. lasiogyna, F. v. M., fragm. IV, 119 (1864)	—	—	—	—	—	—	N.A.	B.fl.VI,118	M.fr.IV,119.			
APOROSA, Blume, Bijdrag. 514 (1824).												
A. Australiana, F. v. M., fragm. XII ined.	—	—	—	—	—	Q.	—	—	M.fr.VIII,141;XII.			
BRIDELIA, Willdenow, Spec. IV, 978 (1805).												
B. exaltata, F. v. M., fragm. III, 32 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.VI,119	M.fr.III,32.			
B. ovata, Decaisne in nouv. ann. du Mus. V, 484 (1835)	—	—	—	—	—	—	N.A.	B.fl.VI,120				
B. tomentosa, Blume, Bijdrag. 597 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.VI,120	M.fr.VI,182.			
B. faginea, F. v. M. in Benth. Fl. Austr. VI, 120 (1873)... ...	—	—	—	—	—	Q.	—	B.fl.VI,120				
CLEISTANTHUS, J. Hooker, icon. pl. t. 779 (1847). (Lebediera, Amanoa.)												
C. Cunninghamii, J. Mueller in De Cand., prodr. XV, 506 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,122				
C. apodus, Bentham, Fl. Austr. VI, 122 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,122				
C. Dallachyanus, Baillon in Benth. Fl. Austr. VI, 122 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,122	M.fr.VI,182.			
C. semiopacus, F. v. M. in Benth. Fl. Austr. VI, 123 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,123				
CROTON, Linné, gen. pl. 288 (1737).												
C. Schultzii, Bentham, Fl. Austr. VI, 124 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,124				
C. insularis, Baillon, Adans. II, 217 (1862)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,124				
C. phlebalioides, F. v. M. in Benth. Fl. Austr. VI, 125 (1873) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,125	M.fr.IV,140.			
C. opponens, F. v. M. in Benth. Fl. Austr. VI, 125 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,125				
C. tomentellus, F. v. M., fragm. IV, 141 (1864)	—	—	—	—	—	—	N.A.	B.fl.VI,126	M.fr.IV,141.			
C. Verreauxii, Baillon, Etud. Euph. 357 (1858)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,126	M.fr.IV,141;VI,186.			
C. acronychioides, F. v. M., fragm. IV, 142 (1864)	—	—	—	—	—	Q.	—	B.fl.VI,127	M.fr.VI,186.			
C. triacros, F. v. M., fragm. VI, 185 (1867)	—	—	—	—	—	Q.	—	B.fl.VI,127	M.fr.VI,185.			
C. Arnhemicus, F. v. M. in Linnaea XXXIV, 112 (1865) ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,127	M.fr.VI,186.			
ALEURITES, R. & G. Forster, char. gen. 111, t. 56 (1776).												
A. triloba, R. & G. Forster, char. gen. 111, t. 56 (1776)... ...	—	—	—	—	—	Q.	—	B.fl.VI,128	M.fr.VI,182.			
CLAOXYLON, Adr. de Jussieu, Euphorb. gen. 43, t. 14 (1824).												
C. angustifolium, J. Mueller in Linnaea XXXIV, 165 (1865) ...	—	—	—	—	—	Q.	—	B.fl.VI,129				
C. tenerifolium, F. v. M. in Baillon, Adans. VI, 323 (1866) ...	—	—	—	—	—	Q.	-	B.fl.VI,130	M.fr.VI,183.			
C. australe, Baillon, Etud. Euphorb. 493 (1858)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,130	M.fr.IV,142.			
C. Hillii, Bentham, Fl. Austr. VI, 131 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,131				
ACALYPHA, Royen in Linné, hort. Cliffort. 495 (1737).												
A. nemorum, F. v. M. in Linnaea XXXIV, 38 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,132				

A. eremorum, F. v. M. in Regensb. Flora 440 (1864)	—	—	—	—	—	—	Q.	—	B.fl.VI,132	
A. capillipes, F. v. M. in Linnaea XXXIV, 40 (1865)	—	—	—	—	—	N.S.W.	—	—	B.fl.VI,133	
ADRIANA, Gaudichaud in Ann. sc. nat. V, 223 (1825). (Trachycaryon.)										
A. tomentosa, Gaudichaud in Ann. sc. nat. V, 223 (1825) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,134	M.fr.IX,2.	
A. quadripartita, Gaudichaud in Freyc. voy. Bot. 489 (1826) ...	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,135		
ALCHORNEA, Solander in Swartz, prodr. 698 (1788). (Coelebogyne, Cladodes.)										
A. ilicifolia, J. Mueller in Linnaea XXXIV, 170 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,136		
A. Thozetiana, F. v. M. in Benth. Fl. Austr. VI, 137 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,137		
TRAGIA, Plumier, gen. 14, t. 12 (1703).										
T. Novae Hollandiae, J. Mueller in Linnaea XXXIV, 180 (1865)	—	—	—	—	—	Q.	—	B.fl.VI,137	M.fr.VI,183.	
MALLOTUS, Loureiro, Fl. Cochinch. II, 635 (1790). (Echinus, Rottlera, Echinocroton.)										
M. ricinoides, J. Mueller in Linnaea XXXIV, 187 (1865) ...	—	—	—	—	—	Q.	—	B.fl.VI,140	M.fr.IV,138;VI,184.	
M. Chinensis, Loureiro, Fl. Cochinchin. II, 635 (1770)	—	—	—	—	—	Q.	—	B.fl.VI,140	M.fr.VI,184.	
M. claoxyloides, J. Mueller in Linnaea XXXIV, 192 (1865) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,140	M.fr.VI,184.	
M. Philippinensis, J. Mueller in Linnaea XXXIV, 196 (1865) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,141	M.fr.VI,183.	
M. angustifolius, Bentham in Fl. Austr. VI, 141 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,142		
M. polyadenos, F. v. M., fragm. VI, 184 (1867)	—	—	—	—	—	Q.	—	B.fl.VI,142	M.fr.VI,184.	
M. repandus, J. Mueller in Linnaea XXXIV, 197 (1865)... ...	—	—	—	—	—	Q.	—	B.fl.VI,142	M.fr.VI,185.	
M. nesophilus, F. v. M. in Linnaea XXXIV, 196 (1865)... ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,143	M.fr.VI,184.	
M. discolor, F. v. M. in Benth. Fl. Austr. VI, 143 (1873) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,143	M.fr.IV,139.	
MACARANGA, Petit-Thouars, Gen. Madag. n. 88 (1809). (Mappa.)										
M. Dallachyi, F. v. M. in Benth. Fl. Austr. VI, 144 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,144	M.fr.VI,184.	
M. subdentata, Bentham, Fl. Austr. VI, 145 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,145		
M. inamoena, F. v. M. in Benth. Fl. Austr. VI, 145 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,145		
M. involucrata, Baillon, Etud. Euphorb. 432 (1858)	—	—	—	—	—	Q.	—	B.fl.VI,146	M.fr.IV,139;VI,183.	
M. Tanaria, J. Mueller in De Candolle, prodr. XV, 997 (1866) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,146	M.fr.VI,183.	
CODIAEUM, Rumphius, herbar. Amboin. IV, 65, t. 25 (1743).										
C. variegatum, Blume, Bijdrag. 605 (1825)...	—	—	—	—	—	Q.	—	B.fl.VI,147	M.fr.VI,182.	
BALOGHIA, Endlicher, prodr. fl. Norfolk. 84 (1833).										
B. lucida, Endlicher, prodr. fl. Norfolk. 84 (1833)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,148	M.fr.VI,182.	
FONTAINEA, Heckel, thése inaugurale, Montpell. (1870).										
F. Pancheri, Heckel, thése inaugurale (1870)	—	—	—	—	—	Q.	—	B.fl.VI,149		
OMALANTHUS, Adr. de Jussieu, Euphorb. gen. 50, t. 16 (1824). (Carumbium, Wartmannia.)										
O. populifolius, Graham in Bot. Mag. 2780 (1827)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,150	M.fr.I,32.	
O. stilligiifolius, F. v. M., fragm. I, 32 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,150	M.fr.I,32.	
SEBASTIANIA, Sprengel, Neue Entd. II, 118 (1821). (Microstachys, Elachocroton.)										
S. chamelaea, J. Mueller in De Cand., prodr. XV, 1175 (1866) ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,151		
EXCAECARIA, Linné, syst. veg. ed. decim. 1288 (1759).										
E. Agallocha, Linné, syst. ed. decim. 1288 (1759)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,152	M.fr.XI,64.	
E. Dallachyana, Baillon, Adans. VI, 324 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,153		
E. parvifolia, J. Mueller in Flora, 433 (1864)	—	—	—	—	—	—	N.A.	B.fl.VI,153		

URTICACEAE.

Ventenat, Table III, 524 (1799).

CELTIS, Tournefort, inst. 612, t. 383 (1700), from Camerarius (1586). (Solenostigma.)												
C. Philippinensis, Blanco, Fl. de Filipinas. 197 (1837)	—	—	—	—	—	—	Q.	N.A.	B.fl.VI,156			
C. paniculata, Planchon in Ann. sc. nat. sér. trois. X, 305 (1848)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,156	M.fr.IV,88.			
C. amblyphylla, F. v. M., fragm. IX, 76 (1875)	—	—	—	—	N.S.W.	—	--	—	M.fr.IX,76.			
ULMUS, Tournefort, inst. 601, t. 372 (1700), from Dodoens (1583).												
U. parviflora, N. Jacquin, pl. rar. hort. Schoenbr. descr. et icon. III, t. 269 (1798)	—	—	—	—	—	Q.	—	—	M.fr.VI,101;XI,133.			
TREMA, Loureiro, Fl. Cochinch. II, 562 (1790). (Sponia.)												
T. cannabina, Loureiro, Fl. Cochinch. II, 689 (1790)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,158	M.fr.VIII,247.			
APHANANTHE, Planchon in Ann. sc. nat. X, 265 (1848).												
A. Philippinensis, Planchon in Ann. sc. nat. X, 337 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,160				
FICUS, Tournefort, inst. 662 (1700), from Plinius. (Urostigma, Covellia.)												
F. colossea, F. v. M. in Benth. Fl. Austr. VI, 163 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,163				
F. pilosa, Reinwardt in Blume, Bijdr. 446 (1825)...	—	—	—	—	—	Q.	—	B.fl.VI,164	M.fr.VI,195.			
F. nesophila, F. v. M., Docum. Interc. Exhib. 26 (1866) ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,164	M.fr.VI,195.			
F. Cunninghamii, Miquel in Ann. Mus. Lugd. Bot. III, 286 (1867)	—	—	—	—	—	Q.	—	B.fl.VI,165	M.fr.VI,195;VIII,246.			
F. Henneana, Miquel in Ann. Mus. Lugd. Bot. III, 216 (1867)...	—	—	—	—	—	Q.	N.A.	B.fl.VI,165				
F. validinervis, F. v. M. in Benth. Fl. Austr. VI, 166 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,166				
F. retusa, Linné, mantissa plant. 129 (1767)	—	—	—	—	—	Q.	N.A.	B.fl.VI,166	M.fr.VI,195.			
F. eugenioides, F. v. M., Docum. Interc. Exhib. 26 (1866) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,166	M.fr.VI,195.			
F. Benjaminea, Linné, mantissa plant. 129 (1767)	—	—	—	—	—	Q.	—	B.fl.VI,167	M.fr.VI,195.			
F. Muelleri, Miquel in Ann. Mus. Lugd. Bot. III, 287 (1867) ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,167	M.fr.VI,195;VIII,246.			
F. leucotricha, Miquel in Ann. Mus. Lugd. Bot. III, 285 (1867)	—	—	—	—	—	—	N.A.	B.fl.VI,167	M.fr.VI,195.			
F. rubiginosa, Desfontaines, catal. pl. hort. Paris. 209 (1804) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,168	M.fr.VI,194;VIII,247.			
F. columnaris, F. v. M. & Moore in proc. Accl. Soc. Vict. III, 71 (1874)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,247.			
F. puberula, Cunningham in Hook. Lond. Journ. VI, 562 (1847)	—	—	—	—	—	—	N.A.	B.fl.VI,169				
F. platypoda, Cunningham in Hook. Lond. Journ. VI, 561 (1847)	—	S.A.	—	—	—	Q.	N.A.	B.fl.VI,169	M.fr.VI,194;VIII,247.			
F. dictyophleba, F. v. M. in Benth. Fl. Austr. VI, 170 (1873)...	—	—	—	—	—	Q.	—	B.fl.VI,170				

F. macrophylla, Desfontaines, catal. pl. hort. Paris. 209 (1804)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,170	M.fr.VI,194;VIII,247.	
F. magnifolia, F. v. M., fragm. IV, 50 (1864)	—	—	—	—	—	Q.	—	B.fl.VI,171	M.fr.IV,50&177;VI,196.	
F. ehretioides, F. v. M. in Benth. Fl. Austr. VI, 171 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,171		
F. pumila, Linné, spec. plant. 1060 (1753)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,171	M.fr.VI,196.	[246.
F. coronulata, F. v. M. in Journ. Bot. Neerland 241 (1861) ...	—	—	—	—	—	—	N.A.	B.fl.VI,172	M.fr.IV,49; VI,196;VIII,	
F. leptoclada, Bentham, Fl. Austr. VI, 172 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,172		
F. depressa, Bentham, Fl. Austr. VI, 172 (1873)...	—	—	—	—	—	Q.	—	B.fl.VI,172		
F. Philippinensis, Miquel in Hooker, Lond. Journ. VII, 435 (1848)	—	—	—	—	—	Q.	—	B.fl.VI,173		
F. mollior, F. v. M. in Benth. Fl. Austr. VI, 173 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,173		
F. stenocarpa, F. v. M. in Benth. Fl. Austr. VI, 174 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,174	M.fr.IX,152.	[114.
F. scabra, G. Forster, florul. ins. Austr. prodr. 76 (1786) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,174	M.fr.VI.196;IX,152;X,	
F. orbicularis, Cunningham in Hook. Lond. Journ. VII, 426 (1848)	—	S.A.	—	—	—	—	N.A.	B.fl.VI,175	M.fr.VI,196.	
F. aculeata, Cunningham in Hook. Lond. Journ. VII, 426 (1848)	—	—	—	—	—	—	N.A.	B.fl.VI,175		
F. opposita, Miquel in Hook. Lond. Journ. VII, 426 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,176	M.fr.VI,196;VIII,246.	
F. scobina, Bentham, Fl. Austr. VI, 176 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,176		
F. hispida, Linné, fil. suppl. 442 (1781)	—	—	—	—	—	Q.	N.A.	B.fl.VI,176		
F. fasciculata, F. v. M. in Benth. Fl. Austr. VI, 177 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,177		
F. casearia, F. v. M. in Benth. Fl. Austr. VI, 177 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,177		
F. glomerata, Willdenow, spec. pl. IV, 1148 (1806)	—	—	—	—	—	Q.	N.A.	B.fl.VI,178	M.fr.VI,195.	
F. pleurocarpa, F. v. M., fragm. VIII, 246 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,246.	
CUDRANIA, Trécul in Ann. des. sc. nat. VIII, 122 (1847).										
C. Javanensis, Trécul in Ann. des. sc. nat. VIII, 123 (1847) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,179	M.fr.VIII,247.	
ANTIARIS, Leschenault in Ann. du Mus. XVI, 476 (1810).										
A. macrophylla, R. Brown in Flind. voy. II, 602, t. 5 (1814) ...	—	—	—	—	—	—	N.A.	B.fl.VI,179	M.fr.VI,194.	
MALAISIA, Blanco, Fl. Filip. 789 (1837).										[247.
M. tortuosa, Blanco, Fl. Filip. 789 (1837)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,180	M.fr.VI,193&255;VIII,	
PSEUDOMORUS, Bureau in Ann. des sc. nat. sér. cinq. XI, 372 (1872).										
P. Brunoniana, Bureau in Ann. des sc. nat. sér. cinq. XI, 372 (1872)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,181		
FATOUA, Gaudichaud in Freyc. voy. Bot. 509 (1826).										
F. pilosa, Gaudichaud in Freyc. voy. Bot. 509 (1826)	—	—	—	—	—	—	N.A.	B.fl.VI,182		
ELATOSTEMMA, R. & G. Forster, char. gen. 105, t. 53 (1776).										
E. reticulatum, Weddell in Ann. des sc. nat. sér. 4, I, 188 (1854)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,183		
E. stipitatum, Weddell in Ann. des sc. nat. sér. 4, I, 190 (1854)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,184		
PROCRIS, Commerçon in A. L. de Jussieu, gen. 403 (1789).										
P. montana, Steudel, nomencl. bot. II. 398 (1841)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.	
BOEHMERIA, N. J. Jacquin, stirp. Amer. hist. 246, t. 157 (1763).										
B. calophleba, Moore & Mueller, fragm. VIII, 11 (1870)... ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,184	M.fr.VIII,11.	
B. australis, Endlicher, prodr. Fl. Norfolk, 38 (1833)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.	
PIPTURUS, Weddell in Ann. des sc. nat. sér. quatr. I, 196 (1854).										
P. propinquus, Weddell in Archiv. du Mus. 447, t. 15 (1856) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,185	M.fr.VI,197.	
POUZOLZIA, Gaudichaud in Freyc. voy. Bot. 503 (1826). (Memorialis, Hyrtanandra.)										
P. Indica, Gaudichaud in Freyc. voy. Bot. 503 (1826)	—	—	—	—	—	Q.	N.A.	B.fl.VI,186	M.fr.IV,87.	
P. quinquenervis, Bennett, Pl. Jav. rar. 66 (1852)	—	—	—	—	—	Q.	—	B.fl.VI,187		
PARIETARIA, Tournefort, inst. 509, t. 289 (1700), from C. Bauhin (1623). (Freirea.)										
P. debilis, G. Forster, florul. ins. Austr. prodr. 73 (1786) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VI,188	M.fr.XI,20.	
AUSTRALINA, Gaudichaud in Freyc. voy. Bot. 497 (1826).										
A. pusilla, Gaudichaud in Freyc. voy. Bot. 505 (1826)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,189	M.fr.VIII,247.	
FLEURYA, Gaudichaud in Freyc. voy. Bot. 497 (1826).										
F. interrupta, Gaudichaud in Freyc. voy. Bot. 497 (1826) ...	—	—	—	—	—	Q.	—	—	M.fr.X,114.	
URTICA, Tournefort, inst. 534, t. 308 (1700), from Plinius.										
U. incisa, Poiret, Encycl. meth. suppl. IV, 223 (1817)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,190		
LAPORTEA, Gaudichaud in Freyc. voy. Bot. 498 (1826).										
L. gigas, Weddell in Arch. du Mus. IX, 129, t. 3 & 4 (1856) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,192		
L. photiniphylla, Weddell in Arch. du Mus. IX, 138 (1856) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,192		
L. moroides, Weddell in Arch. du Mus. IX, 142 (1856)	—	—	—	—	—	Q.	—	B.fl.VI,192	M.fr.VIII,248.	

CUPULIFERAE.

L. Cl. Richard, Anal. de fr. 32 & 92 (1808).

FAGUS, Tournefort, inst. 584, t. 351 (1700), from Camerarius (1586).										
F. Gunnii, J. Hooker, icon. plant., t. 881 (1852)	—	—	T.	—	—	—	—	B.fl.VI,210		
F. Cunninghamii, Hooker, Journ. of Bot. II, 152, t. 7 (1840) ...	—	—	T.	V.	—	—	—	B.fl.VI,210		
F. Moorei, F. v. M., fragm. V, 109 (1866)...	—	—	—	—	N.S.W.	—	—	B.fl.VI,211	M.fr.V,109.	
BALANOPS, Baillon, Hist. des pl. VI, 237 & 258 (1876).										
B. Australiana, F. v. M., fragm. X, 114 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,114.	

CASUARINEAE.

Mirbel in Ann. du Mus. XVI, 451 (1810).

CASUARINA, Rumphius, herb. Amboin. III, 87, t. 58 (1743).										
C. quadrivalvis, Labill., nov. Holl. pl. spec. II, 67, t. 218 (1806)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,195	M.fr.VI,18&19;X,62.	
C. trichodon, Miquel in Lehm. pl. Preiss. I, 641 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,196	M.fr.VI,17.	
C. lepidophloia, F. v. M., fragm. X, 115 (1877)	—	S.A.	—	—	N.S.W.	—	—	—	M.fr.X,115.	
C. glauca, Sieber in Sprengel, syst. III, 803 (1826)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,196	M.fr.VI,18.	

C. Huegeliana, Miquel in Lehm. pl. Preiss. I, 640 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.VI,196	M.fr.VI,19;X,62.
C. equisetifolia, R. & G. Forster, char. gen. 103, t. 52 (1776) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,197	M.fr.VI,17.
C. suberosa, Otto & Dietrich, allgem. Gartenzeit. 155 (1841) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,197	M.fr.VI,20&254.
C. Cunninghamiana, Miquel, Revis. Casuarin. 56, t. 6 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,198	
C. distyla, Ventenat, pl. dans le jard. de Cels. t. 62 (1800) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,198	M.fr.VI,20.
C. inophloia, F. v. M. & Bailey in Melb. Chemist, April (1882)	—	—	—	—	—	Q.	—	—	M.fr.XII.
C. Fraseriana, Miquel, Revis. Casuarin. 50, t. 6 (1848)	W.A.	—	—	—	—	—	—	B.fl.VI,199	M.fr.VI,19.
C. nana, Sieber in Sprengel, syst. III, 804 (1826)...	—	—	—	—	N.S.W.	—	—	B.fl.VI,199	
C. humilis, Otto & Dietrich, allgem. Gartenzeit. 163 (1841) ...	W.A.	—	—	—	—	—	—	B.fl.VI,200	M.fr.VI,18.
C. torulosa, Aiton, hort. Kew. III, 320 (1789)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,200	M.fr.VI,17.
C. decussata, Bentham, Fl. Austr. VI, 200 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,200	
C. Decaisneana, F. v. M., fragm. I, 61 (1858)	—	S.A.	—	—	—	—	N.A.	B.fl.VI,201	M.fr.I,61;X,62.
C. Drummondiana, Miquel, Revis. Casuar. 26, t. 1 (1848) ...	W.A.	—	—	—	—	—	—	B.fl.VI,201	
C. microstachya, Miquel in Lehm. pl. Preiss. I, 642 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.VI,201	M.fr.VI,17;X,62.
C. acutivalvis, F. v. M., fragm. X, 61 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,61.
C. bicuspidata, Bentham, Fl. Austr. VI, 202 (1873)	W.A.	S.A.	T.	—	—	—	—	B.fl.VI,202	
C. thuyoides, Miquel in Lehm. pl. Preiss. I, 641 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,202	M.fr.VI,17.
C. corniculata, F. v. M., fragm. X, 62 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,62.
C. acuaria, F. v. M., fragm. VI, 16 (1868)...	W.A.	—	—	—	—	—	—	B.fl.VI,202	M.fr.VI,16&250.

PIPERACEAE.

L. Cl. Richard in Humb. Bonpl. & Kunth, Nov. gen. I, 46 (1815).

PIPER, Linné, gen. pl. 333 (1737), from C. Bauhin (1623). (Macropiper.)

P. subpeltatum, Willdenow, spec. plant. V, 166 (1810)	—	—	—	—	—	Q.	—	B.fl.VI,204	
P. excelsum, G. Forster, florul. ins. Austr. prodr. 5 (1786) ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,204	
P. Novae Hollandiae, Miquel in Medd. Akad. Vetensk. Amsterd. sec. ser. II, 8 (1866)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,204	
P. Banksii, Miquel in Medd. Akad. Amst. sec. ser. II, 9 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,205	
P. triandrum, F. v. M., fragm. V, 197 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,205	M.fr.V,197.
P. hederaceum, Cunningham in De Cand. prodr. XVI, 366 (1869)	—	—	—	—	N.S.W.	—	—	B.fl.VI,205	

PEPEROMIA, Ruiz & Pavon, fl. Peruv. & Chil. prodr. 8, t 2 (1794).

P. leptostachya, Hooker & Arnott, Bot. Beech. 96 (1841) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,206	M.fr.IX,76.
P. reflexa, A. Dietrich, spec. plant. I, 180 (1831)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,206	
P. Urvilleana, Ach. Richard in d'Urville, voy. Bot. 356 (1832)...	—	—	—	—	—	—	—	—	M.fr.IX,76.

PODOSTEMONEAE.

L. Cl. Richard in Humb. Bonpl. & Kunth, Nov. gen. I, 246 (1815).

Genus and species not yet determined	—	—	—	—	—	Q.	—	—	

NEPENTHACEAE.

Reichenbach, Conspectus regni vegetabilis 45 (1828).

NEPENTHES, Linné, syst. nat. 9 (1735); Linné, gen. pl. 273 (1737).

N. Kennedyana, F. v. M., fragm. V, 154 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,40	M.fr.V,154.

ARISTOLOCHIEAE.

A. L. de Jussieu, in Ann. du Mus. V, 221 (1804).

ARISTOLOCHIA, Tournefort, inst. 162, t. 71 (1700), from Hippocrates, Theophrastos & Dioscorides.

A. deltantha, F. v. M., fragm. VI, 179 (1869)	—	—	—	—	—	Q.	—	B.fl.VI,208	M.fr.VI,179&255.
A. praevenosa, F. v. M., fragm. II, 166 (1861)	—	—	—	—	N.S.W.	—	—	B.fl.VI,208	M.fr.VI,180.
A. pubera, R. Brown, prodr. 349 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,208	
A. Thozetii, F. v. M., fragm. II, 167 (1861)	—	—	—	—	—	Q.	—	B.fl.VI,208	M.fr.II,167.
A. Indica, Linné, spec. plant. 960 (1753)	—	—	—	—	—	Q.	—	B.fl.VI,209	

BALONOPHOREAE.

L. Cl. Richard in Mém. du Mus. VIII, 404 & 429 (1822).

BALONOPHORA, R. & G. Forster, char. gen. 99, t. 50 (1776).

B. fungosa, R. & G. Forster, char. gen. 99, t. 50 (1776)	—	—	—	—	—	Q.	—	B.fl.VI,232	

VINIFERAE.

J. de St. Hilaire, Exp. fam. II, 48, t. 79 (1805).

VITIS, Tournefort, inst. 613, t. 384 (1700), e Latinis. (Cissus.)

V. Baudiniana, F. v. M., fragm. IV, 136 (1864)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,447	M.fr.V,212;VI,178.
V. oblonga, Bentham, Fl. Austr. I, 447 (1863)	—	—	—	—	—	Q.	—	B.fl.I,447	M.fr.VI,178;IX,125.
V. cordata, Wallich in Bentham's flora Hongk. 54 (1861) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,447	M.fr.II,73;VI,178;IX,126.
V. adnata, Wallich in Wight & Arnott, prodr. 126 (1834) ...	—	—	—	—	—	—	N.A.	B.fl.I,448	M.fr.VI,178.
V. nitens, F. v. M., fragm. II, 73 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,448	M.fr.VI,178.
V. brachypoda, F. v. M., fr. IX, 125 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,125.
V. saponaria, Seemann, Bonplandia, 254 (1859)	—	—	—	—	—	Q.	—	B.fl.I,448	M.fr.VI,178.
V. acris, F. v. M., fragm. II, 75 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,449	M.fr.VI,178.
V. trifolia, Linné, spec. plant. 203 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,449	M.fr.II,75;VI,177;IX,126.
V. clematidea, F. v. M., fragm. II, 74 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,449	M.fr.VI,177;IX,126.
V. acetosa, F. v. M., plant. of Vict. I, 94 (1860)	—	—	—	—	—	Q.	N.A.	B.fl.I,449	M.fr.VI,177;IX,126.
V. hypoglauca, F. v. M., pl. of Vict. I, 95 (1860)...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,450	M.fr.VI,177.
V. sterculifolia, F. v. M. in Benth. Fl. Austr. I, 450 (1863) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,450	M.fr.IX,126.
V. penninervis, F. v. M., fr. VI, 177 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VI,177.

V. opaca, F. v. M. in Benth. Fl. Austr. I, 450 (1863) ... ... — — — — N.S.W. Q. — B.fl.I,450 M.fr.VI,177;IX,125.
V. augustissima, F. v. M., fragm. I, 141 (1859) ... ... W.A. — — — — — — B.fl.I,450 M.fr.I,244; II,180; IX,126.

LEEA, Linné, mantiss. 17 (1767).
L. staphylea, Roxburgh, hort. Beng. 18 (1814) ... ... — — — — — Q. — B.fl.I,415 M.fr.VI,178.
L. aculeata, Blume, Bijdrag. 197 (1825) ... ... — — — — — Q. N.A. — M.fr.VI,178.

SAPINDACEAE.

A. L. de Jussieu, Gen. 246 (1789).

CARDIOSPERMUM, Linné, syst. nat. (1735); Linné, gen. pl. 117 (1737).
C. Halicacabum, Linné, sp. pl. 366 (1753) ... ... — — — — — Q. N.A. B.fl.I,453

GANOPHYLLUM, Blume, Mus. Bot. Lugd. I, 230 (1850).
G. falcatum, Blume, Mus. Bot. Lugd. I, 230 (1850) ... ... — — — — — Q. — — M.fr.VII,24;IX,193.

ATALAYA, Blume, Rumphia III, 186 (1847). (Pseudatalaya, Sapindus partly.)
A. coriacea, Radlkofer in Sitzungsber. d. Akad. zu Münch. 326 (1878) — — — — N.S.W. — — —
A. salicifolia, Blume, Rumphia III, 186 (1847) ... ... — — — — — — N.A. B.fl.I,463
A. australis, Radlk. in Sitzungsber. der Akad. zu München. 327 (1878) — — — — — Q. — B.fl.I,464
A. multiflora, Bentham, Fl. Austr. I, 463 (1863) ... ... — — — — N.S.W. Q. — B.fl.I,463 M.fr.IX,90.
A. hemiglauca, F. v. M. in Benth. Fl. Austr. I, 463 (1863) ... — S.A. — — N.S.W. Q. N.A. B.fl.I,463 M.fr.IX,90.
A. variifolia, F. v. M. in Benth. Fl. Austr. I, 463 (1863) ... — — — — — — N.A. B.fl.I,463 M.fr.I,45.

DIPLOGLOTTIS, J. Hooker in H. & B. gen. pl. 395 (1862).
D. Cunninghamii, J. Hooker in H. & B. gen. pl. 395 (1862) ... — — — — N.S.W. Q. — B.fl.I,454 M.fr.V,145;IX,90.

ERIOGLOSSUM, Blume, Bijdr. 229 (1829). (Pancovia partly.)
E. edule, Blume, Bijdr. 229 (1829) ... ... — — — — — — N.A. B.fl.I,454 M.fr.IX,100.

CASTANOSPORA, F. v. Mueller, fragm. IX, 92 (1875).
C. Alphandi, F. v. M., fragm. IX, 92 (1875) ... ... — — — — — Q. — — M.fr.IV,158;IX,92,197.

ALLOPHYLUS, Linné, amoen. acad. 124 (1747). (Schmidelia.)
A. ternatus, Loureiro, Fl. Cochinch. ed. Willd. 286 (1790) ... — — — — — Q. N.A. B.fl.I,455 M.fr.IX,90.

CUPANIA, Plumier, Gen. 45, t. 19 (1703). (Guioa, Ratonia, Aryteria, Elattostachys, Mischocarpus, Euphoria partly.)
C. Wadsworthii, Harpullia, F. v. M., fragm. IV, 1, t. XXVI (1863) — — — — — — Q. — — M.fr.IV,1;IX,89,197.
C. anacardioides, A. Richard, sert. Astrol. 33, t. 13 (1833) ... — — — — N.S.W. Q. N.A. B.fl.I,458 M.fr.IX,91.
C. Robertsonii, F. v. M., fragm. V, 146 (1866) ... ... — — — — — Q. — — M.fr.V,146;IX,94.
C. serrata, F. v. M., fragm. III, 43 (1862)... ... ... — — — — N.S.W. Q. — B fl.I,458 M.fr.III,43;IX,94.
C. foveolata, F. v. M., fragm. IX, 95 (1875) ... ... — — — — — Q. — — M.fr.IX,95.
C. tomentella, F. v. M. in Benth. Fl. Austr. I, 458 (1863) ... — — — — — Q. — B.fl I,458
C. pseudorhus, A. Richard, sert. Astrol. 34, t. 14 (1833) ... — — — — N.S.W. Q. — B.fl.I,459 M.fr.IX,92.
C. xylocarpa, Cunningham in Benth. Fl. Austr. I, 459 (1863) ... — — — — N.S.W. Q. — B.fl.I,459 M.fr.IX,95.
C. erythrocarpa, F. v. M., fragm. V, 7 (1865) ... ... — — — — — Q. — — M.fr.V,7;IX,91.
C. Mortoniana, F. v. M., fragm. V, 177 (1866) ... ... — — — — — Q. — — M.fr.V,177;IX,94.
C. lachnocarpa, F. v. M., fragm. IV, 157 (1864) ... ... — — — — — Q. — — M.fr.IV,156;IX,91.
C. grandissima, F. v. M., fragm. IV, 156 (1864) ... ... — — — — — Q. — — M.fr.IV,156;IX,91.
C. pyriformis, F. v. M., App. Rep. Intercol. Exhib. 25 (1867)... — — — — N.S.W. Q. — B.fl.I,461 M.fr.I,2;II,76;IX,90.
C. anodonta, F. v. M., fragm. II, 76 (1860) ... ... — — — — — Q. — B.fl.I,461 M.fr.I,2;II,76;IX,90.
C. stipitata, F. v. M., fragm. II, 75 (1860) ... ... — — — — N.S.W. — — B.fl.I,461 M.fr.II,75;IX,91.
C. punctulata, F. v. M., fragm. III, 12 (1862) ... ... — — — — — Q. — B.fl.I,458 M.fr.III,12;IX,91.
C. exangulata, F. v. M., fragm. IV, 156 (1864) ... ... — — — — — Q. — — M.fr.IV,156;IX,91.
C. O'Shanesiana, F. v. M., fragm. IX, 96 (1875) ... ... — — — — — Q. — — M.fr.IX,96.
C. Daemeliana, F. v. M., fragm. IX, 96 (1875) ... ... — — — — — Q. — — M.fr.IX,96.
C. Martyana, F. v. M., fragm. V, 6 (1865) ... ... — — — — — Q. — — M.fr.V,6;IX,94.
C. Cordieri, F. v. M., fragm. IX, 93 (1875) ... ... — — — — — Q. — — M.fr.IX,93.
C. tenax, Cunningham in Benth. Fl. Austr. I, 461 (1863) ... — — — — N.S.W. Q. — B.fl.I,461 M.fr.IX,94.
C. semiglauca, F. v. M. in Benth. Fl. Austr. I, 457 (1863) ... — — — — N.S.W. Q. — B.fl.I,457 M.fr.IX,97.

NEPHELIUM, Linné, mantiss. 18 (1767). (Euphoria partly, Alectryon, Spanoghea.)
N. connatum, F. v. M. in Benth. Fl. Austr. I, 465 (1863) ... — — — — N.S.W. Q. — B.fl.I,465 M.fr.IX,98.
N. semicinereum, F. v. M., fragm. IV, 158 (1864) ... ... — — — — — Q. — — M.fr.IV,158.
N. subdentatum, F. v. M. in Benth. Fl. Austr. I, 465 (1863) ... — — — — N.S.W. — — B.fl.I,465 M.fr.IX,99.
N. tomentosum, F. v. M. in Transact. Vict. Inst. II, 64 (1858) — — — — N.S.W. Q — B.fl.I,466 M.fr.IX,99.
N. coriaceum, Bentham, Fl. Austr. I, 466 (1863)... ... ... — — — — — Q. — B.fl.I,466
N. foveolatum, F. v. M. in Benth. Fl. Austr. I, 466 (1863) ... — — — — — Q. — B.fl.I,466 M.fr.IX,99.
N. leiocarpum, F. v. M. in Transact. phil. Inst. Vict. III, 25 (1858) — — — V. N.S.W. Q. — B.fl.I,467 M.fr.IX,98.
N. Beckleri, Bentham, Fl. Austr. I, 467 (1863) ... ... — — — — N.S.W. — — B.fl.I,467
N. divaricatum, F. v. M. in Benth. Fl. Austr. 467 (1863) ... — — — — — Q. — B.fl.I,467 M.fr.IX,98.
N. microphyllum, Bentham, Fl. Austr. I, 468 (1863) ... ... — — — — — Q. — B.fl.I,468
N. Leichhardtii, F. v. M., App. Rep. Intercol. Exhib. 25 (1867) — — — — — Q. — B.fl.I,468 M.fr.IX,99.
N. distyle, F. v. M., fragm. IX, 99 (1875)... ... ... — — — — — Q. — B.fl.I,462 M.fr.IX,99.

HETERODENDRON, Desfontaines in Mém. du Mus. IV, 8, t. 3 (1818).
H. oleaefolium, Desfontaines in Mém. du Mus. IV, 8, t. 3 (1818) W.A. S.A. — V. N.S.W. Q. N.A. B.fl.I,469 M.fr.IX,90;X,82.
H. diversifolium, F. v. M., fragm. I, 46 (1858) ... ... — — — — N.S.W. Q. — B.fl.I,469 M.fr.I,46;IX,90;X,82.

HARPULLIA, Roxburgh, Fl. Ind. ed. Carey II, 44 (1826).
H. alata, F. v. M., fragm. II, 103 (1861) ... ... — — — N.S.W. Q. — B.fl.I,470 M.fr.II,103;IX,89.
H. Hillii, F. v. M. in Transact. Vict. Inst. III, 26 (1859) ... — — — — N.S.W. Q. — B.fl.I,470
H. Leichhardtii, F. v. M. in Benth. Fl. Austr. I, 470 (1863) ... — — — — — — N.A. B.fl.I,470
H. pendula, Planchon in Transact. Vict. Inst. III, 26 (1859) ... — — — — N.S.W. Q. — B.fl.I,471 M.fr.IX,89&187.

AKANIA, J. Hooker in B. & H. Gen. pl. 409 (1862).
A. Hillii, J. Hooker in B. & H. Gen. pl. 409 (1862) ... ... — — — — N.S.W. Q. — B.fl.I,471 M.fr.III,44;IX,89&197.

DIPLOPELTIS, Endlicher in Huegel. enum. 13 (1837).

D. Huegelii, Endlicher in Hueg. enum. 13 (1837)	W.A.	—	—	—	—	—	N.A.	B.fl.I,456	M.fr.III,12;IX,89.
D. Stuartii, F. v. M., fragm. III, 12 (1862)	—	S.A.	—	—	—	—	N.A.	B.fl.I,456	M.fr.III,12&167;IX,89.

DODONAEA, Linné, gen. pl. 341 (1737). (Empleurosma).

D. triquetra, Wendland, Bot. Beobacht. p. 44 (1798)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,474	M.fr.I,73;IX,88.
D. lanceolata, F. v. M., fragm. I, 73 (1858)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,475	M.fr.I,73.
D. petiolaris, F. v. M., fragm. III, 13 (1862)	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.I,475	M.fr.IX,89.
D. viscosa, Linné, mantissa X, altera 149 & 228 (1771) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,475	M.fr.IX,80.
D. peduncularis, Lindley in Mitch. Trop. Austr. 361 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,478	M.fr.IX,87.
D. procumbens, F. v. M. in Transact. Vict. Inst. I, 3 (1854) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,478	
D. ericifolia, G. Don, gen. syst. I, 674 (1831)	—	—	T.	—	—	—	—	B.fl.I,478	M.fr.IX,87.
D. filifolia, Hooker in Mitch. Trop. Austr. 241 (1848)	—	—	—	—	—	Q.	—	B.fl.I,478	M.fr.I,71.
D. lobulata, F. v. M. in Linnaea XXV, 372 (1852)	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.I,479	M.fr.IX,87.
D. ptarmicifolia, Turcz. in Bull. Soc. Mosc. XXV, part III, 165(1852)	W.A.	—	—	—	—	—	—	B.fl.I,479	M.fr.I,97;IX,87.
D. truncatiales, F. v. M., fragm. II, 143 (1861)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,479	M.fr.IX,89.
D. platyptera, F. v. M., fragm. I, 73 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,480	M.fr.IX,87.
D. stenophylla, F. v. M., fragm. I, 72 (1858)	—	—	—	—	—	Q.	—	B.fl.I,480	M.fr.IX,87.
D. pinifolia, Miquel in Lehm. pl. Preiss. I, 227 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,480	M.fr.IX,88.
D. ceratocarpa, Endlicher in Hueg. enum. 13 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,481	M.fr.X,88.
D. divaricata, Bentham, Fl. Austr. I, 481 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,481	M.fr.IX,87.
D. triangularis, Lindley in Mitch. Trop. Austr. 219 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,481	M.fr.IX,87.
D. aptera, Miquel in Lehm. pl. Preiss. I, 225 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,481	M.fr.IX,88.
D. bursarifolia, Behr & F. v. M. in Transact. Vict. Inst. I, 8 (1855)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,482	M.fr.IX,87.
D. Baueri, Endlicher in Hueg. enum. 13 (1837)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,482	M.fr.IX,87.
D. trifida, F. v. M., fragm. IX, 88 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,88.
D. humifusa, Miquel in Lehm. pl. Preiss. I, 266 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,482	M fr.IX,88.
D. hexandra, F. v. M. in Transact. Vict. Inst. I, 117 (1855) ...	—	S.A.	—	—	—	—	—	B.fl.I,483	
D. ericoides, Miquel in Lehm. pl. Preiss. I, 227 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,483	M.fr.IX,88.
D. polyzyga, F. v. M. in fragm. I, 74 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,483	M.fr.I,74.
D. megazyga, F. v. M. in Benth. Fl. Austr. I, 483 (1863) ...	—	—	—	—	N.S.W.	—	—	B.fl.I,483	M.fr.IX,86.
D. macrozyga, F. v. M., fragm. IV, 135 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IX,86.
D. physocarpa, F. v. M., fragm. I, 74 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,484	M.fr.I,74;IX,86.
D. vestita, Hooker in Mitch. Trop. Austr. 265 (1848)	—	—	—	—	—	Q.	—	B.fl.I,484	M.fr.IX,86.
D. pinnata, Smith in Rees' Cycl. XII (1809)	—	—	—	—	N.S.W.	—	—	B.fl.I,484	
D. oxyptera, F. v. M., fragm. I, 74 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,484	M.fr.I,74.
D. humilis, Endlicher, nov. stirp. dec. 26 (1839)	—	S.A.	—	—	—	—	N.A.	B.fl.I,485	M.fr.IX,86.
D. Macrossani, F. v. M. & Scortechini in Melb. Chemist Jan. 1882	—	—	—	—	—	Q.	—	—	M.fr.XII.
D. microzyga, F. v. M. annual Rep. bot. Gard. 12 (1863) ...	—	—	—	—	—	—	—	—	M.fr.IX,86.
D. boronifolia, G. Don, gen. syst. I, 674 (1831)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,485	M.fr.IX,86.
D. multijuga, G. Don, gen. syst. I, 674 (1831)	—	—	—	—	N.S.W.	—	—	B.fl.I,485	M.fr.I,219;II,182;IX,86.
D. larraeoides, Turcz. in Bull. Soc. Mosc. XXXI, 408 (1858) ...	W.A.	—	—	—	—	—	—	B.fl.I,486	
D. inaequifolia, Turcz in Bull. Soc. Mosc. XXXI, 408 (1858) ...	W.A.	—	—	—	—	—	—	B.fl.I,486	M.fr.I,219;IX,89.
D. adenophora, Miquel in Linnaea XVIII, 95 (1845)	W.A.	—	—	—	N.S.W.	Q.	—	B.fl.I,486	M.fr.I,98.
D. tenuifolia, Lindley in Mitch. Trop. Austr. 248 (1848) ...	—	—	—	—	—	—	—	—	M.fr.IX,85.
D. stenozyga, F. v. M., fragm. I, 98 (1859)	W A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,486	M.fr.IX,87.
D. concinna, Bentham, Fl. Austr. I, 487 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,487	M.fr.IX,87.

DISTICHOSTEMON, F. v. Mueller in Hook. Kew misc. IX, 306 (1857).

D. phyllopterus, F. v. M. in Hook. Kew misc. IX, 306 (1857) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,487	M.fr.IX,89.

BLEPHAROCARYA, F. v. Mueller, fragm. XI, 15 (1878).

B. involucrigera, F. v. M., fragm. XI, 16 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,16&137

MALPIGHIACEAE.

Ventenat, Tabl. III, 131 (1799).

RYSSOPTERYS, Blume in Delessert ic. sel. III, 21, t. 35 (1837).

R. Timorensis, Blume in Deless. ic. sel. III, 21, t. 35 (1837) ...	—	—	—	—	—	Q.	—	B.fl.I,285	

TRISTELLATEIA, Petit-Thouars in Roemer. Collect. bot. 206 (1809).

T. Australasica, A. Richard, Sert. Astrol. 38, t. 15 (1833) ...	—	—	—	—	—	Q.	—	B.fl.I,286	

BUESERACEAE.

Kunth in Ann. des sc. nat. II, 346 (1824).

GARUGA, Roxburgh, Pl. of Coromandel III, 5, t. 208 (1819).

G. floribunda, Descaisne, in nouv. annal. du Mus. III, 477 (1834)	—	—	—	—	—	—	N.A.	B.fl.I,377	

CANARIUM, Rumphius, herb. Amboin. II, 145, t. 47, 49, 54 (1741). (Sonzaya.)

C. Australianum, F. v. M., fragm. III, 15 (1860)	—	—	—	—	—	Q.	N.A.	B.fl.I,378	M.fr.VII,25.

ANACARDIACEAE.

R. Brown in Tuckey's Expedit. Congo 431 (1818).

RHUS, Tournefort, Inst. 611, t. 381 (1700), from Hippocr. Theophr. Diosc. & Plin. (Rhodosphaera.)

R. rhodanthema, F. v. M. in Journ. pharm. soc. Vict. pl. I, 43 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,489	M.fr.VII,23.
R. panaciformis, F. v. M., fragm. VII, 22 (1869)	—	—	—	—	—	Q.	—	—	M.fr.VII,22.

EUROSCHINUS, J. Hooker in B. & H. Gen. pl. I, 422 (1862).

E. falcatus, J. Hooker in B. & H. Gen. pl. I, 422 (1862) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,490	M.fr.VII,24.

BUCHANANIA, Sprengel in Schrader's Journ. II, 234 (1800).

B. angustifolia, Roxburgh, Pl. Corom. III, 68, t. 262 (1819) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,490	M.fr.VII,23.
B. mangoides, F. v. M., fragm. VII, 23 (1869)	—	—	—	—	—	Q.	—	—	M.fr.VII,23.
B. arborescens, Blume, Mus. Bot. Lugd. Batav. I, 183 (1850) ...	—	—	—	—	—	Q.	—	—	M.fr.VII,23.

SEMECARPUS, Linné fil., suppl. 25 (1781).

S. Anacardium, Linné fil., suppl. 182 (1781)	—	—	–	–	—	Q.	N.A.	B.fl.I,491	M.fr.VII,23.

SPONDIAS, Linné, gen. pl. 365 (1737).

S. Solandri, Bentham, Fl. Austr. I, 492 (1863)	—	—	—	—	—	Q.	—	B.fl.I,492	
S. pleiogyna, F. v. M., fragm. IV, 78 (1864)	—	—	—	—	—	Q.	—	—	M.fr.III,15.

CELASTRINEAE.

R. Brown in App. Flind. Voy. 554 (1814).

EUONYMUS, Tournefort, inst. 617, t. 388 (1700), from Theophr. & Plinius. (Evonymus.)

E. Australianus, F. v. M., fragm. IV, 118 (1864)...	—	—	—	—	—	Q.	—	—	M.fr.IV 118;VI,202.

CELASTRUS, Linné, gen. pl. 59 (1737).

C. Australis, Harvey & F. v. M. in Trans. phil. Soc. Vict. I, 41 (1854)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,398	M.fr.III,93.
C. Muelleri, Bentham, Fl. Austr. I, 399 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,399	
C. dispermus, F. v. M. in Transact. phil. Inst. Vict. III, 31 (1859)	—	—	—	—	—	Q.	—	B.fl.I,399	M.fr.VI,203.
C. bilocularis, F. v. M. in Transact. phil. Inst. Vict. III, 31 (1859)	—	—	—	—	—	Q.	—	B.fl.I,399	M.fr.VI,204.
C. Cunninghamii, F. v. M. in Trans. phil. Inst. Vict. III, 30 (1859)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,399	M.fr.VI,203.

GYMNOSPORIA, Wight & Arnott, prodr. fl. paenius. Ind. I, 159 (1834.)

G. montana, Wight & Arnott, prodr. I, 159 (1834)	—	—	—	—	—	Q.	—	B.fl.I,400	

LEUCOCARPUM, A. Richard, sert. Astrolabe 46 (1834). (Denhamia, Leucocarpon.)

L. Oleaster, F. v. M. in Transact. Roy. Soc. of N.S.W. 15 (1881)	—	—	—	—	—	Q.	—	B.fl.I,401	M.fr.VI,203.
L. obscurum, A. Richard, sert. Astrol. 46, t. 18 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.I,401	M.fr.VI,203.
L. pittosporoides, F. v. M. in Trans. Roy. Soc. of N.S.W. 15 (1881)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,402	M.fr.VI,203.

ELAEODENDRON, J. F. Jacquin in Nov. Act. Helv. I, 36 (1787).

E. Australe, Ventenat, Jard. Malm. t. 117 (1804)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,402	M.fr.III,61;IV,175;VI,204.

CARYOSPERMUM, Blume, Mus. bot. Lugd. I, 175 (1850).

C. arborescens, F. v. M., fragm. VI, 202 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VI,202.

SIPHONODON, Griffith in Calc. journ. of Nat. Hist. IV, 150 (1843).

S. Australe, Bentham, Fl. Austr. I, 403 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,403	M.fr.VI,204.

HIPPOCRATEA, Linné, gen. pl. 363 (1737).

H. obtusifolia, Roxburgh, flor. Ind. edit. Carey, I, 170 (1820)...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,404	

SALACIA, Linné, mantiss. II, 159 (1771).

S. prinoides, De Candolle, prodr. I, 571 (1824)	—	—	—	—	—	Q.	—	—	M.fr.VI,202.

STACKHOUSIEAE.

R. Brown in App. to Flind. Voy. 555 (1814).

STACKHOUSIA, Smith in Transact. Linn. Soc. IV, 218 (1798). (Tripterococcus, Plokiostigma.)

S. pulvinaris, F. v. M. in Transact. phil. Soc. Vict. I, 101 (1855)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,405	M.fr.III,88.
S. linarifolia, Cunningham in Field's N.S.Wales 365 (1825) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,406	M.fr.III,87;XI.27.
S. pubescens, A. Richard, sert. Astrol. 89, t. 33 (1834)	W.A.	—	—	—	—	—	—	B.fl.I,407	
S. Huegelii, Endlicher in Hueg. enum. 17 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,407	
S. flava, Hooker, icon. pl. t. 269 (1840)	W.A.	S.A.	T.	V.	—	—	—	B.fl.I,407	
S. muricata, Lindley, Bot. Reg. XXII, 1917 (1837)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,408	M.fr.VIII,36.
S. viminea, Smith in Rees' Cycl. XXXIII (1816)...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.I,408	M.fr.VIII,36.
S. elata, F. v. M., fragm. III, 86 (1862)	W.A.	—	—	—	—	—	—	B.fl.I,408	M.fr.III,86.
S. scoparia, Bentham, Fl. Austr. I, 409 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,409	
S. spathulata, Sieber in Spreng. syst. cur. post. 124 (1827) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,406	M.fr.III,86.
S. megaloptera, F. v. M., fragm. VIII, 35 (1873)...	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,119.
S. Brunonis, Bentham, Fl. Austr. I, 409 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.I,409	M.fr.VIII,36.

MACGREGORIA, F. v. Mueller in Caruel, Giorn. Bot. Italian. 128 (1873).

M. racemigera, F. v. M. in Caruel, Giorn. Bot. Ital. 128 (1873)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,161;XI,134.

FRANKENIACEAE.

A. de St. Hilaire in Bull. de la Soc. Philom. 22 (1815).

FRANKENIA, Linné, gen. pl. 92 (1737).

F. laevis, Linné, spec. plant. 331 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,151	
F. bracteata, Turczan. in Bull. Soc. Mosc. XXVII, 367 (1854)...	W.A.	—	—	—	—	—	—	B.fl.I,150	
F. glomerata, Turczan. in Bull. Soc. Mosc. XXVII, 368 (1854)	W.A.	—	—	-	—	—	—	B.fl.I,151	
F. parvula, Turczan. in Bull. Soc. Mosc. XXVII, 368 (1854) ...	W.A.	—	—		-	—	—	B.fl.I,152	
F. Drummondii, Bentham, Fl. Austr. I, 152 (1863)	W.A.	—	-	-	-	—	—	B.fl.I,152	
F. tetrapetala, Labillard., nov. Holl. pl. spec. I, 88, t. 114 (1804)	W.A.	—	—	-	—	—	—	B.fl.I,152	
F. punctata, Turczan. in Bull. Soc. Mosc. XXVII, 367 (1854) ...	W.A.	—	—	—	—	—	—	B.fl.I,153	

PLUMBAGINEAE.

A. L. de Jussieu, gen. plant. 92 (1789).

PLUMBAGO, Tournefort, inst. 140, t. 58 (1700).

P. Zeilanica, Linné, sp. pl. 151 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,267	M.fr.XI,7.

STATICE, Linné, gen. pl. 88 (1737).

S. Taxanthema, Roemer & Schultes, syst. veg. VI, 798 (1820)...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,267	M.fr.XI,7.
S. salicorniacea, F. v. M., fragm. XI, 7 (1878)	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,7.

AEGIALITIS, R. Brown, prodr. 426 (1810). (Aegialinites).

A. annulata, R. Brown, prodr. 426 (1810)	—	—	–	—	—	Q.	—	B.fl.IV,266	

PORTULACEAE.

A. L. de Jussieu, Gen. 312 (1789).

PORTULACA, Tournefort, inst. 236, t. 118 (1700), from l'Obel (1581).

P. oleracea, Linné, spec. plant. 445 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,169	M.fr.X,71.
P. napiformis, F. v. M. in Benth. Fl. Austr. I, 169 (1863) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,169	M.fr.X.72.
P. australis, Endlicher, Atakta 7, t. 6 (1833)	—	—	—	—	—	Q.	N.A.	B.fl.I,169	M.fr.X,72.
P. filifolia, F. v. M., fragm. I, 169 (1859)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,169	M.fr.III,164;X,72.
P. digyna, F. v. M., fragm. I, 170 (1859)	—	—	—	—	—	—	N.A.	B.fl.I,170	M.fr.I,170.
P. oligosperma, F. v. M., fragm. I, 170 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.I,170	M.fr.III,164;X,72.
P. bicolor, F. v. M., fragm. I, 171 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.I,170	M.fr.III,164;X,72.
P. Armitii, F. v. M., fragm. X, 97 (1877)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,97.

CLAYTONIA, Gronovius in Linné, gen. pl. 339 (1737). (Calandrinia, Talinum.)

C. spergularina; Calandrinia, F. v. M., fragm. I, 175 (1859) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,176	M.fr.I,175;X,71.
C. ptychosperma; Calandrinia, F. v. M., fragm. IV, 137 (1864)	—	S.A.	—	—	—	Q.	—	—	M.fr.IV,137;X,70
C. strophiolata, F. v. M., fragm. XI, 82 (1880)	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,82.
C. Lehmanni, F. v. M.; Calandr., Endl. in Lehm. pl. Preiss. II, 235 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,172	M.fr.X,71.
C. uniflora, F. v. M.; Calandr. in Trans. Inst. Vict. III, 41 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.I,172	M.fr.X,71.
C. Balonnensis, F. v. M.; Calandrinia, Lindley in Mitch. Trop. Austr. 148 (1848)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,172	M.fr.X,71.
C. polyandra, F. v. M.; Calandr., Benth. Fl. Austr. I, 172 (1863)	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.I,172	M.fr.X,71.
C. pleiopetala; Calandrinia, F. v. M., fragm. X, 70 (1877) ...	—	S.A.	—	—	—	Q.	—	—	M.fr.X,70.
C. quadrivalvis; Calandrinia, F. v. M., fragm. I, 176 (1859) ...	—	—	—	—	—	—	N.A.	B.fl.I,173	M.fr.I,176.
C. liniflora, F. v. M.; Calandr., Fenzl. in Hueg. enum. 52 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,173	M.fr.X,71.
C. gracilis, F. v. M.; Calandrinia, Benth. Fl. Austr. I, 173 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,173	
C. polypetala, F. v. M.; Calandr., Fenzl. in Hueg. enum. 51 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,174	
C. Pickeringi, F. v. M.; Calandr., A. Gray, Bot. Wilkes' Exped. 144 (1854)	—	—	—	—	N.S.W.	—	—	—	M.fr.X,69;XI,136.
C. volubilis, F. v. M.; Calandr., Benth. Fl. Austr. I, 174 (1863)	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,174	M.fr.X,68;XI,27.
C. calyptrata, F. v. M., fragm. III, 89 (1862)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,174	M.fr.IV,175;X,68.
C. pumila; Calandrinia, F. v. M., fragm. X, 68 (1877)	W.A.	S.A.	—	—	—	Q.	—	—	M.fr.X,68.
C. composita, F. v. M.; Calandrinia, Nees in Lehm. pl. Preiss. I, 247 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,175	M.fr.X,69.
C. corrigioloides; Calandr., F. v. M., native pl. of Vict. I, 135 (1879)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,175	M.fr.X,69.
C. brevipedata; Calandr., F. v. M., fragm. X, 69 (1877) ...	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,69;XI,136.
C. granulifera, F. v. M.; Calandr., Benth. Fl. Austr. I, 176 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,176	
C. pogonophora; Calandrinia, F. v. M., fragm. X, 69 (1877) ...	—	—	—	—	—	Q.	—	—	M.fr.X,69.
C. pygmaea, F. v. M., fragm. III, 89 (1862)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,176	M.fr.I,175;X,71.
C. Australasica, J. Hooker, icon. plant. t. 293 (1840)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,177	M.fr.III,89;X,71.

MONTIA, Micheli, nov. pl. gen. 17, t. 13 (1729).

M. fontana, Linné, spec. plant. 87 (1753)	—	—	T.	—	—	—	—	B.fl.I,177	

CARYOPHYLLEAE.

Linné, philosoph. bot. 31 (1751).

STELLARIA, Linné, spec. plant. 421 (1753).

S. pungens, Brongniart, Voy. sur la Coquille t. 78 (1826) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,157	
S. glauca, Withering, Arr. veg. in Great Brit. I, 420 (1776) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,158	
S. flaccida, Hooker, Comp. Bot. Mag. I, 275 (1835)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,158	
S. multiflora, Hooker, Comp. Bot. Mag. I, 275 (1835)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,159	M.fr.XI,27.

SAGINA, Linné, syst. nat. 8 (1735); Linné, gen. pl. 118 (1737).

S. procumbens, Linné, spec. plant. 128 (1753)	—	—	—	V.	N.S.W.	—	—	B.fl.I.160	
S. apetala, Linné, mantiss. 559 (1771)	—	S.A.	T.	V.	—	—	—	B.fl.I,160	

COLOBANTHUS, Bartling, Reliq. Haenk. II, 13, t. 49 (1830).

C. Benthamianus, Fenzl in Ann. Wien. Mus. I, 49 (1838) ...	—	—	—	V.	N.S.W.	—	—	B.fl.I,160	
C. Billardieri, Fenzl in Ann. des Wien. Mus. I, 48 (1838) ...	—	S.A.	T.	V.	—	—	—	B.fl.I,161	

SCLERANTHUS, Linné, gen. plant. 130 (1737). (Mniarum.)

S. pungens, R. Brown, prodr. 412 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,260	
S. diander, R. Brown, prodr. 412 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.V,260	
S. biflorus, J. Hooker, Fl. N. Zeal. I, 74 (1853)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.V,259	
S. mniaroides, F. v. M., pl. of Vict. I, 215 (1862)	—	—	—	V.	N.S.W.	—	—	B.fl.V,259	

SAPONARIA, Linné, gen. plant. 130 (1737), from Camerarius (1586). (Gypsophila.)

S. tubulosa, F. v. M., native pl. of Vict. I, 136 (1879)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,155	

SPERGULARIA, Persoon, synops. pl. I, 504 (1805). (Lepigonum, Arenaria partly.)

S. rubra, Cambessèdes in St. Hilaire, Fl. Brazil. II, 179 (1829) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,162	M.fr.XI,27

DRYMARIA, Willdenow in Roemer & Schultes, syst. V, 406 (1819).

D. filiformis, Bentham, Fl. Austr. I, 162 (1863)	W.A.	S.A.	—	—	—	—	—	B.fl.I,162	

POLYCARPON, Loefling, iter Hispanic. 7 (1758).

P. tetraphyllum, Loefling in Linné, syst. nat. ed. X, 881 (1759)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,163	

POLYCARPAEA, Lamarck in Journ. d'hist. nat. II, 8, t. 25 (1792). (Aylmeyera.)

P. longiflora, F. v. M., Rep. of Babb. Exped. 8 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,164	
P. spirostylis, F. v. M., Rep. of Babb. Exped. 8 (1858) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,165	
P. synandra, F. v. M., Rep. of Babb. Exped. 8 (1858)	—	S.A.	—	—	—	Q.	N.A.	B.fl.I,165	
P. violacea, Bentham, Fl. Austr. I, 165 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,165	
P. Indica, Lamarck in journ. d'hist. nat. II, 8 (1792)		S.A.	—	—	—	Q.	N.A.	B.fl.I,166	

P. breviflora, F. v. M., Rep. of Babb. Exped. 9 (1858) ... ... — — — — — Q. N.A. B.fl.I,166
P. spicata, Arnott in Ann. of nat. hist. III, 91 (1839) ... ... — — — — — — N.A. B.fl.I,167
P. involucrata, F. v. M., Rep. of Babb. Exped. 9 (1858) ... — — — — — — N.A. B.fl.I,167

AMARANTACEAE.

A. L. de Jussieu, Gen. 87 (1789).

GOMPHRENA, Linné, gen. plant. 69 (1737). (Philoxerus.)

G. canescens, R. Brown, prodr. 416 (1810) ... ... — — — — — — N.A. B.fl.V,253
G. flaccida, R. Brown, prodr. 416 (1810) ... ... — — — — — Q. N.A. B.fl.V,254 M.fr.III,123.
G. affinis, F. v. M. in Benth. Fl. Austr. V, 254 (1870) ... ... — — — — — — N.A. B.fl.V,254
G. humilis, R. Brown, prodr. 416 (1810) ... ... — — — — — Q. N.A. B.fl.V,254
G. Brownii, Moquin in De Cand. prodr. XIII, 2, 397 (1849) ... — — — — — Q. N.A. B.fl.V,255
G. brachystylis, F. v. M., fragm. III, 124 (1862) ... ... — — — — — — N.A. B.fl.V,255 M.fr.III,124.
G. leptoclada, Bentham, Fl. Austr. V, 255 (1870) ... ... — — — — — — N.A. B.fl.V,255
G. Maitlandi, F. v. M., fragm. III, 124, t. 23 (1862) ... ... — — — — — — N.A. B.fl.V,256 M.fr.VI,234.
G. pusilla, Bentham, Fl. Austr. V, 256 (1870) ... ... — — — — — — N.A. B.fl.V,256
G. tenella, Bentham, Fl. Austr. V, 256 (1870) ... ... — — — — — — N.A. B.fl.V,256
G. conica, Sprengel, syst. I, 824 (1825) ... ... — — — — — — N.A. B.fl.V,256 M.fr.III,125.
G. conferta, Bentham, Fl. Austr. V, 257 (1870) ... ... — — — — — Q. N.A. B.fl.V,257
G. diffusa, Sprengel, syst. I, 824 (1825) ... ... — — — — — — N.A. B.fl.V,257
G. parviflora, Bentham, Fl. Austr. V, 257 (1870) ... ... — — — — — — N.A. B.fl.V,257

ALTERNANTHERA, Forskael, flor. Aegypt. Arab. 28 (1775). (Telanthera.)

A. triandra, Lamarck, encycl. meth. I, 95 (1783) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.V,249
A. nana, R. Brown, prodr. 417 (1810) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.V,250
A. angustifolia, R. Brown, prodr. 417 (1810) ... ... — — — — — — N.A. B.fl.V,250
A. decipiens, Bentham, Fl. Austr. V, 251 (1870) ... ... — — — — — Q. — B.fl.V,251
A. polycephala, Bentham, Fl. Austr. V, 251 (1870) ... ... — — — — — — N.A. B.fl.V,251
A. leptophylla, Bentham, Fl. Austr. V, 251 (1870) ... ... — — — — — — N.A. B.fl.V,251
A. longipes, Bentham, Fl. Austr. V, 252 (1870) ... ... — — — — — — N.A. B.fl.V,252

ACHYRANTHES, Linné, gen. plant. 34 (1737).

A. aspera, Linné, spec. plant. 204 (1753) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.V,246
A. arborescens, R. Brown, prodr. 417 (1810) ... ... — — — — N.S.W. — — — M.fr.IX,169.

NYSSANTHES, R. Brown, prodr. 418 (1810).

N. erecta, R. Brown, prodr. 418 (1810) ... ... — — — — N.S.W. Q. — B.fl.V,247
N. diffusa, R. Brown, prodr. 418 (1810) ... ... — — — — N.S.W. Q. — B.fl.V,247

PTILOTUS, R. Brown, prodr. 415 (1810). (Trichinium, Psilotrichum, Goniotriche, Hemisteirus, Arthrotrichum.)

P. conicus, R. Brown, prodr. 415 (1810) ... ... — — — — — — N.A. B.fl.V,242
P. corymbosus, R. Brown, prodr. 415 (1810) ... ... — — — — — — N.A. B.fl.V,243 M.fr.III,125.
P. grandiflorus, F. v. M., fragm. I, 237 (1859) ... ... W.A. — — — — — — B.fl.V,243 M.fr.I,237.
P. spicatus, F. v. M. in Benth. Fl. Austr. V, 243 (1870) ... ... — — — — — Q. N.A. B.fl.V,243
P. Murrayi, F. v. M., fragm. III, 145 (1863) ... ... — S.A. — — — — N.A. B.fl.V,243 M.fr.III,145.
P. gomphrenoides, F. v. M., fragm. VI, 233 (1868) ... ... — — — — — — N.A. B.fl.V,244
P. latifolius, R. Brown in App. Sturt's Exped. 25 (1849) ... — S.A. — — — — N.A. B.fl.V,244 M.fr.VI,232.
P. macrotrichus, F. v. M., fragm. IV, 90 (1864) ... ... W.A. — — — — — N.A. B.fl.V,244 M.fr.VI,232.
P. villosiflorus, F. v. M., fragm. III, 125 (1863) ... ... — — — — — — N.A. B.fl.V,245 M.fr.III,125.
P. humilis, F. v. M., fragm. VI, 229 (1868) ... ... W.A. — — — — — — B.fl.V,245 M.fr.III,161; VI,229.
P. obovatus, F. v. M., fragm. VI, 228 (1868) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,220 M.fr.XI,27.
P. incanus, Poiret, Encycl. méth. suppl. IV, 620 (1817) ... — — — — — — N.A. B.fl.V,221 M.fr.VI,228.
P. parviflorus, Trich., Lindley in Mitch. three Exp. II, 13 (1838) ... — — — — N.S.W. Q. — B.fl.V,222
P. astrolasius, F. v. M., fragm. VI, 233 (1869) ... ... — — — — — — N.A. B.fl.V,222 M.fr.VI,233.
P. rotundifolius, F. v. M., fragm. VI, 230 (1868) ... ... — — — — — — N.A. B.fl.V,223 M.fr.III,122.
P. dissitiflorus, F. v. M., fragm. IV, 89 (1864) ... ... — — — — — — N.A. B.fl.V,223 M.fr.IV,89.
P. distans, Poiret, Encycl. suppl. IV, 620 (1817) ... ... — — — — — Q. N.A. B.fl.V,223 M.fr.VI,228.
P. alopecuroideus, F. v. M., fragm. VI, 227 (1868) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,224 M.fr.VI,227.
P. nobilis, F. v. M., fragm. VI, 227 (1868) ... ... — S.A. — V. N.S.W. — — B.fl.V,224
P. macrocephalus, Poiret, Encycl. suppl. IV, 620 (1817) ... — S.A. — V. N.S.W. Q. — B.fl.V,225 M.fr.VI,228; XI,27.
P. hemisteirus, F. v. M., fragm. IV, 90 (1864) ... ... W.A. S.A. — — — — N.A. B.fl.V,226 M.fr.VI,231.
P. parvifolius, F. v. M., fragm. VI, 229 (1869) ... ... — S.A. — — N.S.W. — — B.fl.V,241 M.fr.VI,229.
P. exaltatus, Nees in Lehm. pl. Preiss. I, 630 (1845) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,227
P. Manglesii, F. v. M., fragm. VI, 230 (1869) ... ... W.A. — — — — — N.A. B.fl.V,228 M.fr.VI,230.
P. Beckeri, F. v. M., fragm. VI, 233 (1869) ... ... — S.A. — — — — — B.fl.V,228
P. gomphrenoides, Moquin in De Cand., prodr. XIII, 2, 287 (1849) — S.A. — — — — — B.fl.V,229
P. esquamatus; Trichinium, Bentham, Fl. Austr. V, 229 (1870) W.A. — — — — — — B.fl.V,229
P. declinatus, Nees in Lehm. pl. Preiss. I, 631 (1845) ... ... W.A — — — — — — B.fl.V,230
P. erubescens, Schlechtendal in Linnaea XX, 575 (1847) ... — S.A. — V. N.S.W. — — B.fl.V,230 M.fr.VI,229.
P. divaricatus, F. v. M., fragm. VI, 229 (1868) ... ... W.A. — — — — — N.A. B.fl.V,231 M.fr.VI,229.
P. helipteroides, F. v. M., fragm. VI, 231 (1868) ... ... — S.A. — — — — N.A. B.fl.V,231 M.fr.III,122&161.
P. Stirlingi; Trichinium, Lindley in Bot. Reg. under n. 28 (1839) W.A. — — — — — N.A. B.fl.V,232
P. laxus; Trichinium, Bentham, Fl. Austr. V, 232 (1870) ... W.A. — — — — — — B.fl.V,232
P. axillaris, Trichinium, F. v. M. in Benth. Fl. Austr. V, 232 (1870) — — — — — — N.A. B.fl.V,232
P. striatus; Trichinium, Moquin in Benth. Fl. Austr. V, 233 (1870) — — — — — — N.A. B.fl.V,233
P. auriculifolius; Trichinium, Cunningham in De Cand., prodr. XIII, part II, 287 (1849) ... ... — — — — — — N.A. B.fl.V,233
P. sericostachyus, F. v. M., fragm. VI, 230 (1868) ... ... W.A. — — — — — — B.fl.V,234 M.fr.VI,230
P. roseus; Trichin., Moq. in De Cand., prodr. XIII, p. II, 284 (1849) W.A. — — — — — — B.fl.V,234
P. fusiformis, Poiret, Encycl. suppl. IV, 619 (1817) ... ... — — — — — — N.A. B.fl.V,234
P. gracilis, Poiret, Encycl. suppl. IV, 620 (1817) ... ... — — — — — — N.A. B.fl.V,235 M.fr.VI,228.
P. Drummondii, F. v. M., fragm. VI, 229 (1868) ... ... W.A. — — — — — — B.fl.V,235 M.fr.VI,229.

P. calostachyus, F. v. M., fragm. VI, 231 (1868)	—	—	—	—	—	—	N.A.	B.fl.V.236	M.fr.VI,231.
P. Fraseri ; Trichinium, Cunningham in De Cand., prodr. XIII, part II, 295 (1849)	—	—	—	—	N.S.W.	—	—	B.fl.V,236	
P. spathulatus, Poiret, Encycl. suppl. IV, 620 (1817)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,236	M.fr.VI,228.
P. pyramidatus, F. v. M., fragm. VI, 230 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,237	M.fr.VI,230.
P. holosericeus, F. v. M., fragm. VI, 229 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,237	M.fr.VI,229.
P. lanatus, Cunningham in De Cand., prodr. XIII, p. II, 281 (1849)	—	—	—	—	—	—	N.A.	B.fl.V,238	
P. leucocoma ; Trichinium, Moquin in De Cand., prodr. XIII, part II, 292 (1849)	—	S.A.	—	—	—	—	—	B.fl.V,238	
P. brachyanthus; Trichin., F. v. M. in Benth. Fl. Austr. V, 639 (1870)	—	—	—	—	—	—	N.A.	B.fl.V,239	
P. arthrolasius, F. v. M., fragm. VI, 232 (1868)	—	—	—	—	—	—	N.A.	B.fl.V,239	M.fr.VI,232.
P. Forrestii, F. v. M., pl. of N. W. Austr. 7 (1881)	—	—	—	—	—	—	N.A.	—	
P. aervoides, F. v. M., fragm. VI, 231 (1868)	—	—	—	—	—	Q.	N.A.	B.fl.V,240	M.fr.III,123; VI,231.
P. Roei; Trichinium, F. v. M. in Benth. Fl. Austr. V, 240 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,240	
P. Hoodii, F. v. M., fragm. VIII, 232 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,232.
P. caespitulosus, F. v. M., fragm. VI, 232 (1868)...	W.A.	—	—	—	—	—	—	B.fl.V,240	M.fr.VI,232.
P. helichrysoides, F. v. M., fragm. VI, 231 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,241	M.fr.VI,231.
P. psilotrichoides, F. v. M., fragm. VI, 231 (1869)	W.A.	—	—	—	—	—	—	B.fl.V,209	M.fr.XI,97.
EUXOLUS, Rafinesque, Fl. Tell. 42 (1836).									
E. Mitchellii, F. v. M. in Giles, geogr. trav. Centr. Austr. 214 (1875)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,214	
E. interruptus, Moquin in De Cand., prodr. XIII, part II, 275 (1849)	—	—	—	—	—	Q.	N.A.	B.fl.V,215	
E. viridis, Moquin in De Cand., prodr. XIII, part II, 273 (1849)	W.A.	—	—	—	N.S.W.	Q.	—	B.fl.V,215	
E. macrocarpus; Amarantus, Benth. Fl. Austr. V, 216 (1870) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,216	
AMARANTUS, Dodoens, stirp. hist. pempt. 185 (1583).									
A. leptostachyus, Bentham, Fl. Austr. V, 214 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.V,213	
A. pallidiflorus, F. v. M., fragm. I, 140 (1859)	W.A.	—	—	—	—	Q.	N.A.	B.fl.V,214	M.fr.I,140.
POLYCNEMON, Linné, gen. pl. ed. sec. 21 (1742). (Hemichroa.)									
P. pentandrum, F. v. M., native pl. of Vict. I, 162 (1879) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,211	M.fr.VI,234.
P. diandrum, F. v. M., native pl. of Vict. I, 163 (1879)... ...	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.V,211	M.fr.VI,234.
P. mesembrianthemum, F. v. M., fragm. VIII, 38 (1873) ...	—	S.A.	—	—	—	—	—	—	M.fr.VIII,38.
DEERINGIA, R. Brown, prodr. 413 (1810). (Lestibudesia partly, Celosia partly, Lagrezia.)									
D. celosioides, R. Brown, prodr. 413 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,210	M.fr.II,92.
D. altissima, F. v. M., fragm. II, 92 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,210	M.fr.VI,251.

SALSOLACEAE.

Linné, class. pl. 507 (1738).

RHAGODIA, R. Brown, prodr. 408 (1810).									
R. Billardieri, R. Brown, prodr. 408 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,152	M.fr.VII,10.
R. parabolica, R. Brown, prodr. 408 (1810)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,153	M.fr.VII,10.
R. dioica, Nees in Lehm. pl. Preiss. I, 636 (1845)...	W.A.	—	—	—	—	—	—	B.fl.V,154	
R. Gaudichaudiana, Moquin, Chenop. monogr. enum. 11 (1840)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,154	M.fr.VII,10.
R. crassifolia, R. Brown, prodr. 408 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.V,154	
R. Preissii, Moquin in De Cand. prodr. XIII, part II, 49 (1849)	W.A.	—	—	—	—	—	—	B.fl.V,155	M.fr.VII,10.
R. obovata, Moquin, Chenop. monogr. enum. 10 (1840)... ...	W.A.	—	—	—	—	—	—	B.fl.V,155	
R. spinescens, R. Brown, prodr. 408 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,155	M.fr.VII,10; XI,20.
R. hastata, R. Brown, prodr. 408 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,156	M.fr.VII,10.
R. nutans, R. Brown, prodr. 408 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,156	M.fr.VII,10.
R. linifolia, R. Brown, prodr. 408 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,157	M.fr.VII,10.
CHENOPODIUM, Tournefort, inst. 506, t. 288 (1700). (Blitum, Ambrina.)									
C. nitrariaceum, F. v. M. in Benth. Fl. Austr. V, 158 (1870) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,158	M.fr.XI,20&27.
C. auricomum, Lindley in Mitch. Trop. Austr. 94 (1848) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,159	M.fr.VII,11.
C. triangulare, R. Brown, prodr. 407 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,160	M.fr.VII,10.
C. microphyllum, F. v. M. in Trans. phil. Inst. Vict. II, 74 (1857)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,161	M.fr.VII,11.
C. rhadinostachyum, F. v. M. in Wing's S. Science Rec. May (1880)	—	S.A.	—	—	—	—	—	—	M.fr.XII.
C. carinatum, R. Brown, prodr. 407 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,162	M.fr.VII,11.
C. cristatum, F. v. M., fragm. VII, 11 (1869)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,163	M.fr.XI,27.
C. atriplicinum, F. v. M., fragm. VII, 11 (1869)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,163	M.fr.I,140; XI,27.
DYSPHANIA, R. Brown, prodr. 411 (1810).									
D. plantaginella, F. v. M., fragm. I, 61 (1858)	—	S.A.	—	—	—	—	N.A.	B.fl.V,164	M.fr.I,61.
D. litoralis, R. Brown, prodr. 411 (1810)	—	S.A.	—	—	—	Q.	N.A.	B.fl.V,164	M.fr.VII,11.
D. myriocephala, Bentham, Fl. Austr. V, 165 (1870)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,165	
ATRIPLEX, Tournefort, inst. 506, t. 288 (1700), from l'Obel (1581). (Obione, Theleophyton.)									
A. Moquinianum, Webb in Cand. prodr. XIII, part. II, 97 (1849)	W.A.	—	—	—	—	—	N.A.	B.fl.V,169	
A. stipitatum, Bentham, Fl. Austr. V, 168 (1870)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,168	M.fr.VII,9.
A. paludosum, R. Brown, prodr. 406 (1810)	W.A.	S.A.	T.	V.	—	—	—	B.fl.V,169	M.fr.VII,9.
A. Drummondii, Moquin in Cand. prodr. XIII, part. II, 102 (1849)	W.A.	—	—	—	—	—	—	B.fl.V,170	
A. isatideum, Moquin, Chenop. monogr. enum. 63 (1840) ...	W.A.	—	—	—	—	—	—	B.fl.V,170	
A. nummularium, Lindley in Mitch. Trop. Austr. 64 (1848) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,170	M.fr.VII,10; XI,27.
A. cinereum, Poiret, Encycl. suppl. I, 471 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,171	M.fr.VII,9.
A. rhagodioides, F. v. M. in Trans. phil. Inst. Vict. II, 74 (1857)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,172	
A. incrassatum, F. v. M., Rep. Babb. Exped. 20 (1858)... ...	—	S.A.	—	—	—	—	—	B.fl.V,172	
A. vesicarium, Heward in Benth. Fl. Austr. V, 172 (1870) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,172	
A. hymenothecum, Moquin in Cand. prodr. XIII, part. II, 101 (1849)	W.A.	—	—	—	—	—	—	B.fl.V,173	M.fr.VII,10.
A. humile, F. v. M., fragm. IV, 48 (1866)...	—	—	—	—	—	—	N.A.	B.fl.V,174	M.fr.IV,48.
A. fissivalve, F. v. M., fragm. IX, 123 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,123.
A. velutinellum, F. v. M., Rep. Babb. Exped. 20 (1858) ...	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,174	M.fr.VII,9.
A. angulatum, Bentham, Fl. Austr. V, 174 (1870)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,174	

A. semibaccatum, R. Brown, prodr. 406 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,175	M.fr.VII,9.
A. exilifolium, F. v. M., fragm. VII, 9 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,175	M.fr.VII,9.
A. Muelleri, Bentham, Fl. Austr. V, 175 (1870)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,175	M.fr.VII,9.
A. elachophyllum, F. v. M., fragm. VII, 8 (1870)	—	—	—	—	—	—	N.A.	B.fl.V,176	M.fr.VII,8.
A. microcarpum, Bentham, Fl. Austr. V, 176 (1870)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,176	
A. prostratum, R. Brown, prodr. 406 (1810)	—	S.A.	—	—	—	—	—	B.fl.V,176	
A. pumilio, R. Brown, prodr. 406 (1810)	—	S.A.	—	—	—	—	—	B.fl.V,176	
A. campanulatum, Bentham, Fl. Austr. V, 177 (1870)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,177	
A. leptocarpum, F. v. M. in Trans. phil. Inst. Vict. II, 74 (1857)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,178	M.fr.VII,9;XI 27
A. limbatum, Bentham, Fl. Aust. V, 178 (1870)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,178	
A. halimoides, Lindley in Mitch. Three Exped. I, 282 (1838) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,179	M.fr.VII,10;XI,2
A. holocarpum, F. v. M., Rep. Babb. Exped. 19 (1858)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,179	
A. spongiosum, F. v. M. in Trans. Vict. phil. Inst. II, 74 (1857)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,179	
A. crystallinum, J. Hooker in Lond. journ. of Bot. VI, 279 (1847)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,180	M.fr.VII.9.
KOCHIA, Roth in Schrader's Journ. Bot. I, 303 (1799). (Maireana, Sclerochlamys.)									
K. fimbriolata, F. v. M., fragm. IX, 75 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,75.
K. lobiflora, F. v. M. in Benth. Fl. Austr. V, 184 (1870) ...	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,184	
K. lanosa, Lindley in Mitch. Trop. Austr. 88 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,184	M.fr.VII,13.
K. triptera, Bentham, Fl. Austr. V, 185 (1870)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,185	
K. oppositifolia, F. v. M. in Transact. Vict. Inst. 134 (1855) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,185	M.fr.VII,12.
K. brevifolia, R. Brown, prodr. 409 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,185	M.fr.VII,13.
K. pyramidata, Bentham, Fl. Austr. V, 186 (1870)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,186	
K. eriantha, F. v. M., Rep. Babb. Exped. 20 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,186	M.fr.VII,13.
K. villosa, Lindley in Mitch. Trop. Austr. 91 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,186	M.fr.VII,12.
K. planifolia, F. v. M., fragm. I, 213 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,187	M.fr.I,213.
K. sedifolia, F. v. M. in Transact. Vict. Inst. I, 134 (1855) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,187	M.fr.VII,12;VIII,37.
K. humillima, F. v. M., fragm. IX, 168 (1875)	—	S.A.	—	V.	N.S.W.	—	—	—	M.fr.IX,168.
K. microphylla, F. v. M., fragm. VIII, 148 (1874)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,181	M.fr.VIII,37.
K. dichoptera, F. v. M., fragm. VIII, 37 (1873)	—	—	—	—	—	Q.	—	—	M.fr.VIII,37,163.
K. decaptera, F. v. M., fragm. IX, 75 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,75.
K. ciliata, F. v. M., Rep. of Babb. Exped. 20 (1858)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,188	M.fr.VII,12.
K. brachyptera, F. v. M., second gen. Rep. 15 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,189	M.fr.VII,13.
K. stelligera, F. v. M., fragm. VII, 13 (1869)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,189	M.fr.VII,13.
DIDYMANTHUS, Endlicher, nov. stirp. decad. 7 (1839).									
D. Roei, Endlicher, nov. stirp. decad. 8 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,193	M.fr.VII,11.
BABBAGIA, F. v. M., Rep. on pl. of Babb. Exped. 21 (1858).									
B. dipterocarpa, F. v. M., Rep. of Babb. Exped. 21 (1858) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,192	M.fr.VII,11.
BASSIA, Allioni in Misc. Taurin. III, 117, t. 4 (1766). (Chenolea, Sclerolaena, Anisacantha, Echinopsilon, Kentropsis, Dissocarpus, Eriochiton, Osteocarpum, Coilocarpus.)									
B. carnosa; Chenolea, Benth. Fl. Austr. V, 190 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,190	M.fr.VII,13.
B. Dallachyana; Chenolea, Benth. Fl. Austr. V, 191 (1870) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,191	
B. tricornis; Chenolea, Benth. Fl. Austr. V, 191 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,191	
B. eurotioides; Chenolea, F. v. M. in Benth. Fl. Austr. V, 191 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,191	M.fr.VII,13.
B. Muelleri; Chenolea, Benth. Fl. Austr. V, 191 (1870)	—	—	—	—	—	—	N.A.	B.fl.V,191	
B. sclerolaenoides; Echinopsilon, F. v. M. in Tr. phil. Inst. Vict. II, 75 (1857)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,192	M.fr.VII,13.
B. uniflora; Sclerolaena, R. Brown, prodr. 410 (1810)	W.A.	S.A.	—	—	—	—	—	B.fl.V,194	
B. diacantha; Anisacantha, Nees in Lehm. pl. Preiss. I, 635 (1844)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,194	M.fr.VII,14;X,91.
B. lanicuspis; Anisacantha, F. v. M., fragm. II, 170 (1861) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,195	M.fr.II,140,170;VII,14.
B. bicornis; Sclerolaena, Lindl. in Mitch. Three Exp. II. 47 (1838)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,195	M.fr.VII,14.
B. biflora; Sclerolaena, R. Brown, prodr. 410 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,196	M.fr.VII,11;X,91.
B. paradoxa; Sclerolaena, R. Brown, prodr. 410 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,196	M.fr.VII,11;X,91.
B. quinquecuspis; Anisac., F. v. M. in Tr. Vict. Inst. I, 134 (1855)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,199	M.fr.VII,14;X,91.
B. Drummondii; Anisacantha, Benth. Fl. Austr. V, 199 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,199	
B. Birchii; Anisacantha, F. v. M., fragm. VIII, 163 (1874) ..	—	—	—	—	—	Q.	—	—	M.fr.VIII,163.
B. divaricata; Anisacantha, R. Brown, prodr. 410 (1810) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,200	M.fr.VII,14;X,92.
B. bicuspis; Anisac., F. v. M. in Trans. Vict. Inst. I, 133 (1855)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,200	M.fr.VII,13.
B. glabra; Kentropsis, F. v. M., fragm. I, 139 (1859)	—	—	—	—	—	—	N.A.	B.fl.V,200	
B. echinopsila; Anisacantha, F. v. M., fragm. VII, 14 (1870) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,201	M.fr.I,139;X,92.
B. enchylaenoides; Chenolea, F. v. M., fragm. X, 92 (1876) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,182	M.fr.X,92.
B. micrantha; Enchylaena, Benth. Fl. Austr. V, 181 (1870) ...	W.A.	—	—	—	—	—	—	B.fl.V,181	
B. brevicuspis; Anisacantha, F. v. M., fragm. IV, 150 (1864) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,198	M.fr.VII,14;VIII,38.
B. salsuginosa; Osteocarpum, F. v. M., second gen. Rep. 15 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,197	M.fr.VII,12;X,92.
ENCHYLAENA, R. Brown, prodr. 408 (1810).									
E. tomentosa, R. Brown, prodr. 408 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,181	M.fr.VII,12;XI,27.
THRELKELDIA, R. Brown, prodr. 409 (1810).									
T. diffusa, R. Brown, prodr. 410 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,197	M.fr.VII,12.
T. proceriflora, F. v. M., fragm. VIII, 38 (1873)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VIII,38.
SALICORNIA, Tournefort, coroll. 51 (1703). (Halocnemum, Arthrocnemum, Tecticornia, Pachycornia).									
S. robusta, F. v. M., fragm. VI, 251 (1868)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,202	M.fr.I,139;VII,14.
S. arbuscula, R. Brown, prodr. 411 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,203	M.fr.VII,15.
S. cinerea, F. v. M., fragm. VI, 251 (1869)	—	—	—	—	—	Q.	N.A.	B.fl.V,203	M.fr.VII,15.
S. leiostachya, Bentham, Fl. Austr. V, 203 (1870)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,203	
S. bidens, Bentham, Fl. Austr. V, 204 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,204	
S. tenuis, Bentham, Fl. Austr. V, 204 (1870)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,204	
S. australis, Solander in G. Forster, prodr. 88 (1786)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,205	M.fr.VII,15;XI,20,27.
SUAEDA, Forskael, Fl. Aegypt. Arab. 69 (1779). (Chenopodium partly, Schoberia, Chenopodina.)									
S. maritima, Dumortier, Fl. Belg. 22 (1827)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,206	M.fr.XI,20.

SALSOLA, Linné, gen. pl. 67 (1737), from Cisalpini.

S. Kali, Linné, spec. plant. 222 (1753) ... . W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,207 M.fr.XI,20,27.

FICOIDEAE.

A. L. de Jussieu, Gen. 315 (1789).

MESEMBRIANTHEMUM, Breyne, prodr. fasc. rar. pl. II, 67 (1689).

M. aequilaterale, Haworth, misc. nat. 77 (1803) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,324 M.fr.XI,20,27.
M. australe, Solander in G. Forster, prodr. 90 (1786) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,324 M.fr.V,158;X,83;XI,20,27

TETRAGONIA, Linné, syst. nat. 9 (1735); Linné, gen. pl. 144 (1737). (Tetragonella.)

T. expansa, Murray in Comm. Goetting. VI, 13, t. 5 (1783) ... — S.A. T. V. N.S.W. Q. — B.fl.III,325 M.fr.VII,129;X,84.
T. diptera, F. v. M., fragm. XI, 8 (1878) ... ... W.A. — — — — — N.A. — M.fr.XI,8.
T. implexicoma, J. Hooker, Fl. Tasman. I, 148 (1859) ... ... W.A. S.A. T. V. N.S.W. — N.A. B.fl.III,326 M.fr.X,84;XI,8.

AIZOON, Linné, gen. pl. 161 (1737). (Aizoum.)

A. quadrifidum, F. v. M., fragm. II, 148 (1861) ... ... W.A. S.A. — — N.S.W. — — B.fl.III,327 M.fr.VII,129;X,83.
A. zygophylloides, F. v. M., fragm. VII, 129 (1871) ... ... W.A. S.A. — — — — — — M.fr.X,83.

GUNNIA, F. v. M., Rep. on pl. of Babb. Exped. 9 (1858).

G. septifraga, F. v. M., Rep. of Babb. Exped. 9 (1858) ... ... — S.A. — — — — — B.fl.III,327
G. Drummondii, Bentham, Fl. Austr. III, 327 (1866) ... ... W.A. — — — — — — B.fl.III,327

SESUVIUM, Linné, syst. ed. decim. 1058 (1759).

S. portulacastrum, Linné, syst. ed. dec. 1058 (1759) ... ... — — — — N.S.W. Q. N.A. B.fl.III,328 M.fr.VII,129;X,83.

ZALEYA, N. L. Burmann, Fl. Indic. 110, t. 31 (1768).

Z. decandra, N. L. Burmann, Fl. Ind. 110, t. 31 (1768) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.III,329 M.fr.I,172;X,83.

TRIANTHEMA, Sauvage, Méth. fol. 127 (1751). (Ancistrostigma.)

T. turgidifolia, F. v. M., fragm. X, 83 (1876) ... ... — — — — — — N.A. — M.fr.X,83.
T. crystallina, Vahl, symbol. I, 32 (1790) ... ... — S.A. — — — Q. N.A. B.fl.III,330 M.fr.I,171-2.
T. glaucifolia, F. v. M., fragm. I, 172 (1859) ... ... — — — — — Q. — — M.fr.I,172.
T. pilosa, F. v. M., fragm. I, 174 (1859) ... ... — S.A. — — — — N.A. B.fl.III,330 M.fr.X,83.
T. oxycalyptra, F. v. M., fragm. I, 173 (1859) ... ... — — — — — — N.A. B.fl.III,330 M.fr.I,173.
T. rhynchocalyptra, F. v. M., fragm. I, 174 (1859) ... ... — — — — — — N.A. B.fl.III,330 M.fr.I,174.
T. cypseloides, Bentham, Fl. Austr. III, 331 (1866) ... ... — — — — N.S.W. Q. N.A. B.fl.III,331 M.fr.X,83.

POMATOTHECA, F. v. M., fragm. X, 72 (1876).

P. humillima, F. v. M., fragm. X, 72 (1876) ... ... — S.A. — — N.S.W. — — — M.fr.X,72.

MACARTHURIA, Huegel. enum. pl. Nov. Holl. Austr. occ. 11 (1837).

M. apetala, Harvey in Hook. Kew. misc. VII, 55 (1855) ... W.A. — — — — — — B.fl.III,332 M.fr.XI,20.
M. australis, Huegel, enum. pl. 11 (1837) ... ... W.A. — — — — — — B.fl.III,332
M. Neo-Cambrica, F. v. M., fragm. V, 28 (1866)... ... ... — — — — N.S.W. Q. — B.fl.III,332 M.fr.V,28.

MOLLUGO, Linné, gen. plant. 336 (1737). (Glinus, Trigastrotheca.)

M. Ginus, A. Richard, tentam. fl. Abyss. I, 48 (1847) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,333
M. orygioides, F. v. M. in Benth. Fl. Austr. III, 333 (1866) ... — S.A. — — N.S.W. Q. — B.fl.III,333
M. Spergula, Linné, spec. pl. ed. alt. 131 (1762) .. .. W.A. S.A. — V. N.S.W. Q. N.A. B.fl.III,334
M. trigastrotheca, F. v. M., pl. of Vict. I, 201 (1860) ... . — — — — — — N.A. B.fl.III,334
M. Cerviana, Seringe in De Cand., prodr. I, 392 (1824) ... .. — S.A. — V. N.S.W. — — B.fl.III,334 M.fr.II,148.

POLYGONACEAE.

A. L. de Jussieu, gen. pl. 82 (1789), from B. de Jussieu (1759).

EMEX, Necker, elem. bot. II, 214 (1790). (Perhaps immigrated.)

E. australis, Steinheil in Ann. des sc. nat. ser. 2, IX, 195, t. 7 (1839) W.A. S.A. — — — — — B.fl.V,262

RUMEX, Linné, gen. pl. 105 (1737), from Plinius.

R. Brownii, Campdera, Monogr. Rum. 81 (1819)... ... ... — S.A. T. V. N.S.W. Q. — B.fl.V,263
R. flexuosus, Solander in G. Forst., fl. ins. Austr. prodr. 90 (1786) — S.A. — — N.S.W. — — B.fl.V,264
R. crystallinus, Lange, ind. sem. hort. Hafn. 28 (1861) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.V,265 M.fr.IV,48.
R. bidens, R. Brown, prodr. 421 (1810) ... ... — S.A. T. V. N.S.W. Q. — B.fl.V,265

POLYGONUM, l'Obel, stirp. hist. 228 (1576), from Dioscorides and Plinius.

P. plebejum, R. Brown, prodr. 420 (1810) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.V,267
P. strigosum, R. Brown, prodr. 420 (1810)... ... ... — — T. V. N.S.W. Q. — B.fl.V,268
P. prostratum, R. Brown, prodr. 419 (1810) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.V,268
P. hydropiper, Linné, spec. plant. 361 (1753) ... ... — S.A. — V. N.S.W. Q. — B.fl.V,269
P. minus, Hudson, Fl. Anglica I, 148 (1762) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.V,269
P. subsessile, R. Brown, prodr. 419 (1810) ... ... — — T. V. N.S.W. Q. — B.fl.V,269
P. barbatum, Linné, spec. plant. 362 (1753) ... ... — — — — N.S.W. Q. — B.fl.V,270
P. articulatum, R. Brown, prodr. 420 (1810) ... ... — — — — — Q. — B.fl.V,270
P. lapathifolium, Linné, spec. plant. 360 (1753) ... ... — S.A. T. V. N.S.W. Q. — B.fl.V,270
P. orientale, Linné, spec. plant. 362 (1753)... ... ... — — — — N.S.W. Q. — B.fl.V,271
P. attenuatum, R. Brown, prodr. 420 (1810) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.V,272

MUEHLENBECKIA, Meissner, gen. pl. vasc. 316 (1840).

M. adpressa, Meissner, gen. pl. comm. 227 (1843) ... ... W.A. S.A. T. V. N.S.W. — — B.fl.V,273 M.fr.XI,20.
M. australis, Meissner, gen. pl. comm. 227 (1843) ... ... — — — — N.S.W. — — — M.fr.IX,169.
M. gracillima, Meissner in De Cand., prodr. XIV, 145 (1856) ... — — — — N.S.W. Q. — B.fl.V,274
M. rhyticarya, F. v. M., fragm. V, 92 (1866) ... ... — — — — N.S.W. Q. — B.fl.V,274
M. axillaris, J. Hooker in Lond. Journ. VI, 278 (1847) ... ... — — T. V. N.S.W. — — B.fl.V,275
M. polybotrya, Meissner in Lehm. pl. Preiss. I, 623 (1845) ... W.A. — — — — — — B.fl.V,475 M.fr.IV,130.
M. polygonoides, F. v. M., fragm. V, 73 (1865) ... ... — S.A. — V. N.S.W. — — B.fl.V,275 M.fr.V,73.
M. stenophylla, F. v. M., fragm. I, 138 (1859) ... ... — — — V. — — — B.fl.V,275 M.fr.I,138.
M. Cunninghamii, F. v. M., fragm. V, 91 (1865) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,275 M.fr.V,95.

PHYTOLACCEAE.

R. Brown in Tuck. Exped. Cong. App. V, 454 (1818).

MONOCOCCUS, F. v. M., fragm. I, 47 (1858).

M. echinophorus, F. v. M., fragm. I, 47 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,144	M.fr.I,47.

DIDYMOTHECA, J. Hooker in Lond. Journ. Bot. VI, 278 (1847). (Cyclotheca.)

D. thesioides, J. Hooker in Lond. Journ. V, 279 (1847)	W.A.	S.A.	T.	—	—	—	—	B.fl.V,145	
D. pleiococca, F. v. M., fragm. I, 202 (1859)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,146	M.fr.I,202.

GYROSTEMON, Desfontaines in Mém. du Mus. VI, 16, t. 6 (1820).

G. brachystigma, F. v. M. in Benth Fl. Austr. V, 146 (1870) ...	W.A.	—	—	—	—	—	—	B.fl.V,146	
G. ramulosus, Desfontaines in Mém. du Mus. V, 17, t. 6 (1820)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.V,147	

CODONOCARPUS, Cunningham in Hook. Bot. Misc. I, 244 (1830). (Hymenotheca.)

C. pyramidalis, F. v. M., plants of Vict. I, 201 (1862)	—	S.A.	—	—	—	—	—	B.fl.V,148	
C. australis, Cunningham in Hook. bot. misc. I, 244 (1830) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,148	
C. cotinifolius, F. v. M., pl. of Vict. I, 200 (1862)	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.V,149	

TERSONIA, Moquin in De Cand., prodr. XIII, part sec. 40 (1849).

T. brevipes, Moquin in De Cand., prod. XIII, part II, 40 (1849)	W.A.	—	—	—	—	—	—	B.fl.V,149	
T. subvolubilis, Bentham, Fl. Austr. V, 150 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,150	

CYPSELOCARPUS, F. v. M., fragm. VIII, 36 (1873).

C. haloragoides, F. v. M., fragm. VIII, 36 (1873)	W.A.	—	—	—	—	—	—	B.fl.V,198	M.fr.VIII,36.

NYCTAGINEAE.

A. L. de Jussieu, Gen. 90 (1789).

BOERHAAVIA, Vaillant, serm. de struct. fl. 50 (1718).

B. diffusa, Linné, spec. fl. Zeylan. 4 (1747)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,277	
B. repanda, Willdenow, sp. pl. I, 22 (1797)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.V,278	

PISONIA, Plumier, nov. pl. Amer. gen. 7, t. 11 (1703).

P. aculeata, Linné, sp. pl. 1026 (1753)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,279	M.fr.VI,197.
P. inermis, G. Forster, prodr. 75 (1786)	—	—	—	—	—	Q.	N.A.	B.fl.V,280	M.fr.VI,197.
P. Brunoniana, Endlicher, prodr. fl. Norf. 43 (1833)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,280	M.fr.I,20;IV,169.

CHORIPETALEAE PERIGYNAE.

F. v. M. in Woolls pl. of the neighb. of Sydney 18 (1880).

CONNARACEAE.

R. Brown, Narr. exped. Cong. App. V, 431 (1878).

ROUREA, Aublet, Hist. des pl. de la Guian. I, 467, t. 187 (1775).

R. brachyandra, F. v. M., fragm. VIII, 6 (1872)	—	—	—	—	—	Q.	—	—	M.fr.VIII,6.

TRICHOLOBUS, Blume, Mus. Bot. Lugd. I, 237 (1850).

T. connaroides, F. v. M., fragm. VIII, 224 (1874)	—	—	—	—	—	Q.	—	—	M.fr.V,224;X,120.

LEGUMINOSAE.

Haller, enum. stirp. Helv. II, 565 (1742).

JANSONIA, Kippist in Gard. Chron. 19 (1847). (Cryptosema.)

J. formosa, Kippist in Transact. Linn. Soc. XX, 384, t. 16 (1851)	W.A.	—	—	—	—	—	—	B.fl.II,8	

BRACHYSEMA, R. Brown in Ait. hort. Kew. sec. edit. III, 10 (1811). (Leptosema, Kaleniczenkia, Burgesia.)

B. praemorsum, Meissner in Lehm. pl. Preiss. I, 25 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,10	
B. lanceolatum, Meissner in Lehm. pl. Preiss. I, 24 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,10	
B. latifolium, R. Brown in Ait. hort. Kew. sec. ed. III, 10 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,10	
B. undulatum, Ker in Bot. Regist. t. 642 (1822)	W.A.	—	—	—	—	—	—	B.fl.II,11	M.fr.IV,11.
B. subcordatum, Bentham, Fl. Austr. II, 11 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,11	
B. bracteolosum, F. v. M., fragm. IV, 10 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,11	M.fr.IV,10.
B. bossiaeoides, Bentham, Fl. Austr. II, 12 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,12	
B. oxylobioides, Bentham, Fl. Austr. II, 12 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,12	
B. uniflorum, R. Brown in Benth., Fl. Austr. II, 12 (1864) ...	—	—	—	—	—	—	N.A.	B.fl.II,12	
B. aphyllum, Hooker, Bot. Mag. t. 4481 (1849)	W.A.	—	—	—	—	—	—	B.fl.II,12	M.fr.I,222.
B. macrocarpum, Bentham, Fl. Austr. II, 13 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,13	
B. tomentosum, Bentham, Fl. Austr. II, 13 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,13	
B. Chambersii, F. v. M. in Benth., Fl. Austr. II, 13 (1864) ...	—	S.A.	—	—	—	—	N.A.	B.fl.II,13	
B. daviesioides, Bentham, Fl. Austr. II, 13 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,13	

OXYLOBIUM, Andrews, Bot. Reposit. t. 492 (1809). (Chorizematis subgenus, Callistachys, Podolobium.)

O. Callistachys, Bentham, Fl. Austr. II, 16 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,16	M.fr.IV,18.
O. ellipticum, R. Brown in Ait. hort. Kew. sec. ed. III, 10 (1811)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,16	
O. alpestre, F. v. M. in Transact. phil. Soc. Vict. I, 38 (1854) ...	—	—	—	V.	N.S.W.	—	—	B.fl.II,17	
O. lineare, Bentham, Fl. Austr. II, 17 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,17	M.fr.IV,17.
O. carinatum, Bentham, Fl. Austr. II, 18 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,18	
O. spathulatum, Bentham, Fl. Austr. II, 18 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,18	
O. obtusifolium, Sweet, Fl. Austr. t. 5 (1827)	W.A.	—	—	—	—	—	—	B.fl.II,18	
O. cordifolium, Andrews, Bot. Regist. t. 492 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.II,19	M.fr.IV,17.
O. microphyllum, Bentham, Fl. Austr. II, 19 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,19	
O. Pulteneae, De Candolle, prodr. II, 104 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,19	M.fr.IV,19.
O. hamulosum, Bentham in A. Gray, Bot. Wilk. Exped. t. 379	—	—	—	—	N.S.W.	—	—	B.fl.II,20	
O. scandens, Bentham in Ann. Wien. Mus. II, 70 (1839)... ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,20	
O. procumbens, F. v. M. in Transact. phil. Soc. Vict. I, 37 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.II,20	
O. tricuspidatum, Meissner in Lehm. pl. Preiss. I, 30 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,21	

O. spectabile, Endlicher, nov. stirp. dec. 2 (1839)	W.A.	—	—	—	—	—	—	B.fl.II,21	
O. atropurpureum, Turczanin. in Bull. Soc. Mosc. XXVI, 250 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,22	
O. retusum, R. Brown in Bot. Regist. t. 913 (1825)	W.A.	—	—	—	—	—	—	B.fl.II,22	
O. virgatum, Bentham, Fl. Austr. II, 22 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,22	
O. reticulatum, Meissner in Lehm. pl. Preiss. I, 29 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,23	
O. capitatum, Bentham in Hueg. enum. 28 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,23	
O. cuneatum, Bentham in Bot. Regist. XXV, App. XII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.II,23	
O. acutum, Bentham, Fl. Austr. II, 24 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,24	
O. parviflorum, Bentham in Bot. Regist. XXV, App. XII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,24	
O. heterophyllum, Bentham, Fl. Austr. II, 25 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,25	
O. aciculiferum, F. v. M., fragm. I, 75, implied (1859)	—	—	—	—	—	Q.	—	B.fl.II,25	M.fr.I,75.
O. trilobatum, F. v. M., annual report 23 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,25	M.fr.IV,19.
CHORIZEMA, Labillardière, Voy. I, 405 (1799). (Orthotropis.)									
C. Dicksonii, Graham in Maund's Botanist, t. 106 (1841) ...	W.A.	—	—	—	—	—	—	B.fl.II,27	
C. nervosum, Th. Moore in Gard. Companion (1852)	W.A.	—	—	—	—	—	—	B.fl.II,27	
C. varium, Bentham in Bot. Regist. XXV, t. 49 (1839)	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. cordatum, Lindley, Bot. Regist. t. 10 (1838)	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. ilicifolium, Labillardière, Voy. I, 405, t. 21 (1799)	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. rhombeum, R. Brown in Ait. hort. Kew. sec. ed. III, 9 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. diversifolium, A. de Candolle, Pl. rar. jard. Genèv. 44, t. 8 (1836)	W.A.	—	—	—	—	—	—	B.fl.II,29	
C. angustifolium, Bentham in Hueg. enum. 28 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,29	
C. reticulatum, Meissner in Lehm. pl. Preiss. I, 34 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,30	
C. trigonum, Turcz. in Bull. de la Soc. Mosc. XXVI, 254 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,30	
C. humile, Turcz. in Bull. de la Soc. Mosc. XXVI, 254 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,30	
C. parviflorum, Bentham in Ann. des Wien. Mus. II, 71 (1838)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,30	
C. cytisoides, Turczaninow in Bull. Soc. Mosc. XXVI, 256 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,31	
C. Henchmanni, R. Brown in Bot. Regist. t. 986 (1826)	W.A.	—	—	—	—	—	—	B.fl.II,31	
C. ericifolium, Meissner in Lehm. pl. Preiss. II, 209 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.II,32	
GASTROLOBIUM, R. Brown in Ait. hort. Kew. sec. edit. III, 16 (1811).									
G. pyramidale, Th. Moore in Gard. Comp. I, 81 (1852)	W.A.	—	—	—	—	—	—	B.fl.II,98	
G. Lehmanni, Meissner in Lehm. pl. Preiss. I, 70 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,98	M.fr.X,35.
G. pulchellum, Turcz. in Bull. de la Soc. Mosc. XXVI, 274 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,98	
G. stipulare, Meissner in Lehm. pl. Preiss. II, 218 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.II,99	
G. Brownii, Meissner in Lehm. pl. Preiss. I, 71 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,99	
G. reticulatum, Bentham, Fl. Austr. II, 99 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,99	M.fr.X,35.
G. elachistum, F. v. M., fragm. IX, 67 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,67.
G. truncatum, Bentham, Fl. Austr. II, 99 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,99	
G. spathulatum, Bentham in Bot. Regist. XX, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,100	
G. plicatum, Turczanin. in Bull. Soc. Mosc. XXVI, 274 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,100	
G. tricuspidatum, Meissner in Lehm. pl. Preiss. I, 66 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,100	
G. obovatum, Bentham in Bot. Regist. XXV, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,101	M.fr.X,35.
G. epacridioides, Meissner in Lehm. pl. Preiss. I, 72 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,102	
G. trilobum, Bentham in Bot. Regist. XXV, App. XIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.II,101	
G. ilicifolium, Meissner in Lehm. pl. Preiss. I, 67 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,101	
G. villosum, Bentham in Bot. Regist. XXV, App. XIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.II,102	
G. polystachyum, Meissner in Lehm. pl. Preiss. II, 217 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.II,102	
G. ovalifolium, Henfrey in Gard. Comp. I, 41 (1852)	W.A.	—	—	—	—	—	—	B.fl.II,102	
G. grandiflorum, F. v. M., fragm. III, 17 (1862)	—	—	—	—	—	Q.	N.A.	B.fl.II,103	M.fr.III,17;IV,174;X,35.
G. pycnostachyum, Bentham, Fl. Austr. II, 103 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,103	
G. spinosum, Bentham in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,103	M.fr.X,35.
G. rotundifolium, Meissner in Lehm. pl. Preiss. II, 216 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.II,104	
G. microcarpum, Meissner in Lehm. pl. Preiss. I, 70 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,104	
G. oxylobioides, Bentham in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,104	
G. calycinum, Bentham in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,104	M.fr.X,35.
G. Callistachys, Meissner in Lehm. pl. Preiss. II, 216 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.II,105	
G. stenophyllum, Turczanin. in Bull. Soc. Mosc. XXVI, 275 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,105	
G. crassifolium, Bentham, Fl. Austr. II, 105 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,105	
G. parvifolium, Bentham, Fl. Austr. II, 106 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,106	
G. hamulosum, Meissner in Lehm. pl. Preiss. II, 218 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.II,106	
G. velutinum, Lindley & Paxton, Flower Gard. III, 76 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,106	M.fr.X,35.
G. bidens, Meissner in Regensb. Bot. Zeit. 29 (1855)	W.A.	—	—	—	—	—	—	B.fl.II,106	
G. bilobum, R. Brown in Ait. hort. Kew. sec. edit. III, 16 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,107	M.fr.X,35.
G. seorsifolium, F. v. M., fragm. X, 35 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,35.
ISOTROPIS, Bentham in Hueg. enum. 28 (1837).									
I. striata, Bentham in Hueg. enum. 28 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,39	M.fr.III,16.
I. Drummondii, Meissner in Lehm. pl. Preiss. I, 31 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,39	
I. juncea, Turczanin. in Bull. Soc. Mosc. XXVI, 251 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,39	M.fr.X,51.
I. canescens, F. v. M., fragm. X, 51 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,51.
I. atropurpurea, F. v. M., fragm. III, 16 (1862)	—	S.A.	—	—	—	—	N.A.	B.fl.II,40	M.fr.III,16;X,51.
I. filicaulis, Bentham in Ann. des Wien. Mus. II, 71 (1838) ...	—	—	—	—	—	Q.	—	B.fl.II,40	M.fr.IV,20.
I. parviflora, Bentham in Ann. des Wien. Mus. II, 71 (1838) ...	—	—	—	—	—	—	N.A.	B.fl.II,40	
I. Wheeleri, F. v. M. in Benth. Fl. Austr. II, 40 (1864) ...	—	S.A.	—	—	N.S.W.	—	—	B.fl.II,40	
MIRBELIA, Smith in Koenig & Sims, Ann. of Bot. I, 511 (1805). (Dichosema, Oxycladium.)									
M. dilatata, R. Brown in Ait. hort. Kew. sec. ed. III, 21 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,33	
M. racemosa, Turczanin. in Bull. Soc. Mosc. XXVI, 282 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,33	M.fr.IV,18.
M. grandiflora, Aiton in Bot. Mag. t. 2771 (1827)	—	—	—	—	N.S.W.	—	—	B.fl.II,34	
M. subcordata, Turczanin. in Bull. Soc. Mosc. XXVI, 282 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,34	
M. ovata, Meissner in Lehm. pl. Preiss. I, 77 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,34	

M. oxylobioides, F. v. M., fragm. II, 154 (1861)	—	—	—	V.	N.S.W.	—	—	B.fl.II,34	M.fr.IV 12.
M. reticulata, Smith in Koen. & Sims, Ann. of Bot. I, 511 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,35	
M. aotoides, F. v. M. in Transact. phil. Inst. Vict. III, 53 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,35	
M. pungens, Cunningham in G. Don, gen. syst. II, 126 (1832) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,35	
M. speciosa, Sieber in De Cand. prodr. II, 115 (1824)	—	—	—	—	N.S.W.	—	—	B.fl.II,36	
M. floribunda, Bentham in Bot. Regist. XXV, App. XII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,36	
M. spinosa, Bentham, Fl. Austr. II, 36 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,36	
M. multicaulis, Bentham, Fl. Austr. II, 37 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,37	
M. microphylla, Bentham, Fl. Austr. II, 37 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,37	
M. daviesioides, Bentham, Fl. Austr. II, 37 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,37	M.fr.IV,11.
M. oxyclada, F. v. M., fragm. IV, 12 (1864)	—	S.A.	—	—	—	—	N.A.	B.fl.II,38	M.fr.IV,12.
GOMPHOLOBIUM, Smith in Transact. Linn. Soc. IV, 220 (1798).									
G. ovatum, Meissner in Lehm. pl. Preiss. I, 35 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,42	
G. amplexicaule, Meissner in Lehm. pl. Preiss. I, 36 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,42	
G. latifolium, Smith in Koenig and Sims, Ann. of Bot. I, 505 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,42	
G. Huegelii, Bentham in Hueg. enum. 29 (1837)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,43	
G. polymorphum, R. Brown in Ait. hort. Kew, sec. ed. III, 11 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,43	
G. obcordatum, Turczanin. in Bull. Soc. Mosc. XXVI, 258 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,44	
G. marginatum, R. Brown in Ait. hort. Kew, sec. ed. III, 11 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,44	
G. grandiflorum, Smith, Exot. Bot. t. 5 (1804)	—	—	—	—	N.S.W.	—	—	B.fl.II,44	
G. virgatum, Sieber in De Cand. prodr. II, 105 (1824)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,45	
G. minus, Smith in Transact. Linn. Soc. IX, 251 (1808)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,45	
G. uncinatum, Cunningham in Annal. des Wien. Mus. II, 72 (1838)	—	—	—	—	N.S.W.	—	—	B.fl.II,46	
G. Baxteri, Bentham, Fl. Austr. II, 46 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,46	
G. aristatum, Bentham in Ann. des Wien. Mus. II, 72 (1838) ...	W.A.	—	—	—	—	—	—	B.fl.II,46	
G. burtonioides, Meissner in Lehm. pl. Preiss. I, 37 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,46	
G. capitatum, Cunningham in Bot. Regist. t. 1563 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.II,47	
G. tomentosum, Labillard., Nov. Holl. pl. sp. I. 106, t. 134 (1804)	W.A.	—	—	—	—	—	—	B.fl.II,47	
G. Preissii, Meissner in Lehm. pl. Preiss. I, 40 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,17	
G. viscidulum, Meissner in Lehm. pl. Preiss. I, 39 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,48	
G. glabratum, De Candolle, prodr. II, 106 (1824)	—	—	—	—	N.S.W.	—	—	B.fl.II,48	
G. nitidum, Solander in Benth. Fl. Austr. II, 48 (1864) ...	—	—	—	—	—	Q.	—	B.fl.II,48	
G. pinnatum, Smith in Transact. Linn. Soc. IX, 251 (1808) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,48	
G. Shuttleworthii, Meissner in Lehm. pl. Preiss. I, 39 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,49	
G. venustum, R. Brown in Ait. hort. Kew. sec. ed. III, 12 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,49	
G. Knightianum, Lindley, Bot. Regist. t. 1468 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,49	
BURTONIA, R. Brown in Ait. hort. Kew, ed. sec. III, 13 (1811).									
B. subulata, Bentham, Fl. Austr. II, 50 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,50	M.fr.III,30;X,35.
B. foliolosa, Bentham, Fl. Austr. II, 50 (1864)	—	—	—	—	—	Q.	—	B.fl.II,50	
B. polyzyga, Bentham, Fl. Austr. II, 51 (1864)	—	S.A.	—	—	—	—	N.A.	B.fl.II,51	M.fr.III,29;X,34.
B. villosa, Meissner in Lehm. pl. Preiss. I, 41 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,51	M.fr.X,35.
B. Hendersonii, Bentham, Fl. Austr. II, 51 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,51	
B. gompholobioides, F. v. M., fragm. X, 34 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,34.
B. scabra, R. Brown in Ait. hort. Kew. sec. ed. III, 12 (1811) ...	W.A.	—	—	—	—	—	—	B.fl.II,51	M.fr.X,35.
B. conferta, De Candolle, prodr. II, 106 (1825)	W.A.	—	—	—	—	—	—	B.fl.II,52	M.fr.X.35.
JACKSONIA, R. Brown in Ait. hort. Kew. sec. ed. III, 12 (1811). (Piptomeris.)									
J. dilatata, Bentham in Ann. des Wien. Mus. II, 74 (1838) ...	—	—	—	—	—	—	N.A.	B.fl.II,54	M.fr.X,37.
J. densiflora, Bentham in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,54	M.fr.X,38.
J. carduacea, Meissner in Bot. Zeit. 25 (1855)	W.A.	—	—	—	—	—	—	B.fl.II,55	
J. floribunda, Endlicher in Ann. des Wien. Mus. II. 197 (1838)	W.A.	—	—	—	—	—	—	B.fl.II,55	
J. odontoclada, F. v. M. in Benth. Fl. Austr. II, 55 (1864) ...	—	—	—	—	—	—	N.A.	B.fl.II,55	M.fr.X,37.
J. ramosissima, Bentham in Mitch. Trop. Austr. 258 (1848) ...	—	—	—	—	—	Q.	—	B.fl.II,56	M.fr.X,37.
J. foliosa, Turcz. in Bull. de la Soc. Imp. de Mosc. XXVI, 260 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,56	
J. spinosa, R. Brown in Ait. hort. Kew. sec. ed. III, 13 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,56	M.fr.X,38&39.
J. stricta, Meissner in Regensb. Bot. Zeit. 27 (1855)	W.A.	—	—	—	—	—	—	B.fl.II,56	
J. hakeoides, Meissner in Lehm. pl. Preiss. I, 45 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,57	M.fr.X,39.
J. furcellata, De Candolle, prodr. II, 107 (1824)	W.A.	—	—	—	—	—	—	B.fl.II,57	M.fr.X,39.
J. horrida, De Candolle, prodr. II, 107 (1824)	W.A.	—	—	—	—	—	—	B.fl.II,57	
J. sericea, Bentham in Hueg. enum. 31 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,58	
J. Sternbergiana, Huegel, Bot. Archiv. t. 3 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,58	M.fr.X,39.
J. vernicosa, F. v. M. in Benth. Fl. Austr. II, 58 (1864) ...	—	—	—	—	—	—	N.A.	B.fl.II,58	M.fr.X,37.
J. rhadinoclona, F. v. M., fragm. X, 37 (1876)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.X,37.
J. thesioides, Cunningh. in den Ann. des Wien. Mus. II, 74 (1838)	—	—	—	—	—	Q.	N.A.	B.fl.II,59	M.fr.IV,161;X,37.
J. compressa, Turcz. in Bull. de la Soc. de Mosc. XXVI, 260 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,59	M.fr.X,38.
J. scoparia, R. Brown in Ait. hort. Kew. sec. ed. III, 13 (1811)	W.A.	—	—	—	N.S.W.	Q.	—	B.fl.II,59	M.fr.X,37.
J. cupulifera, Meissner in der Bot. Zeitung 27 (1855)	W.A.	—	—	—	—	—	—	—	M.fr.X,36.
J. restioides, Meissner in Lehm. pl. Preiss. I, 46 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,60	
J. velutina, Bentham, Fl. Austr. II, 60 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,60	
J. Lehmanni, Meissner in Lehm. pl. Preiss. I, 46 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,60	
J. racemosa, Meissner in Lehm. pl. Preiss. II, 212 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.II,60	
J. umbellata, Turczaninow in Bull. Soc. Mosc. XXVI, 261 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,61	M.fr.X,38.
J. nematoclada, F. v. M., fragm. X, 50 (1876)	W.A.	—	—	—		—	—	—	M.fr.X,50.
J. rhadinoclada, F. v. M., fragm. X, 38 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,38.
J. capitata, Meissner in Lehm. pl. Preiss. I, 45 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,61	M.fr.X,38.
J. alata, Bentham in Hueg. enum. 30 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,61	M.fr.X,38.
J. angulata, Bentham, Fl. Austr. II, 62 (1864)	W.A.	—	—	—	—	—	-	B.fl.II,62	
J. Stackhousii, F. v. M. in proc. Linn. Soc. N.S.W., VI, 791 (1881)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.XI,138
J. pteroclada, F. v. M., fragm. X, 37 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,37.
J. macrocalyx, Meissner in der Bot. Zeit. 26 (1855)	W.A.	—	—	—	—	—	—	B.fl.II,62	

J. purpurascens, F. v. M., fragm. IV, 161 (1864)...	—	—	—	—	—	Q.	—	B.fl.II,62	M.fr.IV,161.
J. piptomeris, Bentham, Fl. Austr. II, 62 (1864)...	W.A.	—	—	—	—	—	—	—	
SPHAEROLOBIUM, Smith in Koenig & Sims, Ann. of Bot. I, 509 (1805). (Roea.)									
S. linophyllum, F. v. M., fragm. I, 167 (1859)	W.A.	—	—	—	—	—	—	B.fl.II,64	M.fr.I,167.
S. foliosum, F. v. M., fragm. I, 166 (1859)...	W.A.	—	—	—	—	—	—	B.fl.II,64	M.fr.I,166.
S. gracile, Bentham, Fl. Austr. II, 64 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,64	
S. racemulosum, Bentham, Fl. Austr. II, 64 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,65	
S. alatum, Bentham in Hueg. enum. 32 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,65	
S. vimineum, Smith in Koen. & Sims, Ann. of Bot. I, 509 (1805)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,65	
S. grandiflorum, R. Brown in Benth. Fl. Austr. II, 66 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.II,66	
S. fornicatum, Bentham in Hueg. enum. 32 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,66	
S. medium, R. Brown in Ait. hort. Kew. sec. ed. III, 14 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,66	
S. scabriusculum, Meissner in Lehm. pl. Preiss. II, 214 (1849)...	W.A.	—	—	—	—	—	—	B.fl.II,66	
S. macranthum, Meissner in Lehm. pl. Preiss. II, 213 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.II,67	
S. daviesioides, Turczaninow in Bull. Soc. Mosc. XXVI, 266 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,67	
VIMINARIA, Smith, Exotic Botany 51, t. 27 (1804).									
V. denudata, Smith, Exot. Bot. 51, t. 27 (1804)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,68	
DAVIESIA, Smith in Transact. Linn. Soc. IV, 222 (1798).									
D. cordata, Smith in Transact. Linn. Soc. IX, 259 (1808) ...	W.A.	—	—	—	—	—	—	B.fl.II,72	
D. ovata, Bentham, Fl. Austr. II, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,72	
D. crenulata, Turczanin. in Bull. Soc. de Mosc. XXVI, 265 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,72	M.fr.IV,16.
D. oppositifolia, Endl. in den. Ann. des Wien. Mus. II, 199 (1838)	W.A.	—	—	—	—	—	—	B.fl.II,73	
D. alternifolia, Endlicher in Ann. Wien. Mus. II, 199 (1838) ...	W.A.	—	—	—	—	—	—	B.fl.II,73	
D. elongata, Bentham, Fl. Austr. II, 74 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,74	
D. pedunculata, Bentham in Bot. Regist. XXV, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,74	
D. arthropoda, F. v. M., fragm. VIII, 225 (1874)...	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,225.
D. mollis, Turczaninow in Bull. Soc. Mosc. XXVI, 263 (1853)...	W.A.	—	—	—	—	—	—	B.fl.II,74	
D. concinna, R. Brown in Benth. Fl. Austr. II, 75 (1864) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,75	
D. umbellulata, Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,75	
D. Wyattii, Bailey in pap. Gardn. Assoc. of S. Austr. (1880) ...	—	—	—	V.	N.S.W.	Q.	—	—	M.fr.X,138.
D. latifolia, R. Brown in Ait. hort. Kew. sec. ed. III, 20 (1811)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,76	
D. corymbosa, Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,76	
D. horrida, Meissner in Lehm. pl. Preiss. I, 54 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,77	
D. reclinata, Cunningham in Benth. Fl. Austr. II, 77 (1864) ...	—	—	—	—	—	—	N.A.	B.fl.II,77	
D. obtusifolia, F. v. M., fragm. II, 104 (1861)	W.A.	—	—	—	—	—	—	B.fl.II,77	M.fr.II,104.
D. obovata, Turczaninow in Bull. Soc. Mosc. XXVI, 261 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,78	
D. longifolia, Bentham in Bot. Regist. XXV, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,78	
D. chordophylla, Meissner in Lehm. plant. Preiss. I, 48 (1844)...	W.A.	—	—	—	—	—	—	B.fl.II,78	
D. nematophylla, F. v. M. in Benth. Fl. Austr. II, 78 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.II,78	
D. daphnoides, Meissner in Lehm. plant. Preiss. I, 54 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,79	
D. nudiflora, Meissner in Lehm. plant. Preiss. I, 53 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,79	
D. rhombifolia, Meissner in Lehm. plant. Preiss. I, 56 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,79	
D. Drummondii, Meissner in Lehm. plant. Preiss. I, 53 (1844)...	W.A.	—	—	—	—	—	—	B.fl.II,80	
D. filipes, Bentham in Mitch. Trop. Austr. 363 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,80	
D. cardiophylla, F. v. M., fragm. II, 105 (1861)	W.A.	—	—	—	—	—	—	B.fl.II,79	M.fr.II,105.
D. squarrosa, Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,80	
D. ulicina, Smith in Koen. & Sims, Ann. of Bot. I, 506 (1805) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,81	
D. acicularis, Smith in Koen. & Sims, Ann. of Bot. I, 506 (1805)	—	—	—	—	N.S.W.	—	—	B.fl.II,81	
D. pachyphylla, F. v. M., fragm. IV, 15 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,82	M.fr.IV,15.
D. teretifolia, R. Brown in Benth. Fl. Austr. II, 82 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.II,82	
D. genistifolia, Cunningham in Annal. Wien. Mus. II, 75 (1838)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,82	
D. hakeoides, Meissner in Lehm. plant. Preiss. I, 47 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,82	
D. colletioides, Meissner in Lehm. plant. Preiss. I, 48 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,83	
D. reversifolia, F. v. M., fragm. I, 145 (1859)	W.A.	—	—	—	—	—	—	B.fl.II,83	M.fr.I,145.
D. incrassata, Smith in Transact. Linn. Soc. IX, 255 (1808) ...	W.A.	S.A.	—	—	—	—	—	B.fl.II,83	
D. brevifolia, Lindley in Mitch. Three Exped. II, 201 (1838) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,84	
D. acanthoclona, F. v. M., fragm. X, 32 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,32.
D. Preissii, Meissner in Lehm. plant. Preiss. I, 50 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,84	
D. spinosissima, Meissner in Lehm. plant. Preiss. I, 51 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,85	
D. pachylina, Turczaninow in Bull. Soc. Mosc. XXVI, 263 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,85	
D. quadrilatera, Bentham in Bot. Regist. XXV, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,85	
D. striata, Turczaninow in Bull. Soc. Mosc. XXVI, 264 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,85	M.fr.II,105.
D. polyphylla, Bentham in Bot. Regist. XXV, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,86	
D. microphylla, Bentham, Fl. Austr. II, 86 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,86	
D. flexuosa, Bentham in Hueg. enum. 32 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,86	
D. pectinata, Lindley in Mitch. Three Exped. II, 151 (1838) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,87	
D. trigonophylla, Meissner in Lehm. plant. Preiss. II, 213 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,87	
D. epiphylla, Meissner in Regensb. Bot. Zeit. 27 (1855)... ...	W.A.	—	—	—	—	—	—	B.fl.II,87	
D. euphorbioides, Bentham, Fl. Austr. II, 88 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,88	
D. divaricata, Bentham in Hueg. enum. 31 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,88	
D. paniculata, Bentham in Hueg. enum. 31 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,88	
D. aphylla, F. v. M. in Benth. Fl. Austr. 88 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,88	
D. juncea, Smith in Transact. Linn. Soc. IX, 260 (1808) ...	W.A.	—	—	—	—	—	—	B.fl.II,88	
D. alata, Smith in Transact. Linn. Soc. IX, 259 (1808)	—	—	—	—	N.S.W.	—	—	B.fl.II,89	
D. anceps, Turczaninow in Bull. Soc. Mosc. XXVI, 266 (1853)...	W.A.	—	—	—	—	—	—	B.fl.II,89	
AOTUS, Smith in Koenig & Sims, Ann. of Bot. I, 504 (1805).									
A. gracillima, Meissner in Lehm. pl. Preiss. I, 59 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,90	
A. villosa, Smith in Koen. & Sims, Ann. of Bot. I, 504 (1805) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,90	
A. mollis, Bentham in Mitch. Trop. Austr. 236 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,91	

A. Preissii, Meissner in Lehm. plant. Preiss. II, 214 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.II,91	
A. phylicoides, F. v. M. in Benth. Fl. Austr. II, 72 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.II,92	
A. lanigera, Cunningham in Annal. des Wien. Mus. II, 78 (1838)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,92	
A. genistoides, Turczaninow in Bull. Soc. Mosc. XXVI, 268 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,92	
A. carinata, Meissner in Lehm. plant. Preiss. II, 215 (1849) ...	W.A.	—	—	—	—	—	—	B.fl.II,93	
A. passerinoides, Meissner in Lehm. plant. Preiss. I, 61 (1844)...	W.A.	—	—	—	—	—	—	B.fl.II,93	
A. Tietkensii, F. v. M., fragm. X, 33 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,33.
A. cordifolia, Bentham in Hueg. enum. 33 (1837)...	W.A.	—	—	—	—	—	—	B.fl.II,93	
PHYLLOTA, De Candolle, prodr. II, 113 (1825). (Urodon.)									
P. barbata, Bentham in Hueg. enum. 33 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,94	
P. gracilis, Turczaninow in Bull. Soc. Mosc. XXVI, 267 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,94	
P. Sturtii, Bentham, Fl. Austr. II, 95 (1864)	—	S.A.	—	—	—	—	—	B.fl.II,95	
P. phylicoides, Bentham in Ann. des Wien. Mus. II, 77 (1838)...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,95	M.fr.X,33.
P. humifusa, Bentham, Fl. Austr. II, 95 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,95	
P. diffusa, F. v. M., fragm. I, 8 (1858)	—	—	T.	—	—	—	—	B.fl.II,119	M.fr.I,8.
P. pleurandroides, F. v. M. in Transact. phil. Inst. Vict. I, 38 (1854)	—	S.A.	—	V.	—	—	—	B.fl.II,96	M.fr.X,33.
P. Luehmanni, F. v. M., fragm. X, 33 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,33.
P. Urodon, F. v. M. in papers of the Roy. Soc. of Tasm. 120 (1877)	W.A.	—	—	—	—	—	—	B.fl.II,124	
PULTENAEA, Smith, Specim. of the Bot. of New Holl. 35, t. 12 (1793). (Euchilus, Spadostyles, Bartlingia.)									
P. daphnoides, Wendland, Bot. Beob. 49 (1798)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,112	
P. stricta, Sims, Bot. Mag. 1588 (1813)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,112	
P. retusa, Smith in Koen. & Sims, Ann. of Bot. I, 502 (1805) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,113	
P. Benthamii, F. v. M. in Transact. phil. Inst. Vict. I, 38 (1854)	—	—	—	V.	—	—	—	B.fl.II,113	
P. pycnocephala, F. v. M. in Benth. Fl. Austr. II, 114 (1864) ...	—	—	—	—	N.S.W.	—	—	B.fl.II,114	
P. myrtoides, Cunningham in Annal. des Wien. Mus. II, 81 (1838)	—	—	—	—	—	Q.	—	B.fl.II,114	
P. mucronata, F. v. M., fragm. I, 8 (1858)...	—	S.A.	—	V.	—	—	—	B.fl.II,114	M.fr.I,8.
P. polifolia, Cunningham in Field's N. S. Wales 346 (1825) ...	—	—	—	—	N.S.W.	—	—	B.fl.II,115	
P. petiolaris, Cunningham in Ann. des Wien. Mus. II, 82 (1838)	—	—	—	—	—	Q.	—	B.fl.II,115	
P. paleacea, Willdenow, spec. plant. II, 506 (1799)	—	—	—	V.	N.S.W.	—	—	B.fl.II,115	
P. Gunnii, Bentham in Ann. des Wien. Mus. II, 82 (1839) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.II,116	
P. scabra, R. Brown in Ait. hort. Kew. sec. ed. III, 18 (1811)...	—	—	—	V.	N.S.W.	—	—	B.fl.II,116	
P. Hartmanni, F. v. M., fragm. VIII, 166 (1874)...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VIII,166.
P. microphylla, Sieber in De Cand. prodr. II, 11 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,117	
P. Skinneri, F. v. M., fragm. VIII, 166 (1874)	W.A.	—	—	—	—	—	—	—	M.fr.VIII,166.
P. Drummondii, Meissner in Lehm. plant. Preiss. II, 219 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,117	
P. pinifolia, Meissner in Lehm. plant. Preiss. II, 220 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.II,118	
P. pedunculata, Hooker, Bot. Magaz. t. 2859 (1828)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,118	
P. conferta, Bentham, Fl. Austr. II, 118 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,118	
P. aciphylla, Bentham in Hueg. enum. 35 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,119	
P. ochreata, Meissner in Lehm. plant. Preiss. I, 73 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,120	
P. aspalathoides, Meissner in Lehm. plant. Preiss. I. 73 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,120	
P. obcordata, Bentham, Fl. Austr. II, 120 (1864)...	W.A.	—	—	—	—	—	—	B.fl.II,120	
P. rotundifolia, Bentham, Fl. Austr. II, 121 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,121	M.fr.I,145.
P. calycina, Bentham, Fl. Austr. II, 121 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,121	
P. spinulosa, Bentham, Fl. Austr. II, 121 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,121	
P. tenella, Bentham, Fl. Austr. II, 122 (1864)	—	—	—	V.	—	—	—	B.fl.II,122	
P. ternata, F. v. M., fragm. I, 8 (1858)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,122	M.fr.I,8;IV,10.
P. styphelioides, Cunningham in G. Don, gen. syst. II, 124 (1832)	—	—	—	V.	N.S.W.	—	—	B.fl.II,122	
P. altissima, F. v. M. in Bentham, Fl. Austr. II, 123 (1864) ...	—	—	—	V.	N.S.W.	—	—	B.fl.II,123	
P. obovata, Bentham, Fl. Austr. II, 123 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,123	
P. incurvata, Cunningham in Field's N.S. Wales 346 (1825) ...	—	—	—	—	N.S.W.	—	—	B.fl.II,124	
P. subumbellata, Hooker, Bot. Mag. t. 3254 (1833)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,124	
P. stipularis, Smith, Specim. of the Bot. of New Holl. 35, t. 12 (1793)	—	—	—	—	N.S.W.	—	—	B.fl.II,125	
P. glabra, Bentham, Fl. Austr. II, 124 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,125	
P. dentata, Labillardière, Nov. Holl. pl. spec. I, 103, t. 131 (1804)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,125	
P. aristata, Sieber in De Cand. prodr. II, 112 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,126	
P. plumosa, Sieber in De Cand. prodr. II, 111 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,126	
P. viscosa, R. Brown in Benth. Fl. Austr. II, 127 (1864) ...	—	—	—	V.	N.S.W.	—	—	B.fl.II,127	
P. echinula, Sieber in De Cand. prodr. II, 112 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,127	
P. hibbertioides, J. Hooker, Fl. Tasm. I, 89 (1860)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,127	
P. rosea, F. v. M., fragm. II, 15 (1860)	—	—	—	V.	—	—	—	B.fl.II,128	M.fr.II,15.
P. mollis, Lindley in Mitch. Three Exped. II, 260 (1838) ..	—	S.A.	—	V.	—	—	—	B.fl.II,128	
P. strobilifera, Meissner in Lehm. pl. Preiss. I, 75 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,128	
P. ericifolia, Bentham in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,129	
P. verruculosa, Turcz. in Bull. Soc. Mosc. XXVI, 278 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,129	
P. empetrifolia, Meissner in Lehm. plant. Preiss. 1, 76 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,129	
P. adunca, Turczaninow in Bull. Soc. Mosc. XXVI, 279 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,130	
P. neurocalyx, Turcz. in Bull. Soc. Mosc. XXVI, 281 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,130	
P. rigida, R. Brown in Benth. Fl. Austr. II, 130 (1864)	—	S.A.	—	—	—	—	—	B.fl.II,130	
P. juniperina, Labillardière, Nov. Holl. pl. spec. I, 102, t. 130 (1804)	—	—	T.	V.	—	—	—	B.fl.II,131	
P. acerosa, R. Brown in Benth. Fl. Austr. II, 131 (1864) ...	—	S.A.	—	—	—	—	—	B.fl.II,131	
P. humilis, Bentham in J. Hook. Fl. Tasm. I, 91 (1860)... ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,131	
P. parviflora, Sieber in De Cand. prodr. II, 111 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.II,132	
P. setulosa, Bentham, Fl. Austr. II, 132 (1864)	—	—	—	—	—	Q.	—	B.fl.II,132	
P. vestita, R. Brown in Ait. hort. Kew. sec. ed. III, 19 (1811)...	W.A.	S.A.	—	—	—	—	—	B.fl.II,132	
P. procumbens, Cunningham in Field's N. S. Wales 347 (1838)...	—	—	—	—	N.S.W.	—	—	B.fl.II,133	
P. hispidula, R. Brown in Bentham, Fl. Austr. II, 133 (1864) ...	—	—	—	—	N.S.W.	—	—	B.fl.II,133	
P. laxiflora, Bentham, Fl. Austr. I, 133 (1864)	—	S.A.	—	V.	—	—	—	B.fl.II,133	
P. largiflorens, F. v. M. in Benth. Fl. Austr. II, 134 (1864) ...	—	S.A.	—	V.	—	—	—	B.fl.II,134	

P. villosa, Willdenow, spec. plant. II, 507 (1799)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,134	
P. foliolosa, Cunningham in Ann. des Wien. Mus. II, 83 (1838)	—	—	—	V.	N.S.W.	—	—	B.fl.II,135	
P. flexilis, Smith in Koen. & Sims, Ann. of Bot. I, 502 (1805) ...	—	—	—	—	N.S.W.	—	—	B.fl.II,135	
P. euchila, De Candolle, prodr. II, 112 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,135	
P. selaginoides, J. Hooker, Fl. Tasm. I, 87 (1860)	—	—	T.	—	—	—	—	B.fl.II,136	
P. densifolia, F. v. M. in Transact. Vict. Inst. 119 (1854) ...	—	S.A.	—	—	—	—	—	B.fl.II,136	
P. elliptica, Smith in Transact. Linn. Soc. IX, 246 (1808) ...	—	—	—	—	N.S.W.	—	—	B.fl.II,136	
P. subspicata, Bentham, Fl. Austr. II, 137 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,137	
P. villifera, Sieber in De Cand. prodr. II, 111 (1825)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,137	
P. involucrata, Bentham, Fl. Austr. II, 138 (1864)	—	S.A.	—	—	—	—	—	B.fl.II,138	
P. Muelleri, Bentham, Fl. Austr. II, 138 (1864)	—	—	—	V.	—	—	—	B.fl.II,138	
P. prostrata, Bentham in J. Hook. Fl. Tasm. I, 89 (1860) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,138	
P. canaliculata, F. v. M. in Transact. Vict. Inst. 119 (1854) ...	—	S.A.	—	V.	—	—	—	B.fl.II,139	
P. fasciculata, Bentham in Ann. des Wien. Mus. II, 82 (1838) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.II,139	
P. tenuifolia, R. Brown in Bot. Mag. t. 2086 (1819)	W.A.	S.A.	T.	V.	—	—	—	B.fl.II,139	M.fr.I,9.
LATROBEA, Meissner in Lehm. plant. Preiss. II, 219 (1847). (Leptocytisus.)									
L. pungens, Bentham, Fl. Austr. II, 140 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,140	M.fr.II,106.
L. genistoides, Meissner in Lehm. plant. Preiss. II, 219 (1847)...	W.A.	—	—	—	—	—	—	B.fl.II,141	
L. Brunonis, Meissner in Lehm. plant. Preiss. II, 219 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.II,141	
L. tenella, Bentham, Fl. Austr. II, 141 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,141	
L. hirtella, Bentham, Fl. Austr. II, 142 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,142	
L. diosmifolia, Bentham, Fl. Austr. II, 142 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,142	
EUTAXIA, R. Brown in Ait. hort. Kew. sec. ed. III, 16 (1811). (Sclerothamnus.)									
E. cuneata, Meissner in Lehm. plant. Preiss. I, 65 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,143	
E. myrtifolia, R. Brown in Ait. hort. Kew. sec. ed. III, 16 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,144	
E. epacridoides, Meissner in Lehm. plant. Preiss. I, 64 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,144	
E. virgata, Bentham in Hueg. enum. 34 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,144	
E. densifolia, Turczaninow in Bull. Soc. Mosc. XXVI, 271 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,145	
E. dillwynioides, Meissner in Lehm. plant. Preiss. I, 63 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,145	
E. parvifolia, Bentham in Hueg. enum. 34 (1837)...	W.A.	—	—	—	—	—	—	B.fl.II,145	
E. empetrifolia, Schlechtendal, Linnaea XX, 667 (1847) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,145	M.fr.I,7.
DILLWYNIA, Smith in Koenig & Sims, Ann. of Bot. I, 510 (1805).									
D. hispida, Lindley in Mitch. Three Exped. II, 251 (1838) ...	—	S.A.	—	V.	—	—	—	B.fl.II,147	
D. ericifolia, Smith in Koen. & Sims, Ann. of Bot. I, 510 (1805)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,147	
D. floribunda, Smith in Koen. & Sims, Ann. of Bot. I, 510 (1805)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,149	
D. Preissii, Bentham, Fl. Austr. II, 149 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,149	
D. brunioides, Meissner in Lehm. pl. Preiss. I, 62 (1844) ...	—	—	—	—	N.S.W.	—	—	B.fl.II,150	
D. juniperina, Sieber in Hueg. enum. pl. N. Holl. occ. 33 (1837)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,150	
D. pungens, Mackay in den Annal. des Wien. Mus. II, 79 (1838)	W.A.	—	—	—	—	—	—	B.fl.II,150	
D. cinerascens, R. Brown in Bot. Mag. t. 2247 (1821)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,151	
D. divaricata, Bentham, Fl. Austr. II, 151 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,151	
D. patula, F. v. M., fragm. IV, 16 (1864)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,151	M.fr.IV,16.
EUCHILOPSIS, F. v. M. in Melbourne Chemist, June (1882).									
E. linearis, F. v. M. in Melb. Chem., June (1882)	W.A.	—	—	—	—	—	—	B.fl.II,67	
PLATYLOBIUM, Smith in Transact. Linn. Soc. II, 350 (1794). (Cheilococca.)									
P. formosum, Smith in Transact. Linn. Soc. II, 350 (1794) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,153	M.fr.IX,157.
P. obtusangulum, Hooker, Bot. Mag. t. 3258 (1833)	—	S.A.	T.	V.	—	—	—	B.fl.II,153	
P. triangulare, R. Brown in Ait. hort. Kew. sec. ed. III, 266 (1811)	—	S.A.	T.	V.	—	—	—	B.fl.II,152	M.fr.IX,158.
BOSSIAEA, Ventenat, Jardin de Cels I, 7, t. 7 (1800). (Lalage, Scottia.)									
B. dentata, Bentham, Fl. Austr. II, 156 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,156	
B. Aquifolium, Bentham, Fl. Austr. II, 157 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,157	
B. strigillosa, Bentham, Fl. Austr. II, 157 (1864)... ...	W.A.	—	—	—	—	—	—	B.fl.II,157	
B. cordigera, Bentham in J. Hook. Fl. Tasm. I, 95, t. 16 (1860)...	W.A.	—	T.	V.	—	—	—	B.fl.II,157	M.fr.III,100;IX,45.
B. lenticularis, Sieber in De Cand. prodr. II, 117 (1825) ...	—	—	—	—	N.S.W.	—	—	B.fl.II,157	
B. Kiamensis, Bentham, Fl. Austr. II, 158 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,158	
B. ornata, F. v. M., fragm. IV, 13 (1863)	W.A.	—	—	—	—	—	—	B.fl.II,158	M.fr.IX,45.
B. eriocarpa, Bentham in Hueg. enum. 36 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,159	M.fr.IX,45.
B. divaricata, Turczaninow in Bull. Soc. Mosc. XXVI, 285 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,159	M.fr.IV,12.
B. calycina, Bentham, Fl. Austr. II, 159 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,159	
B. foliosa, Cunningham in Field's N. S. Wales, 347 (1825) ...	—	—	—	V.	N.S.W.	—	—	B.fl.II,160	M.fr.III,100;IX,45.
B. cinerea, R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,160	M.fr.III,98.
B. biloba, Bentham in Hueg. enum. 36 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,160	M.fr.IX,45.
B. Preissii, Meissner in Lehm. pl. Preiss. I, 82 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,161	M.fr.IX,45.
B. concinna, Bentham, Fl. Austr. II, 161 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,161	
B. carinalis, Bentham in Mitch. Trop. Austr. 290 (1848) ...	—	—	—	—	—	Q.	—	B.fl.II,161	M.fr.IX,45.
B. rupicola, Cunningham in Benth. Fl. Austr. II, 162 (1864) ...	—	—	—	—	—	Q.	—	B.fl.II,162	
B. linophylla, R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812)	W.A.	—	—	—	—	—	—	B.fl.II,162	M.fr.IX,45.
B. disticha, Lindley, Bot. Regist. t. 55 (1841)	W.A.	—	—	—	—	—	—	B.fl.II,162	
B. prostrata, R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,162	M.fr.III,97.
B. Neo-Anglica, F. v. M., fragm. V, 106 (1866)	—	—	—	—	N.S.W.	—	—	—	M.fr.V,106.
B. buxifolia, Cunningham in Field's N. S. Wales, 348 (1825) ...	—	—	—	V.	N.S.W.	—	—	B.fl.II,163	M.fr.I,9;IX,45.
B. Brownii, Bentham, Fl. Austr. II, 163 (1864)	—	—	—	—	—	Q.	—	B.fl.II,163	
B. rhombifolia, Sieber in De Cand., prodr. II, 117 (1825) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,164	M.fr.IX,45.
B. pulchella, Meissner in Lehm. pl. Preiss. I, 84 (1844)... ...	W.A.	—	—	—	—	—	—	B.fl.II,164	
B. microphylla, Smith in Transact. Linn. Soc. IX, 303 (1808) ...	—	—	—	V.	N.S.W.	—	—	B.fl.II,164	M.fr.III,99.
B. peduncularis, Turczaninow in Bull. Soc. Mosc. XXVI, 287 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,165	
B. heterophylla, Ventenat, Jardin de Cels t. 7 (1800)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,165	M.fr.III,97.
B. rufa, R. Brown in Ait. hort. Kew. sec. ed. III, 267 (1812) ...	W.A.	—	—	—	—	—	—	B.fl.II,165	M.fr.IX,45.

| | | |

INDIGOFERA, Royen, pl. hort. Lugd. 372 (1740), from Linné (1737).

I. linifolia, Retzius, observ. bot. IV, 29 (1786)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,195	M.fr.III,101;IX,44.
I. cordifolia, Heyne in Roth, nov. pl. spec. 357 (1821)	—	—	—	—	—	—	N.A.	B.fl.II,196	
I. enneaphylla, Linné, mantissa altera 272 (1771)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,196	M.fr.III,102;IX,44. [44.
I. glandulosa, Willdenow, spec. plant. III, 1227 (1800)	—	—	—	—	—	Q.	—	B.fl.II,196	M.fr.IV,22;VII,106;IX,
I. haplophylla, F. v. M., fragm. III, 102 (1863)	—	—	—	—	—	—	N.A.	B.fl.II,196	M.fr.IX,44.
I. trifoliata, Linné, amoen. acad. IV, 327 (1759)	—	—	—	—	—	Q.	—	B.fl.II,197	M.fr.III,104.
I. trita, Linné fil., suppl. plant. 335 (1781)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,197	M.fr.III,103.
I. parviflora, Heyne in Wall. numeric. list 5457 (1828)	—	—	—	—	—	Q.	N.A.	B.fl.II,197	M.fr.III,103;IX,44.
I. viscosa, Lamarck, Encycl. méth. III, 247 (1789)	—	S.A.	—	—	—	Q.	N.A.	B.fl.II,198	M.fr.III,104;IX,44.
I. hirsuta, Linné, spec. plant. 751 (1753)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,198	M.fr.III,105;IX,44.
I. pratensis, F. v. M., essay on plants of Burdek. Exped. 10 (1860)	—	—	—	—	—	Q.	—	B.fl.II,198	
I. Schultziana, F. v. M., fragm. VII, 105 (1870)	—	—	—	—	—	—	N.A.	—	M.fr.VII,105. [44.
I. monophylla, De Candolle, prodr. II, 222 (1825)	—	S.A.	—	—	—	—	N.A.	B.fl.II,199	M.fr.III,44;VII,106;IX,
I. rugosa, Bentham, Fl. Austr. II, 199 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,199	
I. saxicola, F. v. M. in Benth. Fl. Austr. II, 199 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,199	
I. australis, Willdenow, spec. plant. III, 1235 (1800)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,199	
I. brevidens, Bentham in Mitch. Trop. Austr. 385 (1848)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,200	
I. Baileyi, F. v. M., fragm. IX, 43 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,43.
I. coronillifolia, Cunningham in Benth. Fl. Austr. II, 201 (1864)	—	S.A.	—	—	N.S.W.	—	—	B.fl.II,201	

PTYCHOSEMA, Bentham in Lindl. Bot. Regist. XXV, App. XVI (1839).

P. pusillum, Bentham in Lindl. Bot. Regist. XXV, App. XVI (1839)	W.A.	—	—	—	—	—	—	B.fl.II,201	
P. anomalum, F. v. M., fragm. IX, 62 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,62.
P. trifoliolatum, F. v. M. in Wing's S. Science Record II, 72 (1882)	W.A.	—	—	—	—	—	—	—	

LAMPROLOBIUM, Bentham, Fl. Austr. II, 202 (1864).

L. fruticosum, Bentham, Fl. Austr. II, 202 (1864)	—	—	—	—	—	Q.	—	B.fl.II,202	M.fr.IX,68.
L. megalophyllum, F. v. M., fragm. IX, 67 (1875)	—	—	—	—	—	—	N.A.	—	M.fr.IX,67.

TEPHROSIA, Persoon, synops. pl. II, 328 (1807).

T. coriacea, Bentham, Fl. Austr. II, 204 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,204	M.fr.IX,65&66.
T. lamprolobioides, F. v. M., fragm. IX, 64 (1875)	—	—	—	—	—	—	N.A.	—	M.fr.IX,64.
T. Forrestiana, F. v. M., fragm. XI, 98 (1880)	—	—	—	—	—	—	N.A.	—	M.fr.XI,98.
T. flammea, F. v. M. in Benth. Fl. Austr. II, 204 (1864)	—	—	—	—	—	Q.	—	B.fl.II,204	
T. reticulata, Bentham, Fl. Austr. II, 205 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,205	M.fr.IX,64.
T. crocea, Bentham, Fl. Austr. II, 205 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,205	
T. oblongata, Bentham, Fl. Austr. II, 205 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,205	
T. porrecta, Bentham, Fl. Austr. II, 206 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,206	
T. polyzyga, F. v. M. in Benth. Fl. Austr. II, 206 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,206	
T. graminifolia, F. v. M. in Benth. Fl. Austr. II, 206 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,206	M.fr.IX,66.
T. simplicifolia, F. v. M. in Benth. Fl. Austr. II, 206 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,206	
T. nematophylla, F. v. M., fragm. IX, 63 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.IX,63.
T. leptoclada, Bentham, Fl. Austr. II, 207 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,207	
T. brachycarpa, F. v. M. in Benth. Fl. Austr. II, 207 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,207	
T. uniovulata, F. v. M., fragm. XI, 70 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,70.
T. Stuartii, Bentham, Fl. Austr. II, 207 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,207	
T. conferta, F. v. M., fragm. IX, 65 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.IX,75.
T. eriocarpa, Bentham, Fl. Austr. II, 207 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,207	
T. phaeosperma, F. v. M. in Benth. Fl. Austr. II, 208 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,208	
T. astragaloides, Bentham, Fl. Austr. II, 208 (1864)	—	—	—	—	—	Q.	—	B.fl.II,208	
T. juncea, Bentham, Fl. Austr. II, 208 (1864)	—	—	—	—	—	Q.	—	B.fl.II,208	M.fr.IX,64.
T. filipes, Bentham, Fl. Austr. II, 208 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,208	
T. remotiflora, F. v. M. in Benth. Fl. Austr. II, 209 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,209	
T. oligophylla, Bentham, Fl. Austr. II, 209 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,209	
T. macrocarpa, Bentham, Fl. Austr. II, 209 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,209	
T. purpurea, Persoon, synops. plant. II, 329 (1807)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,209	
T. Bidwilli, Bentham, Fl. Austr. II, 210 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,210	
T. rosea, F. v. M. in Benth. Fl. Austr. II, 211 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,211	

WISTARIA, Nuttall, Gen. of N. Amer. pl. II, 115 (1818). (Millettia.)

W. megasperma, F. v. M., fragm. I, ad hist. natur. 10 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,211	M.fr.I,10.

SESBANIA, Scopoli, Introd. 308 (1777). (Agati.)

S. grandiflora, Persoon, synops. pl. II, 316 (1807)	—	—	—	—	—	—	N.A.	B.fl.II,212	M.fr.II,88,XI,33.
S. Aegyptiaca, Persoon, synops. pl. II, 316 (1807)	—	—	—	—	—	—	N.A.	B.fl.II,212	M.fr.XI,33.
S. aculeata, Persoon, synops. pl. II, 316 (1807)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,213	M.fr.XI,33.
S. simpliciuscula, F. v. M. in Benth. Fl. Austr. II, 213 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,213	
S. brachycarpa, F. v. M., fragm. XI, 32 (1878)	—	—	—	—	—	—	N.A.	—	M.fr.IX,32.

CARMICHAELIA, R. Brown in Bot. Regist. XI, 912 (1825).

C. exsul, F. v. M., fragm. VII, 126 (1871)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,126,193.

CLIANTHUS, Banks & Solander in G. Don, dichlam. pl. II, 468; indicative (1832). (Donia.)

C. Dampieri, Cunning. in Trans. hort. soc. Lond. sec. ser. I, 522 (1835)	W.A.	S.A.	—	—	N.S.W.	—	N.A.	B.fl.II,214	

STREBLORRHIZA, Endlicher, prodr. fl. Norfolk. 92 (1833).

S. speciosa, Endlicher, prodr. fl. Norf. 93 (1833)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.

SWAINSONIA, Salisbury, parad. Londin. t. 28 (1806). (Cyclogyne, Diplolobium.)

S. Greyana, Lindley, Bot. Regist. t. 66 (1846)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,216	M.fr.IX,154;X,7.
S. galegifolia, R. Brown in Ait. hort. Kew. sec. ed. III, 327 (1811)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,217	
S. coronillifolia, Salisbury, paradis. Lond. t. 28 (1806)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,217	M.fr.XI,70.
S. brachycarpa, Bentham, Fl. Austr. II, 217 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,217	M.fr.IX,155.
S. phacoides, Bentham in Mitch. Trop. Austr. 363 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,217	
S. Burkittii, F. v. M. in Benth. Fl. Austr. II, 218 (1864)	—	S.A.	—	—	N.S.W.	—	—	B.fl.II,218	

S. Burkei, F. v. M. in Benth. Fl. Austr. II, 218 (1864)	—	S.A.	—	—	—	—	N.A.	B.fl.II,218	M.fr.IX,155.
S. oligophylla, F. v. M. in Benth. Fl. Austr. II, 219 (1864) ...	—	S.A.	—	—	N.S.W.	—	N.A.	B.fl.II,219	
S. campylantha, F. v. M. in Gregory's pap. rel. to Leichh. Exp. 6(1858)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,219	M.fr.IX,154.
S. occidentalis, F. v. M., fragm. III, 46 (1862)	W.A.	—	—	—	—	—	N.A.	B.fl.II,219	M.fr.III,46.
S. stenodonta, F. v. M., fragm. XI, 70 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,70.
S. gracilis, Bentham, Fl. Austr. II, 220 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,220	
S. plagiotropis, F. v. M., fragm. IX, 153 (1875)	—	—	—	V.	N.S.W.	—	—	—	M.fr.IX,143.
S. procumbens, F. v. M., fragm. III, 46 (1863)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,220	M.fr.IX,154;XI,27.
S. Drummondii, Bentham, Fl. Austr. II, 220 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,220	
S. canescens, F. v. M., fragm. III, 46 (1863)	W.A.	—	—	—	—	—	—	B.fl.II,221	M.fr.IX,154;XI,70.
S. phacifolia, F. v. M. in South Austr. Register (1850)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,221	M.fr.IX,154.
S. oroboides, F. v. M. in Benth. Fl. Austr. II, 222 (1864) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,222	
S. unifoliolata, F. v. M., fragm. VIII, 226 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,226.
S. lessertiifolia, De Candolle, prodr. II, 271 (1825)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,222	M.fr.IX,154.
S. luteola, F. v. M., fragm. I, 75 (1858)	—	—	—	—	—	Q.	—	B.fl.II,223	M.fr.I,75;XI,70.
S. parviflora, Bentham, Fl. Austr. II, 223 (1864)	—	—	—	—	—	Q.	—	B.fl.II,223	
S. monticola, Cunningham in A. Gray's Bot. Wilk. Exp. I, 411(1854)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,223	M.fr.IX,155.
S. microphylla, A. Gray, Bot. Wilk. Exped. I, 410 (1854) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,223	M.fr.IX,155.
S. laxa, R. Brown in Sturt's Centr. Austr. App. 18 (1849) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,224	
S. colutoides, F. v. M., fragm. X, 6 (1876)...	—	S.A.	—	—	—	—	—	—	M.fr.X,6.
S. Fraseri, Bentham, Fl. Austr. II, 224 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,224	
S. Macullochiana, F. v. M., fragm. VII, 25 (1869)	W.A.	—	—	—	—	—	N.A.	—	M.fr.IX,154.
GLYCYRRHIZA, Tournefort, inst. 389, t. 210 (1700), from Dioscorides and Plinius. (Clidanthera.)									
G. psoraleoides, Bentham, Fl. Austr. II, 225 (1864)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,225	M.fr.III,45.
ORMOCARPUM, Palisot, Fl. d'Ovar. I, 95, t. 58 (1805).									
O. sennoides, De Candolle, prodr. II, 315 (1825)	—	—	—	—	—	Q.	—	B.fl.II,226	
AESCHYMOMENE, Linné, gen. pl. 350 (1737).									
A. Indica, Linné, spec. plant. 713 (1753)	W.A.	—	—	—	—	Q.	N.A.	B.fl.II,227	
A. falcata, De Candolle, prodr. II, 322 (1825)	—	—	—	—	—	Q.	—	B.fl.II,227	
A. Americana, Linné, spec. plant. 713 (1753)	—	—	—	—	—	Q.	—	—	
SMITHIA, Aiton, hort. Kew. III, 496 (1789).									
S. conferta, Smith in Rees, Cyclop. XXXIII (1816)	—	—	—	—	—	Q.	—	B.fl.II,228	
S. sensitiva, Aiton, hort. Kew. III, 496, t. 13 (1789)	—	—	—	—	—	Q.	—	—	M.fr.VII,27;IX,193.
ZORNIA, Gmelin, Syst. Nat. 1076 (1791).									
Z. diphylla, Persoon, synops. II, 318 (1807)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,228	
Z. chaetophora, F. v. M. in Transact. phil. Inst. Vict. III, 56 (1857)	—	—	—	—	—	—	N.A.	B.fl.II,229	
DESMODIUM, Desvaux, Journ. de Bot. III, 122 (1813). (Dicerma, Phyllodium, Pteroloma.)									
D. umbellatum, De Candolle, prodr. II, 325 (1825)	—	—	—	—	—	Q.	—	B.fl.II,230	M.fr.IX,66.
D. pulchellum, Bentham, Fl. Hongk. 83 (1861)	—	—	—	—	—	—	N.A.	B.fl.II,231	
D. biarticulatum, F. v. M., fragm. II, 121 (1861)...	—	—	—	—	—	Q.	N.A.	B.fl.II,231	M.fr.IX,66.
D. acanthocladum, F. v. M., fragm. II, 122 (1861)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,231	M.fr.IX,66.
D. triquetrum, De Candolle, prodr. II, 326 (1825)	—	—	—	—	—	Q.	—	—	M.fr.IX,66.
D. Gangeticum, De Candolle, prodr. II, 327 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.II,232	
D. brachypodum, A. Gray, Bot. Wilk. Exped. 434 (1854) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,232	
D. varians, Endlicher in Ann. des Wien. Mus. I, 185 (1839) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,232	
D. flagellare, Bentham, Fl. Austr. II, 233 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,233	
D. rhytidophyllum, F. v. M. in Benth. Fl. Austr. II, 223 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,233	
D. campylocaulon, F. v. M. in Benth. Fl. Austr. II, 233 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,233	
D. nemorosum, F. v. M. in Benth. Fl. Austr. II, 234 (1864) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,234	
D. neurocarpum, Bentham, Fl. Austr. II, 234 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,234	
D. trichostachyum, Bentham, Fl. Austr. II, 234 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,234	
D. polycarpum, De Candolle, prodr. II, 334 (1825)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,235	M.fr.IX,66.
D. trichocaulon, De Candolle, prodr. II, 335 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,235	
D. Muelleri, Bentham, Fl. Austr. II, 235 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,235	
D. reniforme, De Candolle, prodr. II, 327 (1825)	—	—	—	—	—	Q.	N.A.	—	M.fr.XII.
D. parvifolium, De Candolle, prodr. II, 334 (1825)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,235	M.fr.IX,66. [X,120.
D. dependens, Blume in Miq. Fl. Ind. Batav. I, 249 (1855) ...	—	—	—	—	—	Q.	—	—	M.fr.VIII,225;IX,66&195
PYCNOSPORA, R. Brown in Wight & Arnott, prodr. I, 197 (1834).									
P. hedysaroides, R. Brown in Wight & Arn., prodr. 197 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.II,236	
URARIA, Desvaux, Journ. de bot. III, 122 (1813).									
U. picta, Desvaux, Journ. de Bot. III, 122 (1813)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,237	
U. cylindracea, Bentham, Fl. Austr. II, 237 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,237	
U. lagopoides, De Candolle, prodr. II, 324 (1825)	—	—	—	—	—	Q.	—	B.fl.II,237	
LOUREA, Necker, elem. bot. III, 17 (1790).									
L. obcordata, Desvaux, Journ. de Bot. III, 122 (1813)	—	—	—	—	—	—	N.A.	B.fl.II,238	
ALYSICARPUS, Necker, elem. bot. III, 15 (1790).									
A. vaginalis, De Candolle, prodr. II, 353 (1825)	—	—	—	—	—	Q.	—	B.fl.II,239	
A. longifolius, Wight & Arnott, prodr. 223 (1834)	—	—	—	—	—	—	N.A.	B.fl.II,239	
A. cylindricus, Desvaux in Annal. Soc. Linn. Par. 301 (1825) ..	—	—	—	—	—	Q.	N.A.	B.fl.II,239	
LESPEDEZA, Cl. Richard in Michaux, fl. bor. am. II, 70 (1803).									
L. cuneata, G. Don, gen. syst. II, 307 (1832)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,240	
L. lanata, Bentham, Fl. Austr. II, 241 (1864)	—	S.A.	—	—	—	—	—	B.fl.II,241	
CLITORIA, Linné, gen. plant. 216 (1737), from Petiver.									
C. australis, Bentham, Fl. Austr. II, 242 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,242	

GLYCINE, Linné, gen. pl. 349 (1737). (Leptolobium, Leptocyamus.)

G. falcata, Bentham, Fl. Austr. II, 243 (1864)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,243				
G. clandestina, Wendland, Bot. Reob. 54 (1798)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,243				
G. Latrobeana, Bentham, Fl. Austr. II, 244 (1864)	—	S.A.	T.	V.	—	—	—	B.fl.II,244				
G. tabacina, Bentham, Fl. Austr. II, 244 (1864)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,244				
G. sericea, Bentham, Fl. Austr. II, 245 (1864)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,245				
G. tomentosa, Bentham, Fl. Austr. II, 245 (1864)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,245				

KENNEDYA, Ventenat, Jardin de la Malmaison II, t. 104, 106 (1804). (Caulinia, Moench 1802, Hardenbergia, Physolobium, Zichya, Amphodus.)

K. nigricans, Lindley, Bot. Regist. t. 1715 (1835)	W.A.	—	—	—	—	—	—	B.fl.II,249	M.fr.VII,127.			
K. Beckxiana, F. v. M., fragm. XI, 98 (1880)	W.A.	—	—	—	—	—	—	—	M.fr.XI,98.			
K. rubicunda, Ventenat, Jard. de la Malm. t. 104 (1804) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,249	M.fr.VII,127.			
K. procurrens, Bentham in Mitch. trop. Austr. 365 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,249				
K. prorepens, F. v. M., fragm. VIII, 225 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,225.			
K. prostrata, R. Brown in Ait. hort. Kew. sec. ed. IV, 299 (1812)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,250	M.fr.VII,128.			
K. bracteata, Gaudichaud in Freyc. voy. 286, t. 113 (1826) ...	W.A.	—	—	—	—	—	—	B.fl.II,250				
K. eximia, Lindley in Paxt. Mag. XVI, 35	W.A.	—	—	—	—	—	—	B.fl.II,250				
K. microphylla, Meissner in Lehm. pl. Preiss. I, 91 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,251				
K. parviflora, Meissner in Lehm. pl. Preiss. I, 91 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,251				
K. Stirlingii, Lindley, Bot. Regist. t. 1845 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,252				
K. glabrata, Lindley, Bot. Regist. t. 1838 (1837)...	W.A.	—	—	—	—	—	—	B.fl.II,252	M.fr.VII,128.			
K. macrophylla, F. v. M., fragm. IV, 79 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,252	M.fr.IV,79.			
K. coccinea, Ventenat, Jard. de la Malm. t. 105 (1804)	W.A.	—	—	—	—	—	—	B.fl.II,250	M.fr.VII,128.			
K. monophylla, Ventenat, Jard. de la Malm. t. 106 (1804) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,246	M.fr.VII,128.			
K. Comptoniana, Link, enum. pl. hort. bot. Berol. II, 235 (1822)	W.A.	—	—	—	—	—	—	B.fl.II,247	M.fr.VII,128.			
K. retusa, F. v. M., fragm. V, 106 (1866)	—	—	—	—	—	Q.	—	B.fl.II,247	M.fr.VII,128.			

ERYTHRINA, Linné, gen. pl. 216 (1737).

E. vespertilio, Bentham in Mitch. Trop. Austr. 218 (1848) ...	—	S.A.	—	—	—	Q.	N.A.	B.fl.II,253				
E. Indica, Lamarck, Encyclop. méthod. II, 391 (1786)	—	—	—	—	—	Q.	N.A.	B.fl.II,253				

STRONGYLODON, T. Vogel in Linnaea X, 585 (1836).

S. ruber, T. Vogel in Linnaea X, 585 (1836)	—	—	—	—	—	Q.	—	—				

MUCUNA, Marcgraf, hist. nat. Brazil. 18 (1648).

M. gigantea, De Candolle, prodr. II, 405 (1825)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,254				

GALACTIA, P. Browne, Civil & Nat. Hist. of Jamaica 298 (1756).

G. tenuiflora, Wight & Arnott, prodr. fl. penins. Ind. 206 (1834)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,255				
G. Muelleri, Bentham, Fl. Austr. II, 255 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,255				

CANAVALIA, De Candolle, prodr. II, 403 (1825). (Canavali.)

C. obtusifolia, De Candolle, prodr. II, 404 (1825)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,256				

PHASEOLUS, Tournefort, inst. 412, t. 232 (1700), from Dioscorides & Columella.

P. vulgaris, de l'Obel, stirp. icon. 59 (1581)	—	—	—	—	—	—	N.A.	B.fl.II,257				
P. Truxillensis, Humb. Bonpl. & Kunth, nov. gen. Amer. VI, 451 (1823)	—	—	—	—	—	Q.	N.A.	B.fl.II,257				
P. Mungo, Linné, mantissa plantar. 101 (1767)	—	—	—	—	—	Q.	N.A.	B.fl.II,257				

VIGNA, Savi, Dissert. 6 (1824). (Callicystus.)

V. vexillata, Bentham in Mart. Fl. Bras. Papil. 193, t. 50 (1862)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,258				
V. lutea, A. Gray, Bot. Wilk. Exped. I, 454 (1854)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,259				
V. luteola, Bentham in Mart. Fl. Bras. Papil. 194, t. 50 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,260				
V. lanceolata, Bentham in Mitch. trop. Austr. 350 (1848) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,260				

DOLICHOS, Linné, gen. plant. 222 (1737).

D. biflorus, Linné, spec. plant. 727 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.II,261				

DUNBARIA, Wight & Arnott, prodr. I, 258 (1834).

D. conspersa, Bentham in plant. Junghuhn. (1851)	—	—	—	—	—	Q.	—	B.fl.II,262				

CAJANUS, De Candolle, Catal. hort. Monsp. 85 (1813). (Cajan, Atylosia.)

C. marmoratus, F. v. M.; Atylosia, Benth., Fl. Austr. II, 263 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,263				
C. scarabeoides, F. v. M.; Atyl., Benth. in pl. Jungh. I, 242 (1851)	—	—	—	—	—	Q.	—	B.fl.II,263				
C. reticulatus, F. v. M.; Atylosia, Benth., Fl. Austr. II, 263 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,263				
C. grandifolius, F. v. M., Essay on pl. Burdekin Exped. 9 (1860)	—	—	—	—	—	Q.	N.A.	B.fl.II,264				
C. confertiflorus, F. v. M., Essay on pl. Burdekin Exped. 9 (1860)	—	—	—	—	—	Q.	—	B.fl.II,264				
C. cinereus; Atylosia, F. v. M. in Benth. Fl. Austr. II, 264 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,264				

RHYNCHOSIA, Loureiro, Fl. Cochinch. II, 460 (1790).

R. rhomboidea, F. v. M. in Benth. Fl. Austr. II, 265 (1864) ...	—	—	—	—	—	—	N.A.	B.fl.II,265				
R. acutifolia, Bentham, Fl. Austr. II, 266 (1864)...	—	—	—	—	—	—	N.A.	B.fl.II,266				
R. rostrata, Bentham, Fl. Austr. II, 266 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,266				
R. Cunninghamii, Bentham, Fl. Austr. II, 266 (1864)	—	—	—	—	—	Q.	—	B.fl.II,266				
R. minima, De Candolle, prodr. II, 385 (1825)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,267				

ERIOSEMA, De Candolle, prodr. II, 388 (1825).

E. Chinense, T. Vogel in Meyen's Beitr. zur Bot. 31 (1843) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,268				

FLEMINGIA, Roxburgh in Ait. hort. Kew. sec. ed. IV, 349 (1812).

F. lineata, Roxburgh in Ait. hort. Kew. sec. ed. IV, 350 (1812)	—	—	—	—	—	Q.	N.A.	B.fl.II,268				
F. pauciflora, Bentham, Fl. Austr. II, 269 (1864)...	—	—	—	—	—	—	N.A.	B.fl.II,269				
F. parviflora, Bentham, Fl. Austr. II, 269 (1864)...	—	—	—	—	—	Q.	—	B.fl.II,269				
F. involucrata, Bentham in pl. Jungh. I, 246 (1851)	—	—	—	—	—	Q.	—	B.fl.II,269				

ABRUS, Rauwolf in Dalechamps, hist. general. plant. append. 193 (1587).

A. precatorius, Linné, syst. natur. ed. XII, 533 (1767)	—	—	—	—	—	Q.	N.A.	B.fl.II,270				

DALBERGIA, Linné, fil. suppl. 52 (1781).

D. densa, Bentham in Hook. Lond. Journ. II, 217 (1843) ...	—	—	—	—	—	Q.	—	B.fl.II,271	

LONCHOCARPUS, Humboldt, Bonpland & Kunth, nov. gen. VI, 383 (1823).

L. Blackii, Bentham, Fl. Austr. II, 271 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,272	

DERRIS, Loureiro, Fl. Cochinch. II, 432 (1790). (Brachypterum.)

D. scandens, Bentham in Journ. Linn. soc. IV, suppl. 103 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,272	
D. uliginosa, Bentham in pl. Jungh. I, 252 (1851)	—	—	—	—	—	Q.	N.A.	B.fl.II,272	

PONGAMIA, Lamarck in Ventenat. Jardin de la Malmaison, t. 28 (1803).

P. glabra, Ventenat, Jard. Malm. t. 28 (1803)	—	—	—	—	—	Q.	N.A.	B.fl.II,273	

SOPHORA, Linné, gen. plant. 125 (1737).

S. tomentosa, Linné, spec. plant. 373 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,274	M.fr.VII,27.
S. Fraseri, Bentham, Fl. Austr. II, 274 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,274	M.fr.V,31;VII,27.
S. tetraptera, J. Miller, icon. plant. t. 1 (1780)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,26.

PODOPETALUM, F. v. M. in Melbourne Chemist, June (1882).

P. Ormondi, F. v. M., fragm. XII (inedit.)	—	—	—	—	–	Q.	—	—	

CASTANOSPERMUM, Cunningham & Fraser in Hook. bot. misc. I, 241, t. 51 (1830).

C. Australe, Cunningh. & Fraser in Hook. bot. misc. I, 241, t. 51 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,275	M.fr.VII,27.

BARKLYA, F. v. M., fragm. I, 109 (1859).

B. syringifolia, F. v. M. in Journ. Linn. Soc. III, 158 (1859) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,275	M.fr.I,109;III,163;VIII,27.

CAESALPINA, Plumier, nov. pl. Amer. gen. 28, t. 9 (1703), from Ammand. (Guilandina, Caesalpinia.)

C. Bonducella, Fleming in Asiat. Research XI, 159 (1801) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,276	
C. Nuga, W. T. Aiton, hort. Kew., sec. ed. III, 32 (1811) ...	—	—	—	—	—	Q.	—	B.fl.II,277	M.fr.X,7.
C. sepiaria, Roxburgh, Fl. Indic. II, 360 (1832)	—	—	—	—	—	Q.	—	B.fl.II,277	M.fr.X,7.

MEZONEURON, Desfontaines in Mém. du Mus. IV, 245, t. 10 et 11 (1818).

M. brachycarpum, Bentham, Fl. Austr. II, 278 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,278	M.fr.X,7.
M. Scortechinii, F. v. M. in Wing's South. Sc. Record II, 73 (1882)	—	—	—	—	N.S.W.	Q.	—	—	

PTEROLOBIUM, R. Brown in Salt. Abyssin. 65 (1814).

P. nitens, F. v. M. in Bentham, Fl. Austr. II, 279 (1864) ...	—	—	—	—	—	Q.	—	B.fl.II,279	M.fr.X,7.

PELTOPHORUM, Vogel in Linnaea XI, 406 (1837).

P. ferrugineum, Bentham, Fl. Austr. II, 279 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,279	M.fr.X,7.

CASSIA, Tournefort, inst. 619, t. 392 (1700), from Plumier (1693). (Cathartocarpus.)

C. Brewsteri, F. v. M., fourth ann. Rep. 17 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,282	M.fr.X,8.
C. laevigata, Willdenow, enum. pl. hort. bot. Berol. I, 441 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,282	M.fr.IV,14;X,8.
C. Sophera, Linné, spec. plant. 379 (1853)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,283	M.fr.IV,14;X,9.
C. laxiflora, Bentham, Fl. Austr. II, 283 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,283	
C. magnifolia, F. v. M., fragm. I, 166 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.II,283	M.fr.I,166;X,9.
C. venusta, F. v. M., fragm. I, 165 (1859)...	—	S.A.	—	—	—	Q.	N.A.	B.fl.II,284	M.fr.I,165;X,9.
C. notabilis, F. v. M., fragm. III, 28 (1862)	W.A.	S.A.	—	–	—	—	N.A.	B.fl.II,284	M.fr.III,28;X,9.
C. pleurocarpa, F. v. M., fragm. I, 223 (1859)	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,284	M.fr.I,223;X,9.
C. glauca, Lamarck, Encycl. meth. I, 647 (1783)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,285	M.fr.IV,13;X,9.
C. retusa, Solander in Linnaea XI, 72 (1842)	—	—	—	—	—	Q.	—	B.fl.II,285	
C. australis, Sims in Bot. Mag. t. 2676 (1826)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.II,285	M.fr.X,9.
C. Chatelainiana, Gaudichaud in Freyc. voy. 485, t. 111 (1826)	W.A.	—	—	—	—	—	N.A.	B.fl.II,286	
C. glutinosa, De Candolle, prodr. II, 495 (1825)	—	S.A.	—	—	—	—	N.A.	B.fl.II,286	
C. pruinosa, F. v. M., fragm. III, 48 (1862)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,286	M.fr.III,48.
C. circinata, Bentham in Mitch. trop. Austr. 384 (1848)... ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,286	M.fr.I,165.
C. phyllodinea, R. Brown, App. to Sturt's Centr. Austr. 15 (1849)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,287	
C. eremophila, Cunningh. in T. Vogel, gen. Cass. synops. 47 (1837)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,287	
C. artemisioides, Gaudichaud in De Cand. prodr. II, 495 (1825)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,288	
C. Sturtii, R. Brown, App. to Sturt's Centr. Austr. 14 (1849) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,288	
C. desolata, F. v. M. in Linnaea XXV, 387 (1852)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,289	
C. oligophylla, F. v. M., fragm. III, 49 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,289	M.fr.III,49.
C. cardiosperma, F. v. M., fragm. X, 50 (1877)	W.A.	—	—	—	—	—	—	—	M.fr.X,50.
C. oligoclada, F. v. M., fragm. III, 49 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,289	M.fr.III,49;X,9.
C. heptanthera, F. v. M., fragm. X, 8 (1876)	—	—	—	—	—	—	N.A.	—	M.fr.X,8.
C. leptoclada, Bentham, Fl. Austr. II, 290 (1864)...	—	—	—	—	—	—	N.A.	B.fl.II,290	
C. Absus, Linné, spec. plant. 376 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.II,290	M.fr.III,50;X,9.
C. pumila, Lamarck, Encycl. méth. I, 620 (1783)	—	—	—	—	—	Q.	—	B.fl.II,290	M.fr.III,47;X,10.
C. concinna, Bentham, Fl. Austr. II, 291 (1864)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,291	M.fr.X,10.
C. mimosoides, Linné, spec. plant. 379 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,291	M.fr.III,48;X,10.

PETALOSTYLIS, R. Brown, App. to Sturt's Centr. Austr. 79 (1849).

P. labichoides, R. Brown, App. to Sturt's Centr. Austr. 17 (1849)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,292	M.fr.X,7.

LABICHEA, Gaudichaud in De Cand. prodr. II, 507 (1825).

L. cassioides, Gaudichaud in De Cand. prodr. II, 507 (1825) ...	W.A.	—	—	—	—	—	—	B.fl.II,292	
L. nitida, Bentham, Fl. Austr. II, 293 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,293	M.fr.X,7.
L. lanceolata, Bentham in Hueg. enum. 41 (1837)...	W.A.	—	—	—	—	—	—	B.fl.II,293	M.fr.X,7.
L. rupestris, Bentham in Mitch. Trop. Austr. 342 (1848) ...	—	—	—	—	—	Q.	—	B.fl.II,293	M.fr.X,7.
L. punctata, Bentham in Bot. Regist. XXV, App. XV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.II,294	M.fr.X,8.
L. Buettneriana, F. v. M. in Melb. Chemist, June (1882) ...	—	—	—	—	—	Q.	—	—	

TAMARINDUS, Tournefort, inst. 660, t. 445. App. (1700) from C. Bauhin (1623).

T. Indicus, Linné, spec. plant. 34 (1853)	—	—	—	—	—	—	N.A.	B.fl.II,294	

BAUHINIA, Plumier, nov. gen. pl. Amer. 22, t. 13 (1703).

B. Leichhardtii, F. v. M. in Transact. Vict. Inst. III, 50 (1858)	—	S.A.	—	—	—	—	N.A.	B.fl.II,295	M.fr.X,8.
B. Carronii, F. v. M. in Transact. Vict. Inst. III, 49 (1858) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,295	M.fr.X,8.
B. Hookeri, F. v. M. in Transact. Vict. Inst. III, 51 (1858) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,295	M.fr.X,8.

B. Gilesii, F. v. M. and Bailey in Wing's S. Sc. Record, Jul. (1882)	—	—	—	—	—	—	N.A.	—		
AFZELIA, Smith in Transact. Linn. Soc. IV, 221 (1798).										
A. bijuga, A. Gray, Bot. Wilk. Expl. Exped. 467, t. 51 (1854)...	—	—	—	—	—	Q.	—	—		
CYNOMETRA, Linné in Act. Soc. Ups. 78 (1741).										
C. ramiflora, Linné, spec. plant. 382 (1753)	—	—	—	—	—	Q.	—	B.fl.II,296		
ERYTHROPHLAEUM, Afzelius in Tuckey's Congo 430 (1818). (Laboucheria.)										
E. Laboucherii, F. v. M. in Benth. Fl. Austr. II, 297 (1864) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,297		
ENTADA, Adanson, Familles des plantes II, 318 (1763).										
E. Pursaetha, De Cand., mémoir. sur la fam. des Légum. 12 (1825)	—	—	—	—	—	Q.	—	B.fl.II,298		
ADENANTHERA, Royen in Linné, coroll. 7 (1737).										
A. pavonina, Linné, spec. plant. 384 (1753)	—	—	—	—	—	Q.	—	—	M.fr.V,30.	
A. abrosperma, F. v. M., fragm. V, 30 (1865)	—	—	—	—	—	Q.	—	B.fl.II,298	M.fr.V,31.	
NEPTUNIA, Loureiro, Fl. Cochinch. II, 653 (1790). (Dichrostachys.)										
N. cinerea, F.v.M.; Dichrostachys, Wight & Arnott, prodr. 271 (1834)	—	—	—	—	—	—	N.A.	B.fl.II,299		
N. spicata, F. v. M., fragm. III, 151 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,299	M.fr.III,151.	
N. gracilis; Bentham in Hook. journ. of Bot. IV, 355 (1842) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,300		
N. monosperma, F. v. M. in Benth. Fl. Austr. II, 299 (1864) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,300		
ACACIA, Tournefort, inst. 605, t. 375 (1700) from Dioscorides & Plinius. (Vachellia, Chithonanthus, Tetracheilus, Mimosa partly.)										
I. Alatae.										
A. glaucoptera, Bentham in Schlecht. Linnaea XXVI, 604 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,320		
A. alata, R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813) ...	W.A.	—	—	—	—	—	—	B.fl.II,320		
A. diptera, Lindley, Bot. Regist. XXV. App. XV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.II,321		
A. stenoptera, Bentham in Hook. Lond. Journ. I, 325 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,321		
II. Continuae.										
A. incurva, Bentham in Hook. Lond. Journ. I, 325 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,322		
A. trigonophylla, Meissner in Lehm. pl. Preiss. II, 199 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.II,322	M.fr.IV,3.	
A. continua, Bentham, Fl. Austr. II, 322 (1864)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,322		
A. Peuce, F. v. M., fragm. III, 151 (1863)...	—	S.A.	—	—	—	Q.	—	B.fl.II,323	M.fr.III.151.	
III. Pungentes.										
A. spinescens, Bentham in Hook. Lond. Journ. I, 323 (1842) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,323		
A. cochlearis, H. L. Wendland, comm. de Acac. aphyll. 15 (1820)	W.A.	S.A.	—	—	—	—	—	B.fl.II,324		
A. lanigera, Cunningham in Field's N.S. Wales, 345 (1825) ...	—	—	—	V.	N.S.W.	—	—	B.fl.II,324		
A. phlebocarpa, F. v. M. in Journ. Linn. Soc. III, 119 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,325		
A. trinervata, Sieber in De Cand. prodr. II, 451 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,325		
A. colletioides, Cunningham in Hook. Lond. Journ. I, 336 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,325	M.fr.IV,4.	
A. striatula, Bentham in Hook. Lond. Journ. I, 336 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,326		
A. campylophylla, Bentham in Schlecht. Linnaea XXVI, 605 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,326		
A. teretifolia, Bentham in Hook. Lond. Journ. I, 326 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,326		
A. sulcata, R. Brown in Ait. hort. Kew. sec. ed. V, 460 (1813)	W.A.	—	—	—	—	—	—	B.fl.II,327		
A. costata, Bentham in Hook. Lond. Journ. I, 339 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,327		
A. barbinervis, Bentham in Hook. Lond. Journ. I, 326 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,327		
A. ataxiphylla, Bentham in Schlecht. Linnaea XXVI, 605 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,328		
A. Baxteri, Bentham in Hook. Lond. Journ. I, 327 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,328		
A. aureonitens, Lindley, Bot. Regist. XXV, App. XV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.II,328		
A. quadrisulcata, F. v. M., fragm. III, 127 (1863)	W.A.	—	—	—	—	—	—	B.fl.II,328	M.f.III,127.	
A. erioclada, Bentham in Schlecht. Linnaea XXVI, 606 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,329		
A. siculiformis, Cunningham in Hook. Lond. Journ. I, 337 (1842)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,329		
A. patens, F. v. M. in Journ. Linn. Soc. III, 120 (1858)... ...	—	—	—	—	—	—	N.A.	B.fl.II,329	M.fr.III,46.	
A. laricina, Meissner in Lehm. pl. Preiss. I, 6 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,330		
A. tetragonophylla, F. v. M. in Journ. Linn. Soc. III, 121 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.fl.II,330	M.fr.IV,3.	
A. genistoides, Cunningham in Benth. Fl. Austr. II, 330 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,330		
A. spacelata, Bentham in Hook. Lond. Journ. I, 338 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,331		
A. ingrata, Bentham, Fl. Austr. II, 331 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,331		
A. juniperina, Willdenow, spec. plant. IV, 1049 (1805)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,331		
A. asparagoides, Cunningham in Field's N.S. Wales 343 (1825)...	—	—	—	—	N.S.W.	—	—	B.fl.II,332		
A. tenuifolia, F. v. M. in Transact. phil. Soc. Vict. I, 37 (1854)	—	—	—	V.	—	—	—	B.fl.II,332		
A. diffusa, Edwards, Bot. Regist. t. 634 (1822)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,332		
A. rupicola, F. v. M. in Schlecht. Linnaea XXVI, 610 (1853) ...	—	S.A.	—	V.	—	—	—	B.fl.II,333		
IV. Calamiformes.										
A. volubilis, F. v. M., fragm. X, 98 (1877)...	W.A.	—	—	—	—	—	—	—	M.fr.X,98.	
A. tetragonocarpa, Meissner in Lehm. pl. Preiss. I, 4 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,336		
A. restiacea, Bentham in Hook. Lond. Journ. I, 3 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,336		
A. squamata, Lindley, Bot. Regist. XXV, App. XV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.II,336		
A. brachyphylla, Bentham in Schlecht. Linnaea XXVI, 615 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,337		
A. Bynoeana, Bentham in Schlecht. Linnaea XXVI, 614 (1853)	—	—	—	—	—	—	N.A.	B.fl.II,337	M.fr.IV,9.	
A. triptycha, F. v. M. in Benth. Fl. Austr. II, 337 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.II,337		
A. leptoneura, Bentham in Hook. Lond. Journ. I, 341 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,337		
A. rigens, Cunningham in Loudon, hort. Brit. II, 406 (1830) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,337		
A. papyrocarpa, Bentham, Fl. Austr. II, 338 (1864)	W.A.	S.A.	—	—	—	—	—	B.fl.II,338		
A. Gilesiana, F. v. M. in Melb. Chemist, July (1882) ...	—	S.A.	—	—	—	—	—	—		
A. sessiliceps, F. v. M. in Melb. Chemist, July (1882)	—	S.A.	—	—	—	—	—	—		
A. pugioniformis, Wendl., comm. de Acac. aphyll. 38, t. 9 (1820)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,338		
A. juncifolia, Bentham in Hook. Lond. Journ. I, 341 (1842) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,339		
A. calamifolia, Sweet in Bot. Regist. t. 839 (1824)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,339		
A. scirpifolia, Meissner in Bot. Zeitung 10 (1855)...	W.A.	—	—	—	—	—	—	B.fl.II,339		
A. extensa, Lindley, Bot. Regist. XXV, App. XV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.II,340		

A. gonophylla, Bentham in Schlecht. Linnaea XXVI, 613 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,340	
A. ericifolia, Bentham in Hook. Lond. Journ. I, 345 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,340	
A. uncinella, Bentham in Schlecht. Linnaea XXVI, 613 (1853)...	W.A.	—	—	—	—	—	—	B.fl.II,341	
A. oxyclada, F. v. M. in Benth. Fl. Austr. II, 341 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.II,341	
V. Brunioidae.									
A. cedroides, Bentham in Schlecht. Linnaea XXVI, 615 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,341	
A. lycopodifolia, Cunningham in Hook. icon. plant. t. 172 (1837)	—	S.A.	—	—	—	Q.	N.A.	B.fl.II,342	
A. spondylophylla, F. v. M., fragm. VIII, 243 (1874)	—	S.A.	—	—	—	—	N.A.	—	M.fr.243.
A. galioides, Bentham in Hook. Lond. Journ. I, 344 (1842) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,342	
A. Baueri, Bentham in Hook. Lond. Journ. I, 344 (1842) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,342	M.fr.XI,33.
A. subternata, F. v. M. in Journ. Linn. Soc. III, 124 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,343	
A. minutifolia, F. v. M., fragm. VIII, 243 (1874)...	—	S.A.	—	—	—	—	—	—	M.fr.X,243.
A. brunioides, Cunningham in G. Don, gen. syst. II, 404 (1832)	—	—	—	—	—	Q.	—	B.fl.II,343	
A. conferta, Cunningham in Hook. Lond. Journ. I, 345 (1842)...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,343	
VI. Uninerves.									
A. scabra, Bentham in Schlecht. Linnaea XXVI, 605 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,344	
A. nodiflora, Bentham in Schlecht. Linnaea XXVI, 621 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,344	
A. spinosissima, Bentham in Schlecht. Linnaea XXVI, 621 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,344	
A. ulicina, Meissner in Lehm. pl. Preiss. II, 202 (1846)	W.A.	—	—	—	—	—	—	B.fl.II,345	
A. erinacea, Bentham in Hook. Lond. Journ. I, 360 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,345	
A. Huegelii, Bentham in Hueg. enum. 42 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,345	
A. nervosa, De Candolle, Mem. sur la fam. des Légum. 444 (1825)	W.A.	—	—	—	—	—	—	B.fl.II,346	
A. obovata, Bentham in Hook. Lond. Journ. I, 329 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,346	
A. congesta, Bentham in Hook. Lond. Journ. I, 327 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,346	
A. dermatophylla, Bentham, Fl. Austr. II, 346 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,346	
A. aspera, Lindley in Mitch. Three Exped. II, 139 (1838) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,347	
A. armata, R. Brown in Ait. hort. Kew. sec. ed. V, 463 (1813)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,347	
A. idiomorpha, Cunningham in Hook. Lond. Journ. I, 329 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,348	
A. Shuttleworthii, Meissner in Lehm. pl. Preiss. I, 7 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,348	
A. Gregorii, F. v. M., fragm. III, 47 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,348	M.fr.III,47.
A. pilosa, Bentham in Schlecht. Linnaea XXVI, 607 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,348	
A. crispula, Bentham in Schlecht. Linnaea XXVI, 607 (1853) ..	W.A.	—	—	—	—	—	—	B.fl.II,349	
A. crassistipulea, Bentham in Hook. Lond. Journ. I, 326 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,349	
A. hastulata, Smith in Rees' Cyclopaed. XXXIX, Suppl. (1819)	W.A.	—	—	—	—	—	—	B.fl.II,349	
A. horridula, Meissner in Lehm. pl. Preiss. I, 9 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,350	
A. divergens, Bentham in Hook. Lond. Journ. I, 331 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,350	
A. vomeriformis, Cunningham in Hook. Lond. Journ. I, 332 (1842)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,350	
A. plagiophylla, F. v. M. in journ. Linn. Soc. III, 131 (1858) ..	—	—	—	—	N.S.W.	Q.	—	—	
A. biflora, R. Brown in Ait. hort. Kew. sec. ed. V, 463 (1813)	W.A.	—	—	—	—	—	—	B.fl.II,351	
A. decipiens, R. Brown in Ait. hort. Kew. sec. ed. V, 463 (1813)	W.A.	—	—	—	—	—	—	B.fl.II,351	
A. cuneata, Bentham in Hueg. enumer. 42 (1837)...	W.A.	—	—	—	—	—	—	B.fl.II,351	
A. dilatata, Bentham in Schlecht. Linnaea XXVI, 608 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,352	
A. bidentata, Bentham in Hook. Lond. Journ. I, 333 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,352	
A. acanthoclada, F. v. M., fragm. III, 127 (1863)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,352	M.fr.III,127.
A. obliqua, Cunningham in Hook. Lond. Journ. I, 334 (1842) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,353	
A. acinacea, Lindley in Mitch. Three Exped. II, 267 (1838) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,353	
A. lineata, Cunningham in G. Don, gen. syst. II, 403 (1832) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,353	M.fr.I,5;II,177.
A. lachnophylla, F. v. M. in Wing's S. Sc. Record, July (1882)	W.A.	—	—	—	—	—	—	—	
A. triquetra, Bentham in Hook. Lond. Journ. I, 358 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,354	
A. ligustrina, Meissner in Lehm. pl. Preiss. II, 203 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.II,354	
A. Meissneri, Lehmann, delect. sem. hort. bot. Hamb. (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,354	
A. anceps, De Candolle, Mém. sur la fam. des Légum. 446 (1825)	W.A.	S.A.	—	—	—	—	—	B.fl.II,355	
A. hispidula, Willdenow, spec. plant. IV, 1054 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,355	
A. undulifolia, Fraser in Loddiges, Bot. Cabinet XVI, 1544 (1829)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,355	
A. flexifolia, Cunningham in Hook. Lond. Journ. I, 359 (1842)...	—	—	—	—	N.S.W.	—	—	B.fl.II,356	
A. dura, Bentham in Schlecht. Linnaea XXVI, 622 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,356	
A. spathulata, F. v. M. in Benth. Fl. Austr. II, 356 (1864) ...	W.A.	—	—	—	—	—	N.A.	B.fl.II,356	
A. microcarpa, F. v. M., second gen. Report 7 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,357	M.fr.I,6.
A. montana, Bentham in Hook. Lond. Journ. I, 360 (1842) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,357	
A. verniciflua, Cunningham in Field's N.S.Wales 344 (1825) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,358	
A. leprosa, Sieber in De Cand. prodr. II, 450 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.II,358	
A. stricta, Willdenow, spec. plant. IV, 1052 (1805)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,358	
A. dodonaeifolia, Willdenow, enum. pl. hort. Berol. suppl. 68 (1813)	—	S.A.	—	—	—	—	—	B.fl.II,359	
A. Gnidium, Bentham, Fl. Austr. II, 359 (1864)	—	—	—	—	—	Q.	—	B.fl.II,359	
A. ramosissima, Bentham in Hook. Lond. Journ. I, 356 (1842)...	W.A.	—	—	—	—	—	—	B.fl.II,360	
A. Dempsteri, F. v. M., fragm. XI, 65 (1879)	W.A.	—	—	—	—	—	—	—	M.fr.XI,65.
A. Sentis, F. v. M., second gen. Report 11 (1854)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,360	M.fr.XI,65.
A. dentifera, Bentham in Maund, Botanist IV, t. 179 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.II,361	
A. sclerosperma, F. v. M. in Wing's South. Sc. Rec., July (1882)	W.A.	—	—	—	—	—	N.A.	—	
A. fasciculifera, F. v. M. in Benth. Fl. Austr. II, 361 (1864) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,361	
A. falcata, Willdenow, spec. plant. IV, 1053 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,361	
A. macradenia, Bentham in Mitch. Trop. Austr. 360 (1848) ...	—	—	—	—	—	Q.	—	B.fl.II,362	
A. penninervis, Sieber in De Cand. prodr. II, 452 (1825) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,362	
A. amblyphylla, F. v. M. in Wing's S. Sc. Record, July (1882)	W.A.	—	—	—	—	—	—	—	
A. retinodes, Schlechtendal, Linnaea XX, 664 (1847)	—	S.A.	—	V.	—	—	—	B.fl.II,362	
A. neriifolia, Cunningham in Hook. Lond. Journ. I, 357 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,363	
A. microbotrya, Bentham in Hook. Lond. Journ. I, 353 (1842)...	W.A.	—	—	—	—	—	—	B.fl.II,363	
A. leiophylla, Bentham in Hook. Lond. Journ. I, 351 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,364	
A. cyanophylla, Lindley, Bot. Regist. Misc. 49 (1835)	W.A.	—	—	—	—	—	—	B.fl.II,364	
A. pycnantha, Bentham in Hook. Lond. Journ. I, 351 (1842) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,365	

A. notabilis, F. v. M., fragm. I, 6 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.fl.II,365	M.fr.I,6.
A. gladiiformis, Cunningham in Hook. Lond. Journ. I, 354 (1842)	—	—	—	—	N.S.W.	—	—	B.fl.II,365	
A. obtusata, Sieber in De Cand. prodr. II, 453 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,366	
A. rubida, Cunningham in Field's N.S.Wales, 344 (1825) ...	—	—	—	—	N.S.W.	—	—	B.fl.II,366	
A. amoena, H. L. Wendland, comm. de Acac. aphyll. 10, t. 4 (1820)	—	—	—	V.	N.S.W.	—	—	B.fl.II,366	
A. hakeoides, Cunningham in Hook. Lond. Journ. I, 354 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,367	
A. salicina, Lindley in Mitch. Three Exped. II, 20 (1838) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,367	
A. rostellifera, Bentham in Hook. Lond. Journ. I, 356 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,368	
A. pycnophylla, Bentham, Fl. Austr. II, 368 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,368	
A. Harveyi, Bentham, Fl. Austr. II, 368 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,368	
A. iteaphylla, F. v. M. in Schlecht. Linnaea XXVI, 617 (1853)	—	—	—	—	—	—	—	—	
A. suaveolens, Willdenow, spec. plant. IV, 1050 (1805)... ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,369	
A. subcoerulea, Lindley, Bot. Regist. t. 1075 (1827)	W.A.	—	—	—	—	—	—	B.fl.II,369	
A. Lindleyi, Meissner in Lehm. pl. Preiss. I, 14 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,370	
A. leptopetala, Bentham in Schlecht. Linnaea XXVI, 619 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,370	
A. Dietrichiana, F. v. M. in Melb. Chemist, July (1882) ...	—	—	—	—	—	—	—	—	
A. Murrayana, F. v. M. in Benth. Fl. Austr. II, 370 (1864) ...	—	S.A.	—	—	—	Q.	—	B.fl.II,370	
A. subulata, Bonpland, pl. rar. cult. à Malm. 110 t. 45 (1813)...	—	—	—	—	N.S.W.	—	—	B.fl.II,370	
A. linifolia, Willdenow, spec. plant. IV, 1051 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,371	
A. Leichhardtii, Bentham, Fl. Austr. II, 372 (1864)	—	—	—	—	—	Q.	—	B.fl.II,372	
A. crassiuscula, H. L. Wendland, comm. de Acac. 31, t. 8 (1820)	—	—	T.	—	N.S.W	Q.	—	B.fl.II,372	
A. buxifolia, Cunningham in Field's N.S.Wales, 344 (1825) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,372	
A. lunata, Sieber in De Cand. prodr. II, 452 (1825)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,373	
A. brachybotrya, Bentham in Hook. Lond. Journ. I, 347 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,373	
A. Wattsiana, F. v. M. in Benth. Fl. Austr. II, 374 (1864) ...	—	S.A.	—	—	—	—	—	B.fl.II,374	
A. podalyrifolia, Cunningham in Loudon, hort. Brit. 707 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,374	
A. uncifera, Bentham in Mitch. Trop. Austr. 341 (1848) ...	—	—	—	—	—	Q.	—	B.fl.II,374	
A. vestita, Ker in Bot. Regist. t. 698 (1823)	—	—	—	—	N.S.W.	—	—	B.fl.II,375	
A. cultriformis, Cunningham in G. Don. gen. syst. II, 406 (1832)	—	—	—	—	N.S.W.	—	—	B.fl.II,375	M.fr.XI,123.
A. pravissima, F. v. M., first gen. report 12 (1853)	—	—	—	V.	N.S.W.	—	—	B.fl.II,375	M.fr.I,5.
A. strongylophylla, F. v. M., fragm. VIII, 226 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,226
A. pyrifolia, De Candolle, Mém. sur la fam. des Légum. 447 (1825)	W.A.	—	—	—	—	—	N.A.	B.fl.II,376	M.fr.III,17.
A. myrtifolia, Willdenow, spec. plant. IV, 1054 (1805)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,376	
VII. Plurinerves.									
A. scalpelliformis, Meissner in Lehm. pl. Preiss. II, 200 (1847)...	W.A.	—	—	—	—	—	—	B.fl.II,377	
A. urophylla, Bentham in Bot. Reg. Misc. 24 (1841)	W.A.	—	—	—	—	—	—	B.fl.II,377	
A. Luehmanni, F. v. M., fragm. XI, 116 (1881)	—	—	—	—	—	—	N.A.	—	M.fr.XI,116.
A. sublanata, Bentham in Hueg. enum. 42 (1837)...	—	S.A.	—	—	—	—	—	B.fl.II,378	M.fr.XI,117.
A. pravifolia, F. v. M., fragm. I, 4 (1858)	—	S.A.	—	—	N.S.W.	Q.	—	—	M.fr.I,4.
A. amblygona, Cunningham in Hook. Lond. Journ. I, 332 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,378	M.fr.IV,3.
A. adnata, F. v. M. in Melb. Chemist, July (1882)	W.A.	—	—	—	—	—	—	—	
A. deltoidea, Cunningham in G. Don. gen. syst. II, 401 (1832)...	—	—	—	—	—	—	N.A.	B.fl.II,378	
A. stipulosa, F. v. M. in Journ. Linn. Soc. III, 119 (1858) ...	W.A.	—	—	—	—	—	N.A.	B.fl.II,379	M.fr.XI,117.
A. loxophylla, Bentham in Schlecht. Linnaea XXVI, 625 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,379	
A. setulifera, Bentham in Schlecht. Linnaea XXVI, 625 (1853)	—	—	—	—	—	—	N.A.	B.fl.II,379	
A. translucens, Cunningham in Hook. icon. plant. t. 160 (1837)	—	—	—	—	—	—	N.A.	B.fl.II,379	
A. impressa, F. v. M. in Journ. Linn. Soc. III, 133 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,380	
A. bivenosa, De Candolle, prodr. II, 452 (1825)	W.A.	—	—	—	—	—	N.A.	B.fl.II,380	
A. trineura, F. v. M., pl. of Vict. II, 25 (1860)	—	S.A.	—	V.	—	—	N.A.	B.fl.II,381	M.fr.IV,5.
A. nitidula, Bentham, Fl. Austr. II, 381 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,381	
A. estrophiolata, F. v. M. in Wing's South. Sc. Rec., July (1882)	—	S.A.	—	—	—	—	—	—	
A. heteroclita, Meissner in Lehm. pl. Preiss. I, 318 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.II,381	M.fr.IV,6.
A. elongata, Sieber in De Cand. prodr. II, 451 (1825)	—	—	—	V.	N.S.W.	Q	—	B.fl.II,381	
A. subporosa, F. v. M., pl. of Vict. II, 24 (1860)...	—	—	—	V.	N.S.W.	—	—	B.fl.II,382	M.fr.IV,5.
A. Simsii, Cunningham in Hook. Lond. Journ. I, 368 (1842) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,382	
A. dissoneura, F. v. M. in Wing's S. Sc. Record, July (1882) ...	—	—	—	—	—	—	—	—	
A. leptospermoides, Bentham in Schlecht. Linnaea XXVI, 626 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,383	
A. homalophylla, Cunningham in Hook. Lond. Journ. I, 365 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,383	
A. pendula, Cunningham in Loudon, hort. Brit. 490 (1830) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,383	
A. Osswaldi, F. v. M., pl. of Vict. II, 27 (1860)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,384	M.fr.IV,5.
A. lineolata, Bentham in Schlecht. Linnaea XXVI, 626 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,384	
A. quadrimarginea, F. v. M., fragm. X, 31 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,31.
A. coriacea, De Candolle, Mém. sur la fam. des Légum. 446 (1825)	W.A.	—	—	—	—	—	N.A.	B.fl.II,385	
A. stenophylla, Cunningham in Hook. Lond. Journ. I, 366 (1842)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,385	
A. hemignosta, F. v. M. in Journ. Linn. Soc. III, 134 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,385	M.fr.XI,67
A. sclerophylla, Lindley in Mitch. Three Exped. II, 138 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,386	
A. farinosa, Lindley in Mitch. Three. Exped. II, 145 (1838) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,386	
A. Whanii, F. v. M. in Benth. Fl. Austr. II, 386 (1864) ...	—	—	—	V.	—	—	—	B.fl.II,386	
A. heteroneura, Bentham in Schlecht. Linnaea XXVI, 624 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,387	
A. viscidula, Cunningham in Hook. Lond. Journ. I, 366 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,387	
A. ixiophylla, Bentham in Hook. Lond. Journ. I, 364 (1842) ...	W.A.	—	—	—	N.S.W.	Q.	—	B.fl.II,387	M.fr.IV,6.
A. dictyophleba, F. v. M., fragm. III, 128 (1863)...	—	S.A.	—	—	—	Q.	N.A.	B.fl.II,388	M.fr.III,128.
A. venulosa, Bentham in Hook. Lond. Journ. I, 366 (1842) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,388	
A. cyclops, Cunningham in Loudon, hort. Brit. 407 (1830) ...	W.A.	S.A.	—	—	—	—	—	B.fl.II,388	
A. melanoxylon, R. Brown in Ait. hort. Kew. sec. ed. V, 462 (1813)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,388	
A. oraria, F. v. M., fragm. XI, 66 (1879)	—	—	—	—	—	Q.	—	—	M.fr.XI,66.
A. implexa, Bentham in Hook. Lond. Journ. I, 368 (1842) ...	—	—	—	V.	N.S.W.	Q	—	B.fl.II,389	
A. harpophylla, F. v. M. in Benth. Fl. Austr. II, 389 (1864) ...	—	—	—	—	—	Q.	—	B.fl.II,389	
A. excelsa, Bentham in Mitch. Trop. Austr. 225 (1848)	—	—	—	—	—	Q.	—	B.fl.II,390	M.fr.IV,6.
A. homaloclada, F. v. M., fragm. XI, 34 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,34.

A. complanata, Cunningham in Hook. Lond. Journ. I, 369 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,390	M.fr.XI,34.
A. dineura, F. v. M. in Journ. Linn. Soc. III, 130 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,391	M.fr.XI,65.
A. binervata, De Candolle, prodr. II, 452 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,390	M.fr.XI,66.
A. latescens, Bentham in Hook. Lond. Journ. I, 380 (1842) ...	—	—	—	—	—	—	N.A.	B.fl.II,391	
A. platycarpa, F. v. M. in Journ. Linn. Soc. III, 145 (1858) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,391	M.fr.XI,67.
A. flavescens, Cunningham in Hook. Lond. Journ. I, 381 (1842)	—	—	—	—	—	Q.	—	B.fl.II,391	M.fr.XI,68.
A. retivenea, F. v. M., fragm. III, 128 (1863)	—	S.A.	—	—	—	—	N.A.	B.fl.II,392	M.fr.III,128.
A. triptera, Bentham in Hook. Lond. Journ. I, 325 (1842) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,323	
A. rhigiophylla, F. v. M. in Schlecht. Linnaea XXVI, 611 (1853)	—	S.A.	—	—	—	—	—	B.fl.II,333	
A. Oxycedrus, Sieber in De Cand. prodr. II, 453 (1825)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,334	
A. verticillata, Willdenow, spec. plant. IV, 1049 (1805)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,334	
A. Riceana, Henslow in Maund, Botanist III, t. 135 (1839) ...	—	—	T.	—	—	—	—	B.fl.II,335	
VIII. Juliferae.									
A. amentifera, F. v. M. in Journ. Linn. Soc. III, 141 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,392	
A. Wickhami, Bentham in Hook. Lond. Journ. I, 379 (1842) ...	—	—	—	—	—	—	N.A.	B.fl.II,392	
A. lysiphloia, F. v. M. in Journ. Linn. Soc. III, 137 (1858) ...	—	S.A.	—	—	—	—	N.A.	B.fl.II,393	
A. linarioides, Bentham in Hook. Lond. Journ. I, 371 (1842) ...	—	—	—	—	—	—	N.A.	B.fl.II,393	
A. stipuligera, F. v. M. in Journ. Linn. Soc. III, 144 (1858) ...	—	—	—	—	—	—	N.A	B.fl.II,393	
A. ptychophylla, F. v. M. in Journ. Linn. Soc. III, 142 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,394	
A. stigmatophylla, Cunningham in Hook. Lond. Journ. I, 377 (1842)	—	—	—	—	—	—	N.A.	B.fl.II,394	
A. umbellata, Cunningham in Hook. Lond. Journ. I, 378 (1842)	—	—	—	—	—	—	N.A.	B.fl.II,394	
A. leptophleba, F. v. M. in Journ. Linn. Soc. III, 143 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,395	
A. limbata, F. v. M. in Journ. Linn. Soc. III, 145 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,395	
A. brevifolia, Bentham, Fl. Austr. II, 395 (1864)	—	—	—	—	—	Q.	—	B.fl.II,395	
A. megalantha, F. v. M. in Journ. Linn. Soc. III, 143 (1858) ..	—	—	—	—	—	—	N.A.	B.fl.II,395	
A. gonoclada, F. v. M. in Journ. Linn. Soc. III, 140 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,396	
A. pycnostachya, F. v. M., pl. of Vict. II, 33 (1860)	—	—	—	—	N.S.W.	—	—	B.fl.II,396	
A. subtilinervis, F. v. M., pl. of Vict. II, 32 (1860)	—	—	—	V.	N.S.W.	—	—	B.fl.II,396	M.fr.IV,8.
A. cochliocarpa, Meissner in Bot. Zeitung 10 (1855)	W.A.	—	—	—	—	—	—	B.fl.II,397	
A. Dallachiana, F. v. M., fragm. I, 7 (1858)	—	—	—	V.	—	—	—	B.fl.II,397	M.fr.I,7.
A. alpina, F. v. M., fragm. III, 129 (1863)	—	—	—	V.	—	—	—	B.fl.II,397	M.fr.III,129.
A. longifolia, Willdenow, spec. pl. IV, 1052 (1805)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,397	
A. linearis, Sims, Bot. Mag. t. 2156 (1820)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,399	
A. aciphylla, Bentham in Schlecht. Linnaea, XXVI, 627 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,399	
A. ephedroides, Bentham in Hook. Lond. Journ. I, 370 (1842) ...	W.A.	—	—	—	—	—	—	B.fl.II,399	
A. Burkittii, F. v. M. in Benth. Fl. Austr. II, 400 (1864) ...	—	S.A.	—	—	—	—	—	B.fl.II,400	
A. microneura, Meissner in Lehm. pl. Preiss. I, 19 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,400	
A. cyperophylla, F. v. M. in Benth. Fl. Austr. II, 400 (1864) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,400	
A. multispicata, Bentham, Fl. Austr. II, 400 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,400	
A. pityoides, F. v. M. in Journ. Linn. Soc. III, 135 (1858) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,400	
A. xylocarpa, Cunningham in Hook. Lond. Journ. I, 370 (1842)	—	—	—	—	—	—	N.A.	B.fl.II,401	
A. gonocarpa, F. v. M. in Journ. Linn. Soc. III, 136 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,401	
A. oncinophylla, Lindley, Bot. Regist. XXV, App. XV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.II,401	
A. conjunctifolia, F. v. M., fragm. XI, 68 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,68.
A. drepanocarpa, F. v. M. in Journ. Linn. Soc. III, 137 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,402	
A. arida, Bentham in Hook. Lond. Journ. I, 370 (1842)	—	—	—	—	—	—	N.A.	B.fl.II,402	
A. aneura, F. v. M. in Schlecht. Linnaea, XXVI, 627 (1853) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,402	M.fr.IV,8.
A. cibaria, F. v. M. in Melb. Chemist, July (1882)	W.A.	—	—	—	—	—	—	—	
A. Kempeana, F. v. M. in Melb. Chemist, July (1882)	—	S.A.	—	—	N.S.W.	Q.	—	—	
A. conspersa, F. v. M. in Journ. Linn. Soc. III, 140 (1858) ..	—	—	—	—	—	—	N.A.	B.fl.II,403	
A. Doratoxylon, Cunningham in Field's N.S. Wales, 345 (1825)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,403	
A. acuminata, Bentham in Hook. Lond. Journ. I, 373 (1862) ...	W.A.	—	—	—	—	—	—	B.fl.II,404	M.fr.IV,7.
A. stereophylla, Meissner in Lehm. pl. Preiss. II, 203 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.II,404	
A. signata, F. v. M., fragm. IV, 7 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,404	M.fr.IV,7.
A. delibrata, Cunningham in Hook. Lond. Journ. I, 374 (1842)	—	—	—	—	—	—	N.A.	B.fl.II,404	
A. oligoneura, F. v. M. in Journ. Linn. Soc. III, 139 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,405	
A. torulosa, Bentham in Journ. Linn. Soc. III, 139 (1858) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,405	
A. julifera, Bentham in Hook. Lond. Journ. I, 374 (1842) ...	—	—	—	—	—	Q.	—	B.fl.II,405	
A. Solandri, Bentham, Fl. Austr. II, 406 (1864)	—	—	—	—	—	Q.	—	B.fl.II,406	
A. leptostachya, Bentham, Fl. Austr. II, 406 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,406	
A. glaucescens, Willdenow, spec. plant. IV, 1052 (1805) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,406	
A. Cunninghamii, Hooker, icon. plant. t. 165 (1837)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,407	
A. leptocarpa, Cunningham in Hook. Lond. Journ. I, 376 (1842)	—	—	—	—	—	Q.	—	B.fl.II,407	
A. polystachya, Cunningham in Hook. Lond. Journ. I, 376 (1842)	—	—	—	—	—	Q.	N.A.	B.fl.II,407	
A. holcocarpa, Bentham, Fl. Austr. II, 408 (1864)	—	—	—	—	—	Q.	—	B.fl.II,408	M.fr.XI,69.
A. plectocarpa, Cunningham in Hook. Lond. Journ. I, 375 (1842)	—	—	—	—	—	—	N.A.	B.fl.II,408	
A. pachycarpa, F. v. M. in Journ. Linn. Soc. III, 139 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,408	
A. tumida, F. v. M. in Journ. Linn. Soc. III, 144 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.II,409	
A. loxocarpa, Bentham in Hook. Lond. Journ. I, 377 (1842) ...	—	—	—	—	—	—	N.A.	B.fl.II,409	
A. oncinocarpa, Bentham in Hook. Lond. Journ. I, 378 (1842) ...	—	—			—	—	N.A.	B.fl.II,409	
A. retinervis, Bentham in Hook. Lond. Journ. I, 379 (1842) ...	—	—	—	—	—	—	N.A.	B.fl.II,410	
A. aulacocarpa, Cunningham in Hook. Lond. Journ. I, 378 (1842)	—	—	—	—	—	Q.	—	B.fl.II,410	
A. calyculata, Cunningham in Hook. Lond. Journ. I, 379 (1842)	—	—				Q.	—	B.fl.II,410	
A. crassocarpa, Cunningham in Hook. Lond. Journ. I, 379 (1842)	—	—				Q.	N.A.	B.fl.II,410	M.fr.XI,69.
A. auriculiformis, Cunningham in Hook. Lond. Journ. I, 377 (1842)	—	—	—			Q.	N.A.	B.fl.II,411	
A. latifolia, Bentham in Hook. Lond. Journ. I, 382 (1842) ...	—	—	—	—		—	N.A.	B.fl.II,411	
A. Mangium, Willdenow, spec. plant. IV, 1053 (1805)	—	—	—	—	—	Q.	—	—	M.fr.XI,36.
A. holosericea, Cunningham in G. Don, gen. syst. II, 407 (1832)	—	—	—	—	—	Q.	N.A.	B.fl.II,411	
A. cincinnata, F. v. M., fragm. XI, 35 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,35.
A. dimidiata, Bentham in Hook. Lond. Journ. I, 381 (1842) ...	—	—	—	—	—	—	N.A.	B.fl.II,412	

A. humifusa, A. Cunningham in Lond. Journ. I, 382 (1842) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,412	
A. denticulosa, F. v. M., fragm. X, 32 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,32
IX. Bipinnatae.									
A. Drummondii, Lindley, Bot. Regist. XXV, App. XV (1839)...	W.A.	—	—	—	—	—	—	B.fl.II,419	
A. pulchella, R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813)	W.A.	—	—	—	—	—	—	B.fl.II,416	
A. Mitchelli, Bentham in Hook. Lond. Journ. I, 387 (1842) ...	—	S.A.	—	V.	—	—	—	B.fl.II,417	
A. pentadenia, Lindley, Bot. Regist. t. 1521 (1832)	W.A.	—	—	—	—	—	—	B.fl.II,417	
A. Gilberti, Meissner in Lehm. pl. Preiss. II, 204 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.II,417	
A. nigricans, R. Brown in Ait. hort. Kew. sec. ed. V, 465 (1813)	W.A.	—	—	—	—	—	—	B.fl.II,418	
A. Tayloriana, F. v. M. in Wing's S. Sc. Record, July (1882) ...	W.A.	—	—	—	—	—	—	—	
A. strigosa, Link, enum. pl. hort. Berol. 444 (1822)	W.A.	—	—	—	—	—	—	B.fl.II,418	
A. elata, Cunningham in Hook. Lond. Journ. I, 383 (1842) ...	—	—	—	—	N.S.W.	—	—	B.fl.II,413	
A. pruinosa, Cunningham in Hook. Lond. Journ. I, 383 (1842)...	—	—	—	—	N.S.W.	—	—	B.fl.II,413	
A. spectabilis, Cunningham in Hook. Lond. Journ. I, 383 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,413	
A. polybotrya, Bentham in Hook. Lond. Journ. I, 384 (1842) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,414	
A. discolor, Willdenow, spec. plant. IV, 1068 (1805)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,414	
A. decurrens, Willdenow, spec. plant. IV, 1872 (1805)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,414	
A. dealbata, Link, enum. pl. hort. Berol. 445 (1822)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,415	
A. cardiophylla, Cunningham in Hook. Lond. Journ. I, 385 (1842)	—	—	—	—	N.S.W.	—	—	B.fl.II,415	
A. leptoclada, Cunningham in Hook. Lond. Journ. I, 385 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,416	
A. pubescens, R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813)	—	—	—	—	N.S.W.	—	—	B.fl.II,416	
A. Farnesiana, Willdenow, spec. plant. IV, 1083 (1805)... ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,419	
A. suberosa, Cunningham in Hook. Lond. Journ. I, 499 (1842)...	—	—	—	—	—	—	N.A.	B.fl.II,420	M fr.XI,36.
A. Bidwilli, Bentham in Schlecht. Linnaea XXVI, 629 (1853) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,420	
A. pallida, F. v. M. in Journ. Linn. Soc. III, 14 (1858)	—	—	—	—	—	—	N.A.	B.fl.II,421	
ALBIZZIA, Durazzini in Magazzino Toscano III, part IV, 13 (1772). (Zygia 1756, Pithecolobium, Cathormion, Calliandra Enterolobium, Serianthes.)									
A. lophantha, Bentham in Hook. Lond. Journ. III, 86 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.II,421	
A. Sutherlandi, F. v. M., fragm. VI, 22 (1867)	—	—	—	—	—	—	N.A.	—	M.fr.IX,179.
A. basaltica, Bentham, Fl. Austr. II, 422 (1864)	—	—	—	—	—	Q.	—	B.fl.II,422	
A. Thozetiana, F. v. M. in Benth. Fl. Austr. II, 422 (1864) ...	—	—	—	—	—	Q.	—	B.fl.II,422	M.fr.IV,9&176
A. amoenissima, F. v. M., fragm. VIII, 165 (1874)	—	—	—	V.	N.S.W.	Q.	—	—	M.fr.IX,179.
A. procera, Bentham in Hook. Lond. Journ. III, 88 (1844) ...	—	—	—	—	—	—	N.A.	B.fl.II,422	
A. pruinosa, F. v. M. in Trimen's Journ. of Bot. 9 (1872) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,423	M.fr.IX,179.
A. canescens, Bentham, Fl. Austr. II, 423 (1864)...	—	—	—	—	—	Q.	—	B.fl.II,423	M.fr.IX,179.
A. Tozeri, F. v. M. in Trimen's Journ. of Bot. 10 (1872) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,424	M.fr.IX,179.
A. Hendersoni, F. v. M. in Trimen's Journ. of Bot. 10 (1872) ...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.X,178.
A. monilifera, F. v. M. in Trimen's Journ. of Bot. 10 (1872) ...	—	—	—	—	—	—	N.A.	B.fl.II,424	
A. ramiflora, F. v. M., fragm. IX, 178 (1875)	—	—	—	-	—	Q.	—	—	M.fr.V,10;IX,178.
ARCHIDENDRON, F. v. M., fragm. V, 59 (1865).									
A. Vaillantii, F. v. M., fragm. V, 60 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,9;IX,178.
A. Lucyi, F. v. M., fragm. VI, 201 (1868)...	—	—	—	—	—	Q.	—	—	M.fr.IX,178.

ROSACEAE.

A. L. de Jussieu, gen. pl. 334 (1789), from B. de Jussieu (1759).

PARINARIUM, A. L. de Jussieu, gen. pl. 342 (1789).									
P. Nonda, F. v. M. in Benth. Fl. Austr. II, 426 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,426	
P. Griffithianum, Bentham in Hook. Niger-Flora 334 (1849) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,426	M.fr.VIII,34
STYLOBASIUM, Desfontaines in Mém. du Mus. V, 37, t. 2 (1819). (Macrostigma.)									
S. spathulatum, Desfontaines in Mém. du Mus. V, 37, t. 2 (1819)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.II,427	
S. lineare, Nees in Lehm. pl. Preiss. I, 95 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,427	
GEUM, Linné, gen. plant. 148 (1737) from Plinius. (Sieversia.)									
G. urbanum, Linné, spec. plant. 501 (1753)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,428	
G. renifolium, F. v. M. in Transact. phil. Inst. Vict. II, 66 (1857)	—	—	T.	—	—	—	—	B.fl.II,428	
POTENTILLA, Linné, gen. plant. 147 (1737) from Camerarius (1586).									
P. anserina, Linné, spec. plant. 495 (1753)	—	S.A.	T.	V.	—	—	—	B.fl.II,429	
RUBUS, Tournefort, Inst. 614, t. 385 (1700) from Plinius.									
R. Gunnianus, Hooker, icon. plant. t. 291 (1840)	—	—	T.	—	—	—	—	B.fl.II,430	
R. parvifolius, Linné, spec. plant. 1197 (1753)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,430	M.fr.IV,30.
R. rosifolius, Smith, plantar. icon. ex herb. Linn. t. 60 (1790) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,431	M.fr.IV,32;VIII,34.
R. Moluccanus, Linné, spec. plant. 1197 (1753)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.II,430	M.fr.IV,31;VIII,34.
R. Moorei, F. v. M. in Transact. phil. Inst. Vict. II, 67 (1857)...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,431	M.fr.IV,29&176;VIII,34.
AGRIMONIA, Tournefort, inst. 301, t. 155 (1700) from C. Bauhin (1623).									
A. Eupatoria, Linné, spec. plant. 448 (1753)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,34.
ALCHEMILLA, Linné, gen. pl. 30 (1737) from C. Bauhin (1623). (Alchimilla.)									
A. vulgaris, C. Bauhin, pinax 319 (1623)	—	—	V.	N.S.W.	—	—	B.fl.II,432		
ACAENA, Mutis in Linné, mantiss. II, 200 (1771). (Ancistrum.)									
A. ovina, Cunningham in Field's N.S. Wales, 358 (1825) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,433	
A. sanguisorbae, Vahl, enum. plant. I, 294 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,434	
A. montana, J. Hooker in Hook. Lond. Journ. VI, 276 (1847) ...	—	—	T.	—	—	—	—	B.fl.II,434	

SAXIFRAGEAE.

Ventenat, Tabl. III, 277 (1799).

ARGOPHYLLUM, R. et G. Forster, charact. gen. 29, t. 15 (1776).									
A. Lejourdanii, F. v. M., fragm. IV, 33 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,437	M.fr.VI,188.

ABROPHYLLUM, J. Hooker in Benth. fl. Austr. II, 437 (1864). (Brachynema.)
A. ornans, J. Hooker in Benth. pl. Austr. II, 437 (1864) ... — — — — N.S.W. Q. — B.fl.II,437 M.fr.VI,189;VII,150

CUTTSIA, F. v. M., fragm. V, 47 (1865).
C. viburnea, F. v. M., fragm. V, 47 (1865)... ... ... — — — — N.S.W. Q. — — M.fr.V,47&189;VI,189

COLMEIROA, F. v. M., fragm. VII, 149 (1871).
C. carpodetoides, F. v. M., fragm. VII, 149 (1871) ... ... — — — — N.S.W. — — — M.fr.VII,149.

QUINTINIA, A. de Candolle, Mon. Campan. 92 (1830).
Q. Sieberi, A. De Candolle, Monogr. Camp. 92 (1830) ... ... — — — — N.S.W. Q. — B.fl.II,438 M.fr.III,126;VI,189.
Q. Verdonii, F. v. M., fragm. II, 225 (1861) ... ... — — — — N.S.W. Q. — B.fl.II,438 M.fr.VI,189.
Q. Fawkneri, F. v. M., fragm. VI, 92 (1867) ... ... — — — — — Q. — — M.fr.VII,150.

POLYOSMA, Blume, Bijdr. 658 (1825).
P. Cunninghamii, J. J. Bennett, pl. Jav. rar. 196 (1840) ... — — — — N.S.W. — — B.fl.II,438 M.fr.VI,189.
P. alangiacea, F. v. M., fragm. VIII, 8 (1872) ... ... — — — — — Q. — — M.fr.VIII,8.

ANOPTERUS, Labillardière, Nov. Holl. pl. spec. I, 86, t. 112 (1804.)
A. glandulosus, Labillardière, Nov. Holl. pl. spec. I, 86, t. 112 (1804) — — T. — — — — B.fl.II,439
A. Macleayanus, F. v. M. in Journ. Pharm. Soc. Vict. 43 (1858) — — — — N.S.W. Q. — B.fl.II,439 M.fr.VI,188;VII,150.

CALLICOMA, Andrews, Bot. Reposit. t. 566 (1809). (Calycomis partly.)
C. serratifolia, Andrews, Bot. Reposit. t. 566 (1809) ... ... — — — — N.S.W. Q. — B.fl.II,440 M.fr.V,32.
C. Stutzeri, F. v. M., fragm. V, 34 (1865) ... ... — — — — — Q. — — M.fr.VI,188&252;VII,150.

ANODOPETALUM, Cunningham in Endl. gen. 818 (1839).
A. biglandulosum, Cunningham in J. Hook. Fl. Tasm. I, 148 (1860) — — T. — — — — B.fl.II,440 M.fr.VI,189.

APHANOPETALUM, Endlicher, nov. stirp. decad. I, 34 (1839). (Platyptelea.)
A. resinosum, Endlicher, nov. stirp. dec. I, 35 (1839) ... ... — — — V. N.S.W. Q. — B.fl.II,441 M.fr.I,228.
A. occidentale, F. v. M., fragm. I, 228 (1858) ... ... W.A. — — — — — — B.fl.II,441 M.fr.XI,20.

CERATOPETALUM, Smith, specim. of the Bot. of New Holland, 9 (1793).
C. gummiferum, Smith, specim. of the Bot. of New Holl. t. 3 (1793) — — — — N.S.W. — — B.fl.II,442
C. apetalum, D. Don in Edinb. new phil. journ. (1830) ... — — — — N.S.W. — — B.fl.II,442 M.fr.VI,189;VII,150.

SCHIZOMERIA, D. Don in Edinb. new phil. journ. IX, 94 (1830).
S. ovata, D. Don in Edinb. new phil. journ. IX, 94 (1830) ... — — — — N.S.W. Q. — B.fl.II,443 M.fr.VI,189.

ACROPHYLLUM, Bentham in Maund & Henslow, Botanist II, 95 (1840). (Calycomis partly.)
A. venosum, Bentham in Maund & Henslow, Bot. II, 95 (1840) — — — — N.S.W. — — B.fl.II,443 M.fr.VI,189;VII,150.

WEINMANNIA, Linné, syst. ed. X, 1005 (1759). (Ackama.)
W. paniculosa, F. v. M., fragm. II, 175 (1861) ... ... — — — — N.S.W. Q. — B.fl.II,444 M.fr.II,83; VI,188; VII,
W. Biagiana, F. v. M., fragm. V, 16 (1865) ... ... — — — — — Q. — — M fr.VII,150. [150.
W. lachnocarpa, F. v. M., fragm. VIII, 7 (1872)... ... ... — — — — N.S.W. Q. — — M.fr.VIII,7.
W. rubifolia, Bentham, Fl. Aust. II, 445 (1864) ... ... — — — — N.S.W. Q. — B.fl.II,445 M.fr.II,82&175;VI,181.
W. Benthami, F. v. M., Docum. intercol. Exhibit. 31 (1866) ... — — — — — — — B.fl.II,446 M.fr.V,180;VI,188.

GILLBEEA, F. v. M., fragm. V, 17 (1865).
G. adenopetala, F. v. M., fragm. V, 17 (1865) ... ... — — — — — Q. — — M.fr.V,17;VI,188.

DAVIDSONIA, F. v. M., fragm. VI, 4 (1867).
D. pruriens, F. v. M., fragm. VI, 4 (1867)... ... ... — — — — — Q. — — M.fr.VI,249.

TETRACARPAEA, J. Hooker, icon. plant. t. 264 (1840).
T. Tasmanica, J. Hooker, icon. plant. t. 264 (1840) ... ... — — T. — — — — B.fl.II,445

EUCRYPHIA, Cavanilles, icon. IV, 48, t. 372 (1797). (Carpodontos.)
E. Billardieri, Spach, veg. phan. V, 344 (1836) ... ... — — T. — — — — B.fl.II,446 M.fr.VI,189;VII,150.
E. Moorei, F. v. M., fragm. IV, 2 (1864) ... ... — — — V. N.S.W. — — B.fl.II,447 M.fr.IV,176;VI,189;VII [150

BAUERA, Banks & Kennedy in Andrews, Bot. Rep. t. 198 (1793).
B. rubioides, Andrews, Bot. Reposit. t. 198 (1793) ... ... — S.A. T. V. N.S.W. — — B.fl.II,447 M.fr.IV,23.
B. capitata, Seringe in De Cand. prodr. IV, 13 (1830) ... ... — — — — N.S.W. Q. — B.fl.II,448 M.fr.IV,24.
B. sessiliflora, F. v. M. in Transact. phil. Soc. Vict. I, 41 (1854) — — — V. — — — B.fl.II,448 M.fr.IV,24;VII,150

EREMOSYNE, Endlicher in Hueg. enum. pl. Nov. Holl. austr. occ. 53 (1837).
E. pectinata, Endlicher in Hueg. enum. pl.53 (1837) ... ... W.A. — — — — — — B.fl.II,449 M.fr.VI,189.

CEPHALOTUS, Labillardière, Nov. Holl. pl. spec. II, 7, t. 145 (1806).
C. follicularis, Labillardière, Nov. Holl. pl. spec. II, 7, t. 145 (1806) W.A. — — — — — — B.fl.II,449 M.fr.VI,189.

CRASSULACEAE.

De Candolle, Bull. de la soc. philom. n. 49 (1801).

TILLAEA, Micheli, nov. gen. 22, t. 20 (1729). (Bulliarda, Crassulae subgentis).
T. verticillaris, De Candolle, prodr. III, 382 (1828) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.II,451 M.fr.XI,118.
T. purpurata, J. Hooker in Lond. Journ. of Bot. VI, 472 (1847) W.A. S.A. T. V. N.S.W. — — B.fl.II,451 M.fr.XI,118
T. macrantha, J. Hooker in Hook. icon. plant. t. 310 (1841) ... — S.A. T. V. N.S.W. — — B.fl.II,452 M.fr.XI,117.
T. pedicellosa, F. v. M., fragm. XI, 118 (1881) ... ... W.A. — — — — — — — M.fr.XI,118.
T. intricata, Nees in Lehm. pl. Preiss. I, 278 (1844) ... ... W.A. — — — — — — — M.fr.XI,117.
T. recurva, J. Hooker, Fl. Tasm. I, 146 (1860) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.II,452 M.fr.XI,118.

HAMAMELIDEAE.

R. Brown in Abel, Narr. journ. Chin. 374 (1818).

Genus and species undetermined ... ... — — — — — Q. — —

ONAGREAE.

Adanson, Familles des plantes II, 81 (1763), from B. de Jussieu (1759).

OENOTHERA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 112 (1737).

O. Tasmanica, J. Hooker, Fl. Tasm. I, 119 (1860)	—	—	T.	—	—	—	—	B.fl.III,303			

EPILOBIUM, Dillenius in Linné, syst. nat. 8 (1735), from Gesner (1542).

E. tetragonum, Linné, spec. plant. 348 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,305			

JUSSIEUA, Linné, gen. plant. 126 (1737). (Jussiaea.)

J. repens, Linné, spec. plant. 388 (1753)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,306			
J. suffruticosa, Linné, spec. plant. 388 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,307	M.fr.III,13.		

LUDWIGIA, Linné, coroll. 3 (1737).

L. parviflora, Roxburgh, Fl. Ind. edit. Carey. I, 440 (1820)	—	—	—	—	—	Q.	N.A.	B.fl.III,307	M.fr.III,129;IV,175.		

SALICARIEAE.

Adanson, Familles des plantes II, 232 (1763), from B. de Jussieu (1759).

ROTALA, Linné, mantissa altera 143 (1771).

R. verticillaris, Linné, mantissa altera 143 (1771)	—	—	—	—	—	—	N.A.	B.fl.III,295	M.fr.III,108.		
R. Roxburghiana, Wight, Icon. t. 260 (1840)	—	—	—	—	—	Q.	N.A.	B.fl.III,296	M.fr.XI,118.		
R. diandra, F. v. M. in Transact. Roy. Soc. of N.S.Wales 89 (1880)	—	—	—	—	—	—	N.A.	B.fl.III,296			

AMMANNIA, Houston in Linné, gen. plant. 337 (1737).

A. crinipes, F. v. M. in Transact. phil. Soc. Vict. III, 49 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,296			
A. triflora, R. Brown in Benth. Fl. Austr. III, 297 (1866) ...	—	—	—	—	—	—	N.A.	B.fl.III,297			
A. baccifera, Linné, spec. plant. ed. sec. 175 (1762)	—	S.A.	—	—	—	Q.	N.A.	B.fl.III,297			
A. auriculata, Willdenow, hort. Berol. t. 7 (1806)	—	—	—	—	—	Q.	—	B.fl.III,297			
A. multiflora, Roxburgh, Fl. Ind. edit. Carey. I, 447 (1820) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,298			

LAGERSTROEMIA, Linné, syst. ed. X, 1076 (1759).

L. Indica, Linné, spec. plant. 734 (1753)	—	—	—	—	—	Q.	—	—	M.fr.VIII,35.		
L. Flos Reginae, Retzius, observ. botan. V, 25 (1789)	—	—	—	—	—	Q.	—	—	M.fr.XII.		

LYTHRUM, Linné, syst. nat. 8 (1735); Linné, gen. plant. 136 (1737).

L. Salicaria, Linné, spec. plant. 446 (1753)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,298	M.fr.VII,147.		
L. Hyssopifolia, Linné, spec. plant. 447 (1753)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,299	M.fr.VII,147.		

NESAEA, Commerçon in A. L. de Jussieu, gen. plant. 332 (1789).

N. Robertsii, F. v. M., fragm. VII, 145 (1871)	—	—	—	—	—	Q.	—	—	M.fr.VII,145.		
N. Arnhemica, F. v. M., fragm. VII, 146 (1871)	—	—	—	—	—	—	N.A.	B.fl.III,299	M.fr.III,109;VII,146.		

PEMPHIS, R. et G. Forster, charact. gen. 67, t. 34 (1776).

P. acidula, R. & G. Forster, charact. gen. 67, t. 34 (1776) ...	—	—	—	—	—	Q.	N.A.	B.fl.III,300	M.fr.VII,147.		

LAWSONIA, Linné, gen. plant. 111 (1737).

L. alba, Lamarck, Encycl. méth. III, 106 (1789)	—	—	—	—	—	—	N.A.	B.fl.III,301			

HALORAGEAE.

R. Brown in Flind. voy. II, 549 (1814).

LOUDONIA, Lindley, Bot. Regist. XXV, App. XLII (1839). (Glischrocaryon.)

L. aurea, Lindley, Bot. Regist. XXV, App. XLII (1839) ...	W.A.	S.A.	—	—	—	—	—	B.fl.II,472	M.fr.VIII,162;XI,20.		
L. Behrii, Schlechtendal, Linnaea XX, 648 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,472	M.fr.VIII,162.		
L. Roei, Schlechtendal, Linnaea XX, 648 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,472	M.fr.VIII,162.		

HALORAGIS, R. et G. Forster, charact. gen. 61, t. 31 (1776). (Cercodia, Goniocarpus.)

H. Gossei, F. v. M., fragm. VIII, 161 (1874)	—	S.A.	—	—	—	—	N.A.	—	M.fr.XI,134.		
H. trigonocarpa, F. v. M., fragm. X, 84 (1876)	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,84.		
H. digyna, Labillardière, Nov. Holl. pl. spec. I, 101, t. 129 (1804)	W.A.	—	—	—	—	—	—	B.fl.II,475			
H. mucronata, Bentham, Fl. Austr. II, 475 (1864)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,475			
H. pithyoides, Bentham, Fl. Austr. II, 476 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,476			
H. cordigera, Fenzl in Hueg. enum. 45 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,476			
H. elata, Cunningham in Hueg. enum. 45 (1837)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,476	M.fr.VIII,162.		
H. tenuifolia, Bentham, Fl. Austr. II, 477 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,477			
H. scoparia, Fenzl in Hueg. enum. 45 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,477	M.fr.X,54.		
H. aculeolata, Bentham, Fl. Austr. II, 477 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,477	M.fr.X,54.		
H. foliosa, Bentham, Fl. Austr. II, 477 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,477			
H. confertifolia, F. v. M., fragm. X, 53 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,54.		
H. platycarpa, Bentham, Fl. Austr. II, 478 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,478			
H. ceratophylla, Zahlbruckner in Endl. Atakta, 16, t. 15 (1833)	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.II,478			
H. acutangula, F. v. M. in Transact. Vict. Inst. 125 (1854) ...	—	S.A.	—	—	—	—	—	B.fl.II,478			
H. hexandra, F. v. M., fragm. III, 31 (1862)	W.A.	—	—	—	—	—	—	B.fl.II,478	M.fr.III,31.		
H. odontocarpa, F. v. M., fragm. I, 108 (1859)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,479	M.fr.I,108;X,54.		
H. serra, Brongniart in Duperr. Voy. sur la Coquill. t. 69 (1829)	—	—	—	—	N.S.W.	—	—	B.fl.II,479			
H. glauca, Lindley in Mitch. Trop. Austr. 91 (1848)	—	—	—	—	N.S.W.	—	—	B.fl.II,479			
H. alata, N. J. Jacquin, icon. plant. rar. I, 7, t. 69 (1786) ...	—	—	—	V.	N.S.W.	—	—	B.fl.II,479			
H. racemosa, Labillardière, Nov. Holl. pl. spec. I, 100, t. 128 (1804)	W.A.	—	—	—	—	—	—	B.fl.II,480	M.fr.VIII,162.		
H. rotundifolia, Bentham, Fl. Austr. II, 480 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,480	M.fr.VIII,162.		
H. rudis, Bentham, Fl. Austr. II, 480 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,480			
H. paniculata, R. Brown in Benth. Fl. Austr. II, 481 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.II,481			
H. pusilla, R. Brown in Benth. Fl. Austr. II, 481 (1864) ..	W.A.	—	—	—	—	—	—	B.fl.II,481			
H. intricata, Bentham, Fl. Austr. II, 481 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,481			
H. nodulosa, Walpers, repert. V, 672 (1846)	W.A.	—	—	—	—	—	—	B.fl.II,481			
H. trichostachya, Bentham, Fl. Austr. II, 481 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,481			
H. lanceolata, R. Brown in Benth. Fl. Austr. II, 481 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.II,482			
H. micrantha, R. Brown in Flind. Voy. II, 550 (1814)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,482	M.fr.VIII,162;X,54.		

H. stricta, R. Brown in Benth. Fl. Austr. II, 482 (1864) ...	—	—	—	—	—	Q.	—	B.fl.II,482	
H. heterophylla, Brongniart in Duperr. Voy. t. 68 (1829) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,483	
H. pinnatifida, A. Gray, Bot. Wilk. Expl. Exped. I, 627 (1854)	—	—	—	—	N.S.W.	—	—	B.fl.II,483	
H. leptothoca, F. v. M., fragm. III, 32 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,483	M.fr.VIII,162.
H. tetragyna, R. Brown in Flind. Voy. II, 550 (1814)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,484	M.fr.IV,26.
H. teucrioides, A. Gray, Bot. Wilk. Expl. Exped. I, 625 (1854)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,484	
H. scordioides, Bentham, Fl. Austr. II, 485 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,485	
H. depressa, Walpers, repertor. II, 99 (1843)	—	—	T.	V.	—	—	—	B.fl.II,485	
H. salsoloides, Bentham, Fl. Austr. II, 485 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,485	
MEIONECTES, R. Brown in Flind. Voy. II, 550 (1814).									
M. Brownii, J. Hooker in Hook. icon. plant. t. 306 (1841) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,486	M.fr.VIII,163.
MYRIOPHYLLUM, l'Ecluse, rar. stirp. hist. II, 252 (1583) from Dioscorides. (Pelonastes).									
M. variifolium, J. Hooker in Hook. icon. plant. t. 289 (1840) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,487	
M. elatinoides, Gaudichaud in Ann. des sc. nat. V, 105 (1825) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,487	
M. verrucosum, Lindley in Mitch. Trop. Austr. 384 (1848) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,488	M.fr.VIII,163.
M. latifolium, F. v. M., fragm. II, 87, (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,488	M.fr.II,87;IV,172;X,54.
M. Muelleri, Sonder in Linnaea, XXVIII, 233 (1855)	W.A.	S.A.	—	V.	—	—	—	B.fl.II,488	M.fr.VIII,162.
M. amphibium, Labillardière, Nov. Holl. pl. spec. II, 70, t. 220 (1806)	—	S.A.	T.	V.	—	—	—	B.fl.II,489	M.fr.VIII,163.
M. pedunculatum, J. Hooker in Lond. Journ. VI, 474 (1847) ...	W.A.	S.A.	T.	V.	—	—	—	B.fl.II,489	M.fr.VIII,162.
M. dicoccum, F. v. M. in Transact. phil. Inst. Vict. III, 41 (1858)	—	—	—	—	—	—	N.A.	B.fl.II,489	M.fr.VIII,162.
M. trachycarpum, F. v. M., fragm. II, 87 (1860)	—	—	—	—	—	—	N.A.	B.fl.II,489	M.fr.II,87.
M. gracile, Bentham, Fl. Austr. II, 489 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,489	
M. filiforme, Bentham, Fl. Austr. II, 489 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,489	
M. integrifolium, J. Hooker, Fl. Tasm. I. 123, t. 23 (1860) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,490	M.fr.VIII,162.
M. Drummondii, J. Hooker, Fl. Tasm. I, 123, implied (1860) ...	W.A.	—	—	—	—	—	—	B.fl.II,490	
GUNNERA, Linné, mantiss. plant. 16 (1767). (Milligania.)									
G. cordifolia, J. Hooker, Fl. Tasm. I, 125 (1860)	—	—	T.	—	—	—	—	B.fl.II,491	
CERATOPHYLLUM, Linné, syst. nat. 9 (1735); Linné, gen. plant. 290 (1737).									
C. demersum, Linné, spec. plant. 992 (1753)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,491	M.fr.VIII,163.
CALLITRICHE, Linné, syst. nat. ed. sext. 82 (1748).									
C. verna, Linné, flor. Suec. edit. sec. (1755)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,492	
C. Muelleri, Sonder in Linnaea XXVIII, 229 (1855)	—	—	—	V.	N.S.W.	—	—	—	M.fr.VIII,163.

RHIZOPHOREAE.

R. Brown in Flind. Voy. II, 549 (1814).

RHIZOPHORA, Linné, gen. plant. 137 (1737).										
R. mucronata, Lamarck, Encycl. méth. VI, 169 (1804)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,493	M.fr.IX,159.
CERIOPS, Arnott in Jardine, Ann. of nat. hist. I, 363 (1838).									
C. Candolleana, Arnott in Ann. of nat. hist. I, 364 (1838) ...	—	—	—	—	—	Q.	N.A.	B.fl.II,494	M.fr.IX,159.
BRUGUIERA, Lamarck, Encycl. méth. IV, 696, t. 397 (1796).									
B. Rheedii, Blume, enum. pl. Jav. 92 (1827)	—	—	—	—	—	Q.	N.A.	B.fl.II,494	
B. caryophylloides, Blume, enum. pl. Jav. 93 (1827)	—	—	—	—	—	—	—	—	M.fr.IX,159.
B. gymnorrhiza, Lamarck, Encycl. méth. IV, 696, t. 397 (1796)	—	—	—	—	—	Q.	N.A.	B.fl.II,495	M.fr.IX,159.
CARALLIA, Roxburgh, in Flind. Voy. II, 549 (1814).									
C. integerrima, De Candolle, prodr. III, 33 (1828)	—	—	—	—	—	Q.	N.A.	B.fl.II,495	M.fr.IX,159.

COMBRETACEAE.

R. Brown, prodr. 351 (1810).

TERMINALIA, Linné, mantiss. plant. 21 (1767). (Chuncoa.)									
T. platyptera, F. v. M., fragm. II, 151 (1861)	—	—	—	—	—	—	N.A.	B.fl.II,498	M.fr.II,151;IX,160.
T. volucris, R. Brown in Benth. Fl. Austr. II, 498 (1864) ...	—	—	—	—	—	—	N.A.	B.fl.II,498	
T. oblongata, F. v. M., fragm. II, 152 (1861)	—	—	—	—	—	Q.	—	B.fl.II,499	M.fr.III,167;IX,160.
T. bursarina, F. v. M., fragm. II, 149 (1861)	—	—	—	—	—	—	N.A.	B.fl.II,499	M.fr.II,149.
T. circumalata, F. v. M., fragm. III, 91 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,499	M.fr.III,91.
T. pterocarya, F. v. M., fragm. II, 152 (1861)	—	—	—	—	—	—	N.A.	B.fl.II,500	M.fr.II,152.
T. Thozetii, Bentham, Fl. Austr. II, 500 (1864)	—	—	—	—	—	Q.	—	B.fl.II,500	M.fr.IX,160.
T. Catappa, Linné, mantiss. pl. 128 (1767)...	—	—	—	—	—	Q.	—	—	M.fr.IX,160.
T. melanocarpa, F. v. M., fragm. III, 92 (1862)	—	—	—	—	—	Q.	N.A.	B.fl.II,500	M.fr.III,92;IX,160.
T. Muelleri, Bentham, Fl. Austr. II, 500 (1864)	—	—	—	—	—	Q.	—	B.fl.II,500	M.fr.IX,160.
T. latipes, Bentham, Fl. Austr. II, 501 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,501	
T. edulis, F. v. M., fragm. II, 151 (1861)	—	—	—	—	—	—	N.A.	B.fl.II,501	M.fr.II,151;IX,160.
T. discolor, F. v. M., fragm. III, 92 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,501	M.fr.III,92.
T. porphyrocarpa, F. v. M. in Benth. Fl. Austr. II, 501 (1864)	—	—	—	—	—	Q.	—	B.fl.II,501	
T. platyphylla, F. v. M., fragm. II, 150 (1861)	—	—	—	—	—	—	N.A.	B.fl.II,502	M.fr.II,150.
T. sericocarpa, F. v. M., fragm. IX, 159 (1875)	—	—	—	-	—	Q.	—	—	M.fr.IX,159.
T. microcarpa, Decaisne in Ann. du Mus. III (1834)	—	—	—	—	—	—	N.A.	B.fl.II,502	
T. petiolaris, Cunningham in Benth. Fl. Aust. II, 502 (1864) ...	—	—	—	-	—	—	N.A.	B.fl.II,502	
T. erythrocarpa, F. v. M., fragm. II, 150 (1861)	—	—	—	—	—	—	N.A.	B.fl.II,503	M.fr.II,150.
T. grandiflora, Bentham, Fl. Austr. II, 503 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,503	M.fr.IX,160.
LUMNITZERA, Willdenow, in den Verh. der Ges. Nat. Freunde zu Berl. IV. 186 (1803).									
L. coccinea, Wight & Arnott, prodr. 316 (1834)	—	—	—	—	—	Q.	—	B.fl.II,503	M.fr.IX,160.
L. racemosa, Willd. in Verh. der Nat. Freunde zu Berl. IV, 186 (1803)	—	—	—	—	—	Q.	N.A.	B.fl.II,504	M.fr.IX,160.

MACROPTERANTHES, F. v. M., fragm. III, 91 (1862).

M. montana, F. v. M., fragm. III, 91 (1862)	—	--	—	—	—	Q.	—	B.fl.II,504	M.fr.II,149;III,91&166.
M. Kekwickii, F. v. M., fragm. III, 151 (1863)	—	—	—	—	—	—	N.A.	B.fl.II,504	M.fr.III,151.
M. Leichhardtii, F. v. M., fragm. III, 91 (1862)	—	—	—	—	—	Q.	—	B.fl.II,505	M.fr.III,91.
M. Fitzalani, F. v. M., fragm. VIII, 160 (1874)	—	—	—	--	—	Q.	—	—	M.fr.VIII,160.

GYROCARPUS, N. Jacquin, select. stirp. Amer. hist. 282 (1763).

G. Americanus, N. Jacquin, select. stirp. Amer. hist. 282, t. 178 (1763)	—	—	—	—	—	Q.	N.A.	B.fl.II,505	M.fr.IX,160.

MYRTACEAE.

Adanson, Familles des plantes II, 86 (1763), from B. de Jussieu (1759).

ACTINODIUM, Schauer in Schlecht. Linnaea X, 311 (1835). (Triphelia.)

A. Cunninghamii, Schauer in Lindl. introd. to the nat. syst., sec. ed. 440 (1835)	W.A.	—	—	—	—	—	—	B.fl.III,5	

DARWINIA, Rudge in Transact. Linn. Soc. XI, 299 (1813). (Genetyllis, Chamaelaucium, Homoranthus, Hedaroma, Polyzone, Francisia, Decalophium, Schuermannia, Cryptostemon.)

D. macrostegia, Bentham in Journ. Linn. Soc. IX, 179 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,8	
D. Hookeriana, Bentham in Journ. Linn. Soc. IX, 179 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,9	M.fr.VIII,182.
D. fimbriata, Bentham in Journ. Linn. Soc. IX, 179 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,9	M.fr.VIII,182;IX,177.
D. speciosa, Bentham in Journ. Linn. Soc. IX, 179 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,9	M.fr.XI,10.
D. Meissneri, Bentham in Journ. Linn. Soc. IX, 179 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,9	M.fr.VIII,182;IX,177.
D. helichrysoides, Bentham in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,10	
D. Neildiana, F. v. M., fragm. IX, 177 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,177.
D. oederoides, Bentham in Journ. Linn. Soc. IX, 179 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,10	M.fr.IX,177;XI,9.
D. virescens, Bentham in Journ. Linn. Soc. IX, 179 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,10	M.fr.XI,10.
D. Oldfieldii, Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,10	
D. purpurea, Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,11	M.fr.XI,9. [177;XI,9.
D. citriodora, Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,11	M.fr.I,169;IV,174;IX,
D. thymoides, Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,11	M.fr.VIII,182;IX,176.
D. taxifolia, Cunningham in Field's N.S.Wales 352 (1825) ..	—	—	—	V.	N.S.W.	—	—	B.fl.III,12	M.fr.IX,176.
D. vestita, Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	W.A.	—	—	—	—		—	B.fl.III,12	M.fr.VIII,182;IX,176.
D. pauciflora, Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,12	M.fr.XI,9.
D. diosmoides, Bentham in Journ. Linn. Soc. IX, 180 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,13	M.fr.IX,176.
D. fascicularis, Rudge in Transact. Linn. Soc. XI, 299, t. 22 (1813)	—	—	—	—	N.S.W.	—	—	B.fl.III,13	M.fr.IX,176.
D. pinifolia, Bentham in Journ. Linn. Soc. IX, 181 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,14	
D. rhadinophylla, F. v. M., fragm. IX, 175 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,175.
D. sanguinea, Bentham in Journ. Linn. Soc. IX, 181 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,14	M.fr.IX,176.
D. micropetala, Bentham in Journ. Linn. Soc. IX, 181 (1865) ...	—	S.A.	—	—	—	—	—	B.fl.III,14	M.fr.I,12;IX,176.
D. Schuermanni, Bentham in Journ. Linn. Soc. IX, 181 (1865)...	—	S.A.	—	—	—	—	—	B.fl.III,14	M.fr.I,12.
D. verticordina, Bentham in Journ. Linn. Soc. IX, 181 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,15	M.fr.IV,57.
D. Thomasii, Bentham in Journ. Linn. Soc. IX, 181 (1865) ...	—	—	—	—	—	Q.	—	B.fl.III,15	M.fr.IV,137.
D. virgata, F. v. M., fragm. IX, 176 (1875)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,16	M.fr.IX,176.
D. ciliata, F. v. M.; Chamaelaucium, Desfontaines in Mém. du Mus. Par. V, 40, t. 3 (1821)	W.A.	—	—	—	—	—	—	B.fl.III,36	M.fr.VIII,182.
D. gracilis; Chamaelaucium, F. v. M., fragm. IV, 62 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.III,36	M.fr.IV,62;VIII,182.
D. Forrestii, F. v. M., fragm. XI, 9 (1878)...	W.A.	—	—	—	—	—	—	—	M.fr.XI,9.
D. heterandra, F. v. M.; Cham., Benth., Fl. Austr. III, 36 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,36	
D. Drummondii, F. v. M.; Chamaelaucium, Meissner in Journ. Linn. Soc. I, 44 (1857)	W.A.	—	—	—	—	—	—	B.fl.III,37	
D. Endlicheri, F. v. M.; Chamaelaucium virgatum, Endlicher in Ann. Wien. Mus. II, 193 (1838)	W.A.	—	—	—	—	—	—	B.fl.III,37	
D. brevifolia, F. v. M.; Cham., Benth., Fl. Austr. III, 37 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,37	
D. uncinata, F. v. M.; Cham., Schauer in Pl. Preiss. I, 97 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,37	
D. megalopetala; Cham., F. v. M. in Benth. Fl. Austr. III, 38 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,38	M.fr.VIII,182.
D. Turczaninowii, F. v. M.; Chamaelaucium pauciflorum, Benth., Fl. Austr. III, 38 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,38	M.fr.IX,176;XI,9.
D. axillaris, Cham., F. v. M. in Benth. Fl. Austr. III, 38 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,38	

VERTICORDIA, De Candolle in Dict. class. XI, 400 (1826). (Chrysorrhoea.)

V. Wilhelmii, F. v. M. in Transact. Vict. Inst. 122 (1855) ...	—	S.A.	—	—	—	—	—	B.fl.III,19	M.fr.X,28.
V. densiflora, Lindley in Bot. Regist. XXV, App. VI (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,20	
V. stelluligera, Meissner in Journ. Linn. Soc. I, 38 (1857) ...	W.A.	—	—	—	—	—	—	B.fl.III,20	
V. minutiflora, F. v. M., fragm. IV, 58 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,20	M.fr.IV,38.
V. Fontainesii, De Candolle, prodr. III, 209 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,21	M.fr.VIII,182;X,28.
V. helichrysantha, F. v. M. in Benth. Fl. Austr. III, 21 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,21	
V. Brownii, De Candolle, prodr. III, 209 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,21	
V. fastigiata, Turcz. in Bullet. de l'Ac. de St. Petersb. 402 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,22	M.fr.VIII,122;X,28.
V. Harveyi, Bentham, Fl. Austr. III, 22 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,22	
V. fimbrilepis, Turczaninow in Bull. Mosc. XX, 158 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,22	M.fr.X,28.
V. serrata, Schauer in nov. act. acad. Caes. XIX, suppl. II, 70 (1841)	W.A.	—	—	—	—	—	—	B.fl.III,23	
V. nitens, Schauer in nov. act. acad. Caes. XIX, suppl. II, 71, t. 4 (1841)	W.A.	—	—	—	—	—	—	B.fl.III,23	M.fr.X,28.
V. grandiflora, Endlicher in Ann. Wien. Mus. II, 195 (1838) ...	W.A.	—	—	—	—	—	—	B.fl.III,23	
V. chrysantha, Endlicher in Ann. Wien. Mus. II, 195 (1838) ...	W.A.	—	—	—	—	—	—	B.fl.III,24	M.fr.X,28.
V. Preissii, Schauer in Lehm. pl. Preiss. I, 101 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,24	M.fr.VIII,182.
V. acerosa, Lindley in Bot. Regist. XXV, App. VI (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,25	
V. polytricha, Bentham, Fl. Austr. III, 25 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,25	
V. oxylepis, Turcz. in Bullet. de l'Ac. de St. Petersb. 402 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,25	M.fr.VIII,182.
V. humilis, Bentham, Fl. Austr. III, 26 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,26	
V. penicillaris, F. v. M., fragm. I, 226 (1859)	W.A.	—	—	—	—	—	—	B.fl.III,26	M.fr.I,226.
V. multiflora, Turczaninow in Bull. Mosc. XX, 159 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,26	M.fr.X,28.
V. Huegelii, Endlicher in Hueg. enum. 16 (1837)...	W.A.	—	—	—	—	—	—	B.fl.III,27	

V. insignis, Endlicher in Hueg. enum. 47 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,27	M.fr.X,28.
V. habrantha, Schauer in Lehm. pl. Preiss. I, 100 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,28	M.fr.I,164.
V. monadelpha, Turczaninow in Bull. Mosc. XX, 158 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,28	M.fr.X,28.
V. Lehmanni, Schauer in Lehm. pl. Preiss. I, 99 (1844)... ...	W.A.	—	—	—	—	—	—	B.fl.III,29	
V. Cunninghamii, Schauer in nov. act. acad. Caes. XIX, suppl. II, 55 (1841)	—	—	—	—	—	—	N.A.	B.fl.III,29	M.fr.X,28.
V. picta, Endlicher in Ann. des Wien. Mus. II, 194 (1838) ...	W.A.	—	—	—	—	—	—	B.fl.III,30	M.fr.X,28.
V. pennigera, Endlicher in Hueg. enum. 46 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,30	M.fr.X,28.
V. Drummondii, Schauer in nov. act. acad. Caes. XIX, suppl. II, 56 (1841)	W.A.	—	—	—	—	—	—	B.fl.III,31	
V. Hughani, F. v. M., fragm. XI, 10 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,10.
V. pholidophylla, F. v. M., fragm. I, 227 (1859)	W.A.	—	—	—	—	—	—	B.fl.III,31	M.fr.I,227.
V. spicata, F. v. M., fragm. I, 226 (1859)	W.A.	—	—	—	—	—	—	B.fl.III,32	M.fr.I,226.
V. lepidophylla, F. v. M., fragm. I, 228 (1859)	W.A.	—	—	—	—	—	—	B.fl.III,32	M.fr.I,228.
V. ovalifolia, Meissner in Journ. Linn. Soc. I, 40 (1857)... ...	W.A.	—	—	—	—	—	—	B.fl.III,32	M.fr.X,28.
V. chrysostachya, Meissner in Journ. Linn. Soc. I, 41 (1857) ..	W.A.	—	—	—	—	—	—	B.fl.III,33	M.fr.X,28.
V. oculata, Meissner in Journ. Linn. Soc. I, 41 (1857)	W.A.	—	—	—	—	—	—	B.fl.III,33	
V. grandis, Drummond in Hook. Kew Misc. V, 119 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.III,33	M.fr.X,28;XI,10.
PILEANTHUS, Labillardière, Nov. Holl. pl. specim. II, 11 (1806).									
P. peduncularis, Endl. in Ann. des Wien. Mus. II, 196 (1838) ...	W.A.	—	—	—	—	—	—	B.fl.III,34	M.fr.I,225.
P. Limacis, Labillardière, Nov. Holl. pl. specim. II, 11, t. 149 (1806)	W.A.	—	—	—	—	—	—	B.fl.III,34	
P. filifolius, Meissner in Journ. Linn. Soc. I, 45 (1857)	W.A.	—	—	—	—	—	—	B.fl.III,35	
CALYCOTHRIX, Labillardière, Nov. Holl. pl. specim. II, 8, t. 146 (1806). (Calytrix, Calythrix.)									
C. aurea, Lindley, Bot. Regist. XXV, App. V, t. 3 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,41	
C. puberula, Meissner in Journ. Linn. Soc. I, 48 (1857)	W.A.	—	—	—	—	—	—	B.fl.III,42	
C. flavescens, Cunningham in Bot. Mag. 3323 (1834)	W.A.	—	—	—	—	—	—	B.fl.III,42	
C. asperula, Schauer in Lehm. pl. Preiss. I, 106 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,42	
C. sapphirina, Lindley, Bot. Regist. XXV, App. V (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,43	M.fr.I,224.
C. breviseta, Lindley, Bot. Regist. XXV, App. V (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,43	M.fr.VIII,182.
C. simplex, Lindley, Bot. Regist. XXV, App. V (1839)... ...	W.A.	—	—	—	—	—	—	B.fl.III,43	
C. empetroides, Schauer in nov. act. acad. Caes. XIX, suppl. II, 102 (1841)	W.A.	—	—	—	—	—	—	B.fl.III,44	
C. variabilis, Lindley, Bot. Regist. XXV, App. V (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,44	
C. muricata, F. v. M., fragm. I, 224 (1859)	W.A.	—	—	—	—	—	—	B.fl.III,44	M.fr.I,224.
C. gracilis, Bentham, Fl. Austr. III, 45 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,45	
C. brevifolia, Meissner in Journ. Linn. Soc. I, 46 (1857) ...	W.A.	—	—	—	—	—	—	B.fl.III,45	
C. brachyphylla, Turczaninow in Bull. Mosc. XX, 161 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,45	
C. Leschenaultii, Schauer in Lehm. pl. Preiss. I, 104 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,46	
C. Oldfieldii, Bentham, Fl. Austr. III, 46 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,46	
C. glutinosa, Lindley, Bot. Regist. XXV, App. V (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,46	
C. angulata, Lindley, Bot. Regist. XXV, App. VI (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,47	
C. depressa, Turczaninow in Bull. Soc. Mosc. XX, 162 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,47	
C. tenuifolia, Meissner in Journ. Linn. Soc. I, 46 (1857)... ...	W.A.	—	—	—	—	—	—	B.fl.III,47	
C. strigosa, Cunningham in Bot. Mag. 3323 (1834)	W.A.	—	—	—	—	—	—	B.fl.III,47	
C. plumulosa, F. v. M., fragm. X, 27 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,27.
C. decandra, R. Brown in De Cand. prodr. III, 208 (1828) ...	W.A.	—	—	—	—	—	—	B.fl.III,48	M.fr.I,146.
C. Birdii, F. v. M., fragm. X, 26 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,26.
C. tenuiramea, Turczaninow in Bull. Mosc. XXII, 20 (1849) ...	W.A.	—	—	—	—	—	—	B.fl.III,48	
C. Fraseri, Cunningham in Bot. Mag. 3323 (1834)	W.A.	—	—	—	—	—	—	B.fl.III,48	
C. granulosa, Bentham, Fl. Austr. III, 49 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,49	
C. Creswelli, F. v. M., fragm. X, 27 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,27.
C. microphylla, Cunningham in Bot. Mag. 3323 (1834)	—	—	—	—	—	—	N.A.	B.fl.III,49	
C. longiflora, F. v. M., fragm. I, 12 (1858)	—	S.A.	—	—	—	Q.	—	B.fl.III,49	M.fr.I,12.
C. leptophylla, Bentham, Fl. Austr. III, 50 (1866)	—	—	—	—	—	Q.	—	B.fl.III,50	
C. megaphylla, F. v. M., fragm. I, 146 (1859)	—	—	—	—	—	—	N.A.	B.fl.III,50	M.fr.I,146.
C. tetragona, Labillardière, Nov. Holl. pl. specim. II, 8, t. 146 (1806)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,50	M.fr.IV,36;VIII,183.
C. Sullivani, F. v. M., fragm. IX, 1 (1875)...	—	—	—	V.	—	—	—	—	M.fr.IX,1.
C. conferta, Cunningham in Bot. Mag. 3323 (1834)	—	—	—	—	—	—	N.A.	B.fl.III,51	
C. arborescens, F. v. M. in Transact. phil. Inst. Vict. III, 42 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,51	
C. brachychaeta, F. v. M. in Transact. phil. Inst. Vict. III, 43 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,52	
C. achaeta, F. v. M. in Transact. phil. Inst. Vict. III, 43 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,52	
C. laricina, R. Brown in Benth. Fl. Austr. III, 52 (1866) ...	—	—	—	—	—	—	N.A.	B.fl.III,52	
LHOTZKYA, Schauer in Linnaea X, 309 (1835).									
L. glaberrima, F. v. M., fragm. I, 13 (1858)	—	S.A.	—	—	—	—	—	B.fl.III,53	M.fr.I,13.
L. genetylloides, F. v. M. in Transact. phil. Soc. Vict. I, 16 (1854)	—	S.A.	—	V.	—	—	—	B.fl.III,54	
L. violacea, Lindley, Bot. Regist. XXV, App. VII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,54	M.fr.XI,8.
L. ciliata, F. v. M. in Benth. Fl. Austr. III, 54 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,54	
L. brevifolia, Schauer in Lehm. pl. Preiss. I, 103 (1844)... ...	W.A.	—	—	—	—	—	—	B.fl.III,54	
L. purpurea, F. v. M., fragm. I, 224 (1859)	W.A.	—	—	—	—	—	—	B.fl.III,55	M.fr.I,224.
L. ericoides, Schauer in Lindl. Introd. sec. ed. 439 (1835) ...	W.A.	—	—	—	—	—	—	B.fl.III,55	M.fr.XI,9.
L. acutifolia, Lindley, Bot. Regist. XXV, App. VII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,55	M.fr.XI,8.
L. Harvestiana, F. v. M., fragm. XI, 8 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,8.
THRYPTOMENE, Endlicher in den Ann. des Wien. Mus. II, 192 (1838). (Homalocalyx, Paryphantha, Astraea, Eremopyxis, Micromyrtus.)									
T. mucronulata, Turczaninow in Bull. Mosc. XX, 156 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,58	
T. australis, Endlicher in den Ann. des Wien. Mus. II, 192 (1838)	W.A.	—	—	—	—	—	—	B.fl.III,58	
T. tenella, Bentham, Fl. Austr. III, 59 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,59	M.fr.VIII,182;X,24.
T. prolifera, Turczaninow in Bull. Mosc. XXXV, 324 (1862) ...	W.A.	—	—	—	—	—	—	B.fl.III,59	

T. saxicola, Schauer in Lehm. pl. Preiss. I, 102 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,59	M.fr.IV,75.
T. Johnsonii, F. v. M., fragm. IV, 77 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,60	M.fr.IV,77.
T. racemulosa, Turczaninow in Bull. Mosc. XX, 156 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,60	
T. denticulata, Bentham in Fl. Austr. III, 60 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,60	M.fr.IV,75.
T. baeckeacea, F. v. M., fragm. IV, 65 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,60	M.fr.IV,65.
T. strongylophylla, F. v. M. in Benth. Fl. Austr. III, 61 (1866)	W.A.	—	—	—	—	–	—	B.fl.III,61	
T. hyporhytis, Turczaninow in Bull. Mosc. XXXV, 324 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,61	
T. Maisonneuvii, F. v. M., fragm. IV, 64 (1864)	—	S.A.	—	—	—	—	N.A.	B.fl.III,61	M.fr.IV,64.
T. urceolaris, F. v. M., fragm. X, 25 (1876)	W.A.	—	—	—	—	—	—	–	M.fr.X,25.
T. Mitchelliana, F. v. M., fragm. I, 11 (1858)	—	S.A.	—	V.	—	—	—	B.fl.III,61	M.fr.I,11;IX,62.
T. Miqueliana, F. v. M., fragm. I, 11 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,62	M.fr.I,11.
T. micrantha, J. Hooker in Hook. Kew Misc. V, 299, t. 8 (1853)	—	—	T.	—	—	—	—	B.fl.III,62	
T. stenocalyx, F. v. M., fragm. X, 23 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,23.
T. ericaea, F. v. M., fragm. I, 12 (1858)	—	S.A.	—	—	—	—	—	B.fl.III,62	M.fr.I,12.
T. oligandra, F. v. M., fragm. I, 11 (1858)...	W.A.	—	—	—	—	Q.	N.A.	B.fl.III,63	M.fr.I,11;IV,169.
T. elobata, F. v. M., fragm. IV, 63 (1864)...	W.A.	—	—	—	—	—	—	B.fl.III,64	M.fr.IV,63.
T. racemosa, F. v. M. in Benth. Fl. Austr. III, 64 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,64	
T. trachycalyx, F. v. M., fragm. X, 25 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,25.
T. imbricata, F.v.M.; Micromyrtus, Benth., Fl. Aus. III, 64 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,64	
T. flaviflora, F. v. M., fragm. VIII, 13 (1873)	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,25.
T. Drummondii, F. v. M.; Microm., Benth., Fl. Aus. III, 64 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,64	M.fr.VIII,183.
T. Elliotii, F. v. M., fragm. IX, 62 (1875)...	—	S.A.	—	—	—	—	—	—	M.fr.IX,62.
T. auriculata, F. v. M., fragm. X, 24 (1876)	—	S.A.	—	—	—	—	—	—	M.fr.X,24.
T. hymenonema, F. v. M., fragm. X, 26 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,26.
T. ciliata, F. v. M. in Woolls, pl. of the neighb. of Sydney 23 (1880)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,65	M.fr.I,30&242;IV,63.
T. minutiflora, F.v.M. in Woolls, pl. of the neighb. of Sydn. 23 (1880)	—	—	—	—	N.S.W.	—	—	B.fl.III,65	
T. leptocalyx, F. v. M.; Microm., Benth., Fl. Austr. III, 65 (1866)	—	—	—	—	—	Q.	—	B.fl.III,65	M.fr.I,30.
T. homalocalyx, F. v. M., fragm. IV, 63 (1864)	—	—	—	—	—	—	N.A.	B.fl.III,56	M.fr.IV,63.
T. polyandra, F. v. M., fragm. IV, 77 (1864)	—	—	—	—	—	—	N.A.	B.fl.III,56	M.fr.IV,77.
WEHLIA, F. v. M., fragm. X, 22 (1876).									
W. thryptomenoides, F. v. M., fragm. X, 22 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,22.
W. coarctata, F. v. M., fragm. X, 23 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,23.
BAECKEA, Linné, spec. plant. 358 (1753). (Jungia, Imbricaria, Scholtzia, Schidiomyrtus, Rinzia, Euryomyrtus, Camphoromyrtus, Tetrapora, Harmogia, Oxymyrrhine, Babingtonia, Ericomyrtus, Piptandra, Anticoryne.)									
B. platystemonea, Bentham, Fl. Austr. III, 74 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,74	M.fr.VIII,183.
B. Fumana, F. v. M., fragm. IV, 68 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,74	
B. dimorphandra, F. v. M. in Benth. Fl. Austr. III, 74 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,74	
B. schollerifolia, Lehmann in Lehm. pl. Preiss. II, 369 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,75	M.fr.VIII,183.
B. oxycoccoides, Bentham, Fl. Austr. III, 75 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,75	M.fr.VIII,183.
B. Drummondii, Bentham, Fl. Austr. III, 75 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,75	M.fr.VIII,183.
B. diffusa, Sieber in De Cand. prodr. III, 230 (1828)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,76	M.fr.IV,67.
B. crassifolia, Lindley in Mitch. Three Exped. II, 115 (1838) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,76	M.fr.IV,66.
B. cryptandroides, F. v. M., fragm. X, 29 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,29.
B. tetragona, F. v. M. in Benth. Fl. Austr. III, 77 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,77	
B. ericaea, F. v. M., fragm. I, 31 (1858)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,77	
B. polystemonea, F. v. M., fragm. II, 124 (1861)...	—	S.A.	—	—	—	—	—	B.fl.III,77	M.fr.II,124.
B. crenulata, R. Brown in Flind. voy. 548 (1814)...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,78	M.fr.IV,65.
B. brevifolia, De Candolle, prodr. III, 230 (1828)...	—	—	—	—	N.S.W.	—	—	B.fl.III,78	
B. Gunniana, Schauer in Walp. Rep. II, 920 (1843)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,78	M.fr.IV,66.
B. diosmifolia, Rudge in Transact. Linn. Soc. VIII, 298, t. 13 (1807)	—	—	—	—	N.S.W.	—	—	B.fl.III,79	M.fr.I,29.
B. leptocaulis, J. Hooker, icon. pl. t. 298 (1840)	—	—	T.	—	—	—	—	B.fl.III,79	
B. arbuscula, R. Brown in Benth. Fl. Austr. III, 79 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,79	
B. astarteoides, Bentham, Fl. Austr. III, 80 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,80	
B. linifolia, Rudge in Transact. Linn. Soc. VIII, 297, t. 12 (1807)	—	—	—	V.	N.S.W.	—	—	B.fl.III,80	M.fr.IV,71.
B. stenophylla, F. v. M., fragm. I, 13 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,80	M.fr.I,13.
B. camphorata, R. Brown in Bot. Mag. t. 2694 (1826)	—	—	—	—	N.S.W.	—	—	B.fl.III,81	M.fr.IV,70;IV,177.
B. virgata, Andrews, Bot. Reposit. t. 598 (1810)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.III,81	M.fr.I,69.
B. crenatifolia, F. v. M., fragm. IV, 70 (1864)	—	—	—	V.	—	—	—	B.fl III,82	M.fr.IV,70.
B. Cunninghamii, Bentham, Fl. Austr. III, 82 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,82	
B. densifolia, Smith in Transact. Linn. Soc. III, 260 (1797) ...	—	—	—	—	N.S.W.	—	—	B.fl.III,82	M.fr.IV,71.
B. Behrii, F. v. M., fragm. IV, 68 (1864)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,83	M.fr.IV,68.
B. uncinella, Bentham, Fl. Austr. III, 84 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,84	
B. polyandra, F. v. M., fragm. IV, 12 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,84	M.fr.II,30;IV,72.
B. corynophylla, F. v. M., fragm. IV, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,84	
B. pachyphylla, Bentham, Fl. Austr. III, 85 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,85	
B. crispiflora, F. v. M., fragm. IV, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,85	M.fr.II,31;IV,72.
B. Blackettii, F. v. M., fragm. VIII, 181 (1874)	W.A.	—	—	—	—	—	—	—	M.fr.VIII,181.
B. camphorosmae, Endlicher in Hueg. enum. 51 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,86	
B. pulchella, De Candolle, prodr. III, 230 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,86	
B. ochropetala, F. v. M., fragm. X, 29 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,29.
B. pygmaea, R. Brown in Benth. Fl. Austr. III, 86 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,86	
B. corymbulosa, Bentham, Fl. Austr. III, 87 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,87	M.fr.VIII,183.
B. floribunda, Bentham, Fl. Austr. III, 87 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,87	[183.
B. pentandra, F. v. M., fragm. II, 31; IV, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,87	M.fr.II,31;IV,72;VIII,
B. pentagonantha, F. v. M., fragm. IV, 73 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,87	M.fr.IV,73.
B. robusta, F. v. M., fragm. IV, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,88	M.fr.IV,72. [183.
B. ovalifolia, F. v. M., fragm. IV, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,88	M.fr.II,32;IV,72;VIII,
B. subcuneata, F. v. M., fragm. IV, 73 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,88	M.fr.IV,73.
B. grandiflora, Bentham, Fl. Austr. III, 89 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,89	
B. uberiflora; Scholtzia, F. v. M., fragm. IV, 74 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,67	M.fr.IV,74.

B. involucrata, Endlicher in Hueg. enum. 51 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,68	M.fr.IV,74.
B. spathulata, F. v. M.; Scholtzia, Benth., Fl. Austr. III, 68 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,68	
B. ciliata; Scholtzia, F. v. M., fragm. IV, 76 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,68	M.fr.IV,76.
B. capitata; Scholtzia, F. v. M. in Benth. Fl. Austr. III, 69 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,69	
B. umbellifera; Scholtzia, F. v. M., fragm. IV, 75 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.III,69	M.fr.IV,75.
B. laxiflora, F. v. M.; Scholtzia, Benth., Fl. Austr. III, 69 (1866)	W.A.	—	—	—	-	—	—	B.fl.III,69	
B. leptantha, F. v. M.; Scholtzia, Benth., Fl. Austr. III, 69 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,69	
B. parviflora; Scholtzia, F. v. M., fragm. IV, 76 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,70	M.fr.IV,76.
B. oligandra; Scholtzia, F. v. M. in Benth. Fl. Austr. III, 70 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,70	
B. serpillifolia, F. v. M., fragm. X, 30 (1876)	W.A.	—	—	-	—	—	—	B.fl.III,70	M.fr.VIII,183.
B. teretifolia, F. v. M.; Scholtzia, Benth., Fl. Austr. III, 70 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,70	
ASTARTEA, De Candolle in Dict. class. XI, 400 (1826).									
A. ambigua, F. v. M., fragm. II, 32 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,89	M.fr.II,32.
A. fascicularis, De Candolle, prodr. III, 210 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,90	M.fr.VIII,183.
A. intratropica, F. v. M., fragm. I, 83 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,90	M.fr.I,83.
HYPOCALYMMA, Endlicher in Hueg. enum. pl. Nov. Holl. Austr. occ. 51 (1837).									
H. xanthopetalum, F. v. M., fragm. II, 29 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,92	M.fr.II,29.
H. robustum, Endlicher in Hueg. enum. 50 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,92	
H. longifolium, F. v. M., fragm. II, 28 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,92	M.fr.II,28.
H. strictum, Schauer in Lehm. pl Preiss. I, 111 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,93	
H. tetrapterum, Turczaninow in Bull. Mosc. XXXV, 325 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,93	
H. linifolium, Turczaninow in Bull. Mosc. XXXV, 325 (1862) ...	W.A.	—	—	—	—	—	—	B.fl.III,93	
H. angustifolium, Endlicher in Hueg. enum. 50 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,94	
H. ericifolium, Bentham, Fl. Austr. III, 94 (1866)	W.A.	—	—	—	-	—	—	B.fl.III,94	
H. cordifolium, Lehmann, plant. Preiss. I, 112 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,94	
H. speciosum, Turcz. in Bull. de l'Acad. de St. Pet. 409 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,95	M.fr.VIII,183.
H. Phillipsii, Harvey in nat. hist. Rev. V, 296, t. 22 (1865) ...	W.A.	—	—	—	—	—	—	B.fl.III,95	
H. myrtifolium, Turcz. in Bullet. de St. Petersb. 410 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.III,95	M.fr.VIII,183.
BALAUSTION, Hooker, icones plantarum 852 (1852). (Punicella, Cheynia.)									
B. pulcherrimum, Hooker, icon. plant. 852 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,96	M.fr.VIII,183.
AGONIS, Lindley, Bot. Regist. XXV, p. X (1839). (Billiotia, Billotia.)									
A. spathulata, Schauer in Lehm. pl. Preiss. I, 117 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,97	M.fr.XI,120.
A. floribunda, Turczaninow in Bull. Mosc. II, 20 (1849) ...	W.A.	—	—	—	—	—	—	B.fl.III,97	
A. marginata, Schauer in Lehm. pl. Preiss. I, 117 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,98	M.fr.XI,120.
A. linearifolia, Schauer in Lehm. pl. Preiss. I, 118 (1844) ...	W.A.	-	—	—	—	—	—	B.fl.III,98	M.fr.VIII,183;XI,120.
A. juperina, Schauer in Lehm. pl. Preiss. I, 118 (1844)	W.A.	—	—	-	—	—	—	B.fl.III,98	M.fr.XI,120.
A. parviceps, Schauer in Lehm. pl. Preiss. I, 119 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,99	M.fr.XI,120.
A. obtusissima, F. v. M., fragm. XI, 119 (1881)	W.A.	—	—	—	—	-	—	—	M.fr.XI,119.
A. flexuosa, Schauer in Lehm. pl. Preiss. I, 116 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,99	M.fr.XI,120.
A. undulata, Bentham, Fl. Austr. III, 100 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,100	
A. hypericifolia, Schauer in Lehm. pl. Preiss. I, 117 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,100	M.fr.XI,120.
A. Scortechiniana, F. v. M., fragm. XI, 118 (1881)	—	—	—	—	—	Q.	—	—	M.fr.XI,118.
A. grandiflora, Bentham, Fl. Austr. III, 100 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,100	M.fr.XI,120.
LEPTOSPERMUM, R. et G. Forster, char. gen. 71, t. 36 (1776). (Fabricia, Homalospermum, Pericalymma.)									
L. Fabricia, Bentham, Fl. Austr. III, 102 (1866)	—	—	—	—	—	Q.	—	B.fl.III,102	
L. laevigatum, F. v. M., Annual Report 22 (1858)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,103	M.fr.IV,60.
L. firmum, Bentham, Fl. Austr. III, 104 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,104	
L. flavescens, Smith in Transact. Linn. Soc. III, 262 (1797) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,104	
L. scoparium, R. & G. Forster, char. gen. 48 (1776)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,105	
L. arachnoideum, Smith in Transact. Linn. Soc. III, 263 (1797)	—	—	—	—	N.S.W.	—	—	B.fl.III,105	
L. spinescens, Endlicher in Hueg. enum. 51 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,106	
L. lanigerum, Smith in Transact. Linn. Soc. III, 263 (1797) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,106	
L. parvifolium, Smith in Transact. Linn. Soc. III, 263 (1797) ...	—	—	—	—	N.S.W.	—	—	B.fl.III,107	
L. stellatum, Cavanilles, icon. pl. IV, 16, t. 330	—	—	—	—	N.S.W.	Q.	—	B.fl.III,107	
L. attenuatum, Smith in Transact. Linn. Soc. III, 262 (1797) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,108	
L. myrtifolium, Sieber in De Cand., prodr. III, 238 (1828) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,108	
L. rupestre, J. Hooker in Hook. icon. pl. t. 308 (1841)	—	—	T.	—	—	—	—	B.fl.III,108	
L. myrsinoides, Schlechtendal in Linnaea XX, 653 (1847) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,109	
L. erubescens, Schauer in Lehm. pl. Preiss. I, 121 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,109	M.fr.VIII,183.
L. abnorme, F. v. M. in Benth. Fl. Austr. III, 109 (1866) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,109	M.fr.II,27.
L. Roei, Bentham, Fl. Austr. III, 110 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,110	
L. floridum, Bentham, Fl. Austr. III, 110 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,110	
L. ellipticum, Endlicher in Hueg. enum. 51 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,110	
L. crassipes, Lehmann, ind. sem. hort. Hamb. (1842)	W.A.	—	—	—	—	—	—	B.fl.III,110	
KUNZEA, Reichenbach, conspect. regn. veget. 175 (1828). (Salisia, Pentagonaster.)									
K. micrantha, Schauer in Lehm. pl. Preiss. I, 125 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,112	M.fr.VIII,183.
K. eriocalyx, F. v. M., fragm. II, 28 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,112	M.fr.II,28.
K. Muelleri, Bentham, Fl. Austr. III, 113 (1866)	—	—	—	V.	N.S.W.	—	—	B.fl.III,113	
K. ericifolia, Reichenbach, consp. 175 (1828)	W.A.	—	—	—	—	-	—	B.fl.III,113	
K. Preissiana, Schauer in Lehm. pl. Preiss. I, 125 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,113	
K. recurva, Schauer in Lehm. pl. Preiss. I, 125 (1844)	W.A.	—	—	—	—	-	—	B.fl.III,114	M.fr.VIII,183.
K. micromera, Schauer in Lehm. pl. Preiss. III, 223 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,114	
K. pauciflora, Schauer in Lehm. pl. Preiss. I, 124 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,114	M.fr.VIII,183.
K. calida, F. v. M., fragm. VI, 23 (1867)	—	—	—	—	—	Q.	—	—	M.fr.VI,23.
K. parvifolia, Schauer in Lehm. pl. Preiss. I, 124 (1844) ...	—	—	—	—	N.S.W.	—	—	B.fl.III,115	M.fr.VIII,13.
K. peduncularis, F. v. M. in Transact. Vict. Inst. 144 (1855) ...	—	—	—	V.	N.S.W.	—	—	B.fl.III,115	M.fr.X,55.
K. corifolia, Reichenbach, consp. 175 (1828)	—	—	T.	V.	N.S.W.	-	—	B.fl.III,115	
K. capitata, Reichenbach, consp. 175 (1828)	—	—	—	—	N.S.W.	—	—	B.fl.III,116	
K. pomifera, F. v. M. in Transact. Vict. Inst. 124 (1855) ...	—	S.A.	—	V.	—	—	—	B.fl.III,116	

K. sericea, Turczaninow in Bull. Mosc. XX, 162 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,117	
K. Baxteri, Schauer in Lehm. pl. Preiss. I, 123 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,117	M.fr.III,153.
K. opposita, F. v. M., fragm. VI, 24 (1867)	—	—	—	—	N.S.W.	—	—	—	M.fr.VI,24.
CALLISTEMON, R. Brown in Flind. voy. II, 547 (1814).									
C. speciosus, De Candolle, prodr. III, 224 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,119	M.fr.I,14.
C. phoenicus, Lindley, Bot. Regist. XXV, App. X (1839)	W.A.	—	—	—	—	—	—	B.fl.III,119	M.fr.IV,53.
C. lanceolatus, De Candolle, prodr. III, 224 (1828)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,120	M.fr.IV,53.
C. coccineus, F. v. M., fragm. I, 13 (1858)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,120	M.fr.I,13.
C. salignus, De Candolle, prodr. III, 223 (1828)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,120	M.fr.IV,54.
C. paludosus, F. v. M., fragm. I, 14 (1858)	—	—	—	—	—	Q.	—	—	M.fr.I,14.
C. rigidus, R. Brown in Bot. Regist. t. 393 (1819)	—	—	—	—	N.S.W.	—	—	B.fl.III,121	
C. linearis, De Candolle, prodr. III, 223 (1828)	—	—	—	—	N.S.W.	—	—	B.fl.III,122	
C. pinifolius, De Candolle, prodr. III, 223 (1828)	—	—	—	—	N.S.W.	—	—	B.fl.III,122	
C. teretifolius, F. v. M. in Schlecht. Linnaea XXV, 387 (1852)	—	S.A.	—	—	—	—	—	B.fl.III,122	
C. brachyandrus, Lindley in Journ. Hort. Soc. IV, 112 (1849)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,122	M.fr.IV,52.
MELALEUCA, Linné, mantiss. plant. 14 (1767). (Gymnagathis, Asteromyrtus.)									
M. macronychia, Turcz. in Bullet. de St. Petersb. 420 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,129	M.fr.VIII,183.
M. lateritia, Otto in allgem. Gart. Zeit. II, 257 (1834)	W.A.	—	—	—	—	—	—	B.fl.III,130	
M. calothamnoides, F. v. M., fragm. III, 114 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,130	M.fr.III,114.
M. blaeriaefolia, Turczaninow in Bull. Mosc. XX, 165 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,130	
M. diosmifolia, Andrews, Bot. Reposit. t. 476 (1808)	W.A.	—	—	—	—	—	—	B.fl.III,130	
M. elliptica, Labillardière, Nov. Holl. pl. spec. II, 31, t. 173 (1806)	W.A.	—	—	—	—	—	—	B.fl.III,131	M.fr.VIII,183.
M. hypericifolia, Smith in Transact. Linn. Soc. III, 279 (1797)	—	—	—	—	N.S.W.	—	—	B.fl.III,131	
M. fulgens, R. Brown in Ait. hort. Kew. sec. ed. IV, 415 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,131	
M. acuminata, F. v. M., fragm. I, 15 (1858)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,132	M.fr.I,15.
M. leptoclada, Bentham, Fl. Austr. III, 132 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,132	
M. basicephala, Bentham, Fl. Austr. III, 133 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,133	
M. gibbosa, Labillardière, Nov. Holl. pl. spec. II, 30, t. 172 (1806)	—	S.A.	T.	V.	—	—	—	B.fl.III,133	
M. decussata, R. Brown in Ait. hort. Kew. sec. ed. IV, 415 (1812)	—	S.A.	—	V.	—	—	—	B.fl.III,133	
M. Wilsonii, F. v. M., fragm. II, 124, t. 15 (1861)	—	S.A.	—	V.	—	—	—	B.fl.III,134	M.fr.II,124.
M. thymifolia, Smith in Transact. Linn. Soc. III, 278 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,134	M.fr.X,55.
M. violacea, Lindley, Bot. Regist. XXV, App. VIII (1839)	W.A.	—	—	—	—	—	—	B.fl.III,135	
M. cardiophylla, F. v. M., fragm. I, 225 (1859)	W.A.	—	—	—	—	—	—	B.fl.III,135	M.fr.I,225;III,164.
M. undulata, Bentham, Fl. Austr. III, 135 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,135	
M. depauperata, Turcz. in Bullet. de St. Petersb. 424 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,136	M.fr.III,120;VIII,183.
M. lateriflora, Bentham, Fl. Austr. III, 136 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,136	
M. exarata, F. v. M., fragm. III, 114 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,136	M.fr.III,114;VIII,184.
M. fasciculiflora, Bentham, Fl. Austr. III, 137 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,137	M.fr.VIII,184.
M. teretifolia, Endlicher in Hueg. enum. 49 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,137	M.fr.III,117.
M. alsophila, Cunningham in Benth. Fl. Austr. III, 137 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,137	
M. acacioides, F. v. M., fragm. III, 116 (1862)	—	—	—	—	—	—	N.A.	B.fl.III,138	M.fr.III,116.
M. Baxteri, Bentham, Fl. Austr. III, 138 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,138	
M. symphyocarpa, F. v. M. in Transact. phil. Inst. Vict. III, 44 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,138	
M. angustifolia, Gaertner de fruct. I, 172, t. 35 (1788)	—	—	—	—	—	Q.	—	B.fl.III,139	
M. pauciflora, Turczaninow in Bull. Mosc. XX, 166 (1847)	—	—	—	—	N.S.W.	—	—	B.fl.III,139	
M. squarrosa, Donn, Hort. Cantabr. ed. sec. 101 (1800)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,139	M.fr.X,55.
M. adnata, Turcz. in Bullet. de St. Petersb. 425 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,140	M.fr.III,117;VIII,184.
M. linariifolia, Smith in Transact. Linn. Soc. III, 278 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,140	M.fr.X,55.
M. trichostachya, Lindley in Mitch. Trop. Austr. 277 (1848)	—	S.A.	—	—	—	Q.	N.A.	—	
M. radula, Lindley, Bot. Regist. XXV, App. VIII (1839)	W.A.	—	—	—	—	—	—	B.fl.III,141	M.fr.X,55.
M. pulchella, R. Brown in Ait. hort. Kew. sec. ed. IV, 414	W.A.	—	—	—	—	—	—	B.fl.III,141	
M. conferta, Bentham, Fl. Austr. III, 142 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,142	
M. Leucadendra, Linné, mantiss. 105 (1767)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,142	
M. lasiandra, F. v. M., fragm. III, 115 (1862)	—	—	—	—	—	—	N.A.	B.fl.III,143	M.fr.III,115.
M. genistifolia, Smith in Transact. Linn. Soc. III, 277 (1797)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,143	M.fr.I,15;X,55.
M. styphelioides, Smith in Transact. Linn. Soc. III, 275 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,144	
M. Huegelii, Endlicher in Hueg. enum. 48 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,144	
M. dissitiflora, F. v. M., fragm. III, 153 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,144	M.fr.III,153.
M. linophylla, F. v. M., fragm. III, 115 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,145	M.fr.III,115.
M. parviflora, Lindley, Bot. Regist. XXV, p. VIII (1839)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,145	
M. laxiflora, Turczaninow in Bullet. de St. Petersb. 421 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,145	M.fr.VIII,184;X,55.
M. armillaris, Smith in Transact. Linn. Soc. III, 277 (1797)	—	—	—	V.	N.S.W.	—	—	B.fl.III,146	
M. hamulosa, Turczaninow in Bull. Mosc. XX, 165 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,146	
M. subfalcata, Turczaninow in Bullet. de St. Petersb. 422 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,146	M.fr.III,119;VIII,184.
M. glaberrima, F. v. M., fragm. III, 119 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,147	M.fr.III,119.
M. rhaphiophylla, Schauer in Lehm. pl. Preiss. I, 143 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,147	
M. cymbifolia, Bentham, Fl. Austr. III, 148 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,148	
M. cuticularis, Labillard., Nov. Holl. pl. spec. II, 30, t. 171 (1806)	W.A.	—	—	—	—	—	—	B.fl.III,148	
M. sparsiflora, Turczaninow in Bull. Mosc. XX, 167 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,148	
M. calycina, R. Brown in Ait. hort. Kew. sec. ed. IV, 416 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,148	M.fr.VIII,184.
M. cordata, Turcz. in Bullet. de St. Petersb. 418 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,149	M.fr.VIII,184.
M. globifera, R. Brown in Ait. hort. Kew. sec. ed. IV, 411 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,149	
M. megacephala, F. v. M., fragm. III, 117 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,149	M.fr.III,117.
M. nesophila, F. v. M., fragm. III, 113 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,150	M.fr.III,113.
M. Oldfieldii, F. v. M., fragm. III, 118 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,150	
M. uncinata, R. Brown in Ait. hort. Kew. sec. ed. IV, 414 (1812)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,150	
M. concreta, F. v. M., fragm. III, 118 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,151	M.fr.III,118.
M. filifolia, F. v. M., fragm. III, 119 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,151	M.fr.III,119.
M. hakeoides, F. v. M. in Benth. Fl. Austr. III, 151 (1866)	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,151	
M. glomerata, F. v. M., Rep. Babb. Exped. 8 (1858)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.III,151	

M. pentagona, Labillardière, Nov. Holl. pl. spec. II, 27, t. 166 (1806)	W.A.	—	—	—	—	—	—	B.fl.III,152	M.fr.VIII,184.
M. ciliosa, Turczaninow in Bull. Soc. Mosc. XXXV, 326 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,152	
M. polycephala, Bentham, Fl. Austr. III, 152 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,152	M.fr.VIII,184.
M. spathulata, Schauer in Lehm. pl. Preiss. I, 134 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,153	M.fr.VIII,184.
M. eriantha, Bentham, Fl. Austr. III, 153 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,153	
M. subtrigona, Schauer in Lehm. pl. Preiss. I, 139 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,153	
M. seriata, Lindley, Bot. Regist. XXV, App. VIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,154	
M. scabra, R. Brown in Ait. hort. Kew. sec. ed. IV, 414 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,154	M.fr.VIII,184.
M. urceolaris, F. v. M. in Benth. Fl. Austr. III, 154 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,154	
M. trichophylla, Lindley, Bot. Regist. XXV, App. VIII (1839)	W.A.	—	—	—	—	—	—	B.fl.III,155	M.fr.III,114.
M. holosericea, Schauer in Lehm. pl. Preiss. I, 139 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,155	
M. squamea, Labillardière, Nov. Holl. pl. spec. II, 28, t. 168 (1806)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,155	
M. densa, R. Brown in Ait. hort. Kew. sec. ed. IV, 416 (1812)...	W.A.	—	—	—	—	—	—	B.fl.III,156	M.fr.VIII,184.
M. thymoides, Labillardière, Nov. Holl. pl. spec. II, 27, t. 167 (1806)	W.A.	—	—	—	—	—	—	B.fl.III,156	
M. striata, Labillardière, Nov. Holl. pl. spec. II, 26, t. 165 (1806)	W.A.	—	—	—	—	—	—	B.fl.III,157	
M. polygaloides, Schauer in Lehm. pl. Preiss. I, 142 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,157	
M. incana, R. Brown in Bot. Regist. t. 410 (1819)	W.A.	—	—	—	—	—	—	B.fl.III,157	
M. nodosa, Smith in Transact. Linn. Soc. III, 276 (1797) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,158	
M. pungens, Schauer in Lehm pl. Preiss. I, 138 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,158	M.fr.VIII,184.
M. ericifolia, Smith in Transact. Linn. Soc. III, 276 (1797) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,159	
M. pustulata, J. Hooker in Lond. Journ. VI, 476 (1844) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,160	
M. viminea, Lindley, Bot. Regist. XXV, App. VIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,159	
M. microphylla, Smith in Rees, Cyclop. XXIII (1812)	W.A.	—	—	—	—	—	—	B.fl.III,159	M.fr.III,116.
M. tenella, Bentham, Fl. Austr. III, 160 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,160	
M. leiocarpa, F. v. M., fragm. X, 55 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,55.
M. leiopyxis, F. v. M. in Bentham Fl. Austr. III, 160 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,160	
M. acerosa, Schauer in Lehm. pl. Preiss. I, 137 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,160	M.fr.III,118.
M. pauperiflora, F. v. M., fragm. III, 116 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,161	M.fr.III,116.
M. aspalathoides, Schauer in Lehm. pl. Preiss. I, 140 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,161	
M. cucullata, Turczaninow in Bullet. de St. Petersb. 424 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,161	M.fr.VIII,184.
M. minutifolia, F. v. M. in Transact. phil. Soc. Vict. III, 45 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.III,162	
M. foliolosa, Cunningham in Benth. Fl. Austr. III, 162 (1866)...	—	—	—	—	—	Q.	—	B.fl.III,162	
M. micromera, Schauer in Lehm. pl. Preiss. I, 146 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,162	
M. thyoides, Turczaninow in Bull. Soc. Mosc. XX, 167 (1847)...	W.A.	—	—	—	—	—	—	B.fl.III,162	M.fr.III,114.
M. tamariscina, Hooker in Mitch. Trop. Austr. 262 (1848) ...	—	—	—	—	—	Q.	—	B.fl.III,163	
CONOTHAMNUS, Lindley, Bot. Regist. XXV, App. IX (1839). (Trichobasis.)									
C. trinervis, Lindley, Bot. Regist. XXV, App. IX (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,163	
C. divaricatus, Bentham, Fl. Austr. III, 164 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,164	M.fr.VIII,184.
BEAUFORTIA, R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812). (Schizopleura.)									
B. sparsa, R. Brown in Ait. hort. Kew. sec. ed. IV, 419 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,165	
B. decussata, R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,166	
B. squarrosa, Schauer in nov. act. acad. Caes. XXI, 11 (1844)...	W.A.	—	—	—	—	—	—	B.fl.III,166	
B. orbifolia, F. v. M., fragm. III, 110 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,166	M.fr.III,110.
B. anisandra, Schauer in nov. act. acad. Caes. XXI, 13, t. I, (1844)	W.A.	—	—	—	—	—	—	B.fl.III,167	
B. macrostemon, Lindley, Bot. Regist. XXV, App. X (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,167	
B. heterophylla, Turczaninow in Bullet. de St. Petersb. 427 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,167	M.fr.VIII,184.
B. Schaueri, Preiss in nov. act. acad. Caes. XXI, 18 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,168	
B. purpurea, Lindley, Bot. Regist. XXV, App. X, t. 3 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,168	
B. Dampieri, Cunningham in Bot. Mag. t. 3272 (1833)	W.A.	—	—	—	—	—	—	B.fl.III,168	
B. elegans, Schauer in nov. act. acad. Caes. XXI, 20 (1844) ...	W.A.	—	—	—	—	—	N.A.	B.fl.III,169	
B. interstans, F. v. M., fragm. X, 30 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,30.
B. micrantha, Schauer in nov. act. acad. Caes. XXI, 22 (1844)...	W.A.	—	—	—	—	—	—	B.fl.III,169	M.fr.VIII,184.
REGELIA, Schauer in Linnaea XVII, 243 (1843).									[11.
R. grandiflora, Bentham, Fl. Austr. III, 170 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,170	M.fr.VIII,184; IX,46; XI,
R. ciliata, Schauer in nov. act. acad. Caes. XXI, 11 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,171	M.fr.XI,11.
R. inops, Schauer in Lehm. pl. Preiss. II, 224 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,171	M.fr.XI,11.
PHYMATOCARPUS, F. v. Mueller, fragm. III, 120 (1862).									
P. porphyrocephalus, F. v. M., fragm. III, 121 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,172	M.fr.III,121.
P. Maxwelli, F. v. M., fragm. IX, 45 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,45.
CALOTHAMNUS, Labillardière, Nov. Holl. pl. specim. II, 25 (1806). (Billotia.)									
C. pachystachyus, Bentham, Fl. Austr. III, 173 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,173	
C. longissimus, F. v. M., fragm. III, 112 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,174	M.fr.III,112.
C. blepharantherus, F. v. M., fragm. III, 112 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,174	M.fr.III,112; X,31&55.
C. sanguineus, Labillardière, Nov. Holl. pl. spec. II, 25, t. 164 (1806)	W.A.	—	—	—	—	—	—	B.fl.III,174	M.fr.X,31.
C. torulosus, Schauer in nov. act. acad. Caes. XXI, 25 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,175	M.fr.X,31.
C. gibbosus, Bentham, Fl. Austr. III, 175 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,175	
C. gracilis, R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,175	
C. blepharospermus, F. v. M., fragm. III, 111 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,175	M.fr.III,111.
C. Gilesii, F. v. M., fragm. X, 31 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,31.
C. chrysantherus, F. v. M., fragm. III, 112 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,176	M.fr.III,112.
C. Oldfieldii, F. v. M., fragm. III, 113 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,176	M.fr.III,113.
C. planifolius, Lehmann, delect. sem. hort. bot. Hamb. 7 (1842)	W.A.	—	—	—	—	—	—	B.fl.III,176	M.fr.X,31.
C. lateralis, Lindley, Bot. Regist. XXV, App. IX (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,177	
C. affinis, Turczaninow in Bullet. de St. Petersb. 428 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.III,177	M.fr.III,113; VIII,184.
C. Preissii, Schauer in nov. act. acad. Caes. XXI, 31 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,177	
C. Schaueri, Lehmann, delect. sem. hort. bot. Hamb. 7 (1842)...	W.A.	—	—	—	—	—	—	B.fl.III,178	
C. Lehmanni, Schauer in nov. act. acad. Caes. XXI, 31 (1844)...	W.A.	—	—	—	—	—	—	B.fl.III,178	
C. villosus, R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,178	
C. pinifolius, F. v. M., fragm. III, 153 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,179	M.fr.III,153.

C. rupestris, Schauer in nov. act. acad. Caes. XXI, 26 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,179		
C. quadrifidus, R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,179	M.fr.X,31.	
C. asper, Turczaninow in Bull. Mosc. XXII, 25 (1849)	W.A.	—	—	—	—	—	—	B.fl.III,180	M.fr.X,31.	
C. homalophyllus, F. v. M., fragm. III, 111 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,180	M.fr.X,31.	
LAMARCHEA, Gaudichaud in Freyc. Voy. Bot. 483, t. 110 (1826).										
L. hakeaefolia, Gaudichaud in Freyc. Voy. 484, t. 110 (1826) ...	W.A.	—	—	—	—	—	—	B.fl.III,123		
EREMAEA, Lindley, Bot. Regist. XXV, App. XI (1839).										
E. fimbriata, Lindley, Bot. Regist. XXV, App. XI (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,181	M.fr.XI,11.	
E. acutifolia, F. v. M., fragm. II, 30 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,181	M.fr.II,30.	
E. violacea, F. v. M., fragm. XI, 10 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,10.	
E. pilosa, Lindley, Bot. Regist. XXV, App. XI (1839)	W.A.	—	—	—	—	—	—	B.fl.III,182	M.fr.XI,11.	
E. ebracteata, F. v. M., fragm. II, 29 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,182	M.fr.XI,11.	
E. beaufortioides, Bentham, Fl. Austr. III, 182 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,182	M.fr.XI,11.	
ANGOPHORA, Cavanilles, icon. IV, 21, t. 331 (1797).										
A. cordifolia, Cavanilles, icon. IV, 21, t. 338 (1797)	—	—	—	—	N.S.W.	—	—	B.fl.III,183		
A. subvelutina, F. v. M., fragm. I, 31 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,184	M.fr.I,31.	
A. intermedia, De Candolle, prodr. III, 222 (1828)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,184		
A. lanceolata, Cavanilles, icon. IV, 22, t. 339 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,184		
EUCALYPTUS, L'Heritier, sertum Anglicum 18 (1788). (Eudesmia, Symphyomyrtus.)										
I. Renantherae.										
E. stellulata, Sieber in De Cand., prodr. III, 217 (1828)... ...	—	—	—	V.	N.S.W.	—	—	B.fl.III,200	M.fr.II,45;IX,172.	
E. pauciflora, Sieber in Sprengel, cur. poster. 195 (1827) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,201	M.fr.III,52.	
E. regnans, F. v. M. in Report Acclim. Soc. Vict. 20 (1870) ...	—	—	T.	V.	N.S.W.	—	—	—		
E. amygdalina, Labill., Nov. Holl. pl. spec. II, 14, t. 154 (1806)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,202	M.fr.II,53;VII,42&44.	
E. coccifera, J. Hooker in Lond. Journ. of Bot. VI, 477 (1847)...	—	—	T.	—	—	—	—	B.fl.III,204		
E. obliqua, L'Heritier, sertum Anglicum 18, t. 20 (1788) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,204	M.fr.II,44&171;XI,33.	
E. stricta, Sieber in De Candolle, prodr. III, 218 (1828)... ...	—	—	—	—	N.S.W.	—	—	B.fl.III,217	M.fr.XI,38.	
E. Planchoniana, F. v. M., fragm. XI, 43 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,43.	
E. Baileyana, F. v. M., fragm. XI, 37 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,37.	
E. sepulcralis, F. v. M., Eucalyptographia eight dec. (1882) ...	W.A.	—	—	—	—	—	—	—		
E. buprestium, F. v. M., fragm. III, 57 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,205	M.fr.III,57.	[XI,38.
E. marginata, Smith in Transact. Linn. Soc. VI, 302 (1802) ...	W.A.	—	—	—	—	—	—	B.fl.III,209	M.fr.II,40&174;VII,43;	
E. santalifolia, F. v. M. in Transact. Vict. Inst. I, 35 (1854) ...	W.A.	S.A.	—	—	—	—	—	B.fl.III,206		
E. macrorrhyncha, F. v. M., First general Report, 12 (1853) ...	—	—	—	V.	N.S.W.	—	—	B.fl.III,207	M.fr.XI,45.	
E. capitellata, Smith in White, Journ. Voy. N.S.W. 216 (1790)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,206	M.fr.II,173;XI,38.	
E. eugenioides, Sieber in Spreng. cur. poster. 195 (1827) ...	—	—	—	V.	N.S.W.	—	—	—		
E. piperita, Smith in White, Journ. Voy. N.S.Wales 226 (1790)	—	—	—	V.	N.S.W.	—	—	B.fl.III,207	M.fr.II,173;VII,44;XI,38	
E. pilularis, Smith in Transact. Linn. Soc. III, 284 (1797) ...	—	—	—	V.	N.S.W.	Q	—	B.fl.III,208	M.fr.II,61&172;VII,44;	
E. triantha, Link, enum. pl. hort. bot. Berol. II, 20 (1822) ...	—	—	—	—	N.S.W.	Q.	—	—		[XI,38.
E. haemastoma, Smith in Transact. Linn. Soc. III, 285 (1797) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,212	M.fr.II,51&175;VII,43;	
E. Sieberiana, F. v. M., Eucalyptogr. sec. decade (1879) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,202	M.fr.XI,37.	[XI,38.
E. microcorys, F. v. M., fragm. II, 50 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,212	M.fr.II,50.	
II. Porantherae.										
E. paniculata, Smith in Transact. Linn. Soc. III, 287 (1797) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,211	M.fr.II,174;VII,43.	
E. Leucoxylon, F. v. M. in Transact. Vict. Inst. I, 33 (1854) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,209	M.fr.II,60&175.	
E. melliodora, Cunningham, in Walp. rep. bot. syst. II, 924 (1843)	—	—	—	V.	N.S.W.	—	—	B.fl.III,210	M.fr.VII,43.	
E. polyanthema, Schauer in Walpers, rep. bot. syst. II, 924 (1843)	—	—	—	V.	N.S.W.	—	—	B.fl.III,213	M.fr.VII,44.	
E. populifolia, Hooker, icon. plant. 279 (1852)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,214&243		
E. ochrophloia, F. v. M., fragm. XI, 36 (1878)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.XI.36.	[XI,36.
E. gracilis, F. v. M. in Transact. Vict. Inst. I, 35 (1854) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,222	M.fr.VII,43;VIII,184;	
E. uncinata, Turczaninow in Bull. de Mosc. XXII, 23 (1849) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,216		
E. odorata, Behr in Schlecht. Linnaea XX, 657 (1847)	—	S.A.	—	—	—	—	—	B.fl.III,215	M.fr.II,66.	
E. largiflorens, F. v. M. in Transact. Vict. Inst. I, 34 (1854) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,215	M.fr.VII,43.	
E. Behriana, F. v. M. in Transact. Vict. Inst. I, 34 (1854) ...	—	S.A.	—	V.	—	—	—	B.fl.III,214		
E. hemiphloia, F. v. M., fragm. II, 62 (1860)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,216	M.fr.II,62.	
E. pruinosa, Schauer in Walp. rep. bot. syst. II, 926 (1843) ...	—	—	—	—	—	—	N.A.	B.fl.III,213	M.fr.III,132.	
E. melanophloia, F. v. M. in Journ. Linn. Soc. III, 93 (1858) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,220		
III. Parallelantherae.										
E. Cloeziana, F. v. M., fragm. XI, 44 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,44.	
E. Howittiana, F. v. M. in Wing's S. Sc. Record Aug. (1882) ...	—	—	—	—	—	Q.	—	—		
E. drepanophylla, F. v. M. in Benth. Fl. Austr. III, 221 (1866)	—	—	—	—	—	Q.	—	B.fl.III,221		
E. crebra, F. v. M. in Journ. Linn. Soc. III, 87 (1858)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,221		
E. decipiens, Endlicher in Hueg. enum. 49 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,218		
E. concolor, Schauer in Lehm. pl. Preiss. I, 129 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,247		
E. siderophloia, Bentham, Fl. Austr. III, 220 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,220	M.fr.II,61;VII,44.	
E. microtheca, F. v. M. in Journ. Linn. Soc. III, 87 (1858) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,223		
E. Raveretiana, F. v. M., fragm. X, 99 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,99.	
E. brachyandra, F. v. M. in Journ. Linn. Soc. III, 97 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.III,223		
E. pachyphylla, F. v. M. in Journ. Linn. Soc. III, 98 (1858) ...	—	S.A.	—	—	—	—	N.A.	B.fl.III,237		
E. pyriformis, Turczaninow in Bull. Mosc. XXII, 22 (1849) ...	W.A.	S.A.	—	—	—	—	—	B.fl.III,226	M.fr.II,32;X,5&84.	
E. macrocarpa, Hooker, icones plant. t. 405-407 (1842)	W.A.	—	—	—	—	—	—	B.fl.III,224	M.fr.II,41;XI,38.	
E. Preissiana, Schauer in Lehm. pl. Preiss. I, 131 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,232	M.fr.II,38;VII,42.	
E. alpina, Lindley in Mitch. Three Exped. II, 175 (1838) ...	—	—	—	V.	—	—	—	B.fl.III,225	M.fr.II,68;VII,42.	
E. Globulus, Labillardière, Voy. I, 153, t. 13 (1799)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,225	M.fr.II,68.	
E. megacarpa, F. v. M., fragm. II, 70 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,232	M.fr.VII,43.	
E. cosmophylla, F. v. M. in Transact. Vict. Inst. 32 (1854) ...	—	S.A.	—	—	—	—	—	B.fl.III,225	M.fr.VII,43.	
E. cordata, Labillardière, Nov. Holl. pl. spec. II, 13, t. 152 (1806)	—	—	T.	—	—	—	—	B.fl.III,224		
E. urnigera, J. Hooker in Lond. Journ. of Bot. VI, 477 (1847) ...	—	—	T.	—	—	—	—	B.fl.III,227		

E. longifolia, Link, enum. pl. hort. bot. Berol. II, 29 (1822) ...	—	—	—	V.	N.S.W.	—	—	B.fl.III,226	M.fr.II,50;VII,43.
E. erythronema, Turcz. in Bull. de l'acad. de St. Petersb. 415 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,227	M.fr.VIII,184.
E. caesia, Bentham, Fl. Austr. III, 227 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,227	M.fr.XI,40. [184.
E. tetraptera, Turczaninow in Bull. de Mosc. XXII, 22 (1849)...	W.A.	—	—	—	—	—	—	B.fl.III,228	M.fr.II,34;IV,172;VIII,
E. miniata, A. Cunningham in Walp. rep. bot. syst. II, 925(1843)	—	—	—	—	—	Q.	N.A.	B.fl.III,228	M.fr.XI,38&42.
E. phoenicea, F. v. M. in Journ. Linn. Soc. III, 91 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.III,251	
E. robusta, Smith, Specim. Bot. New Holl. 40, t. 13 (1793) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,228	M.fr.II,43&171;VII,44.
E. botryoides, Smith in Transact. Linn. Soc. III, 286 (1797) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,229	M.fr.II,48&175;VII,43.
E. goniocalyx, F. v. M., fragm. II, 48 (1860)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,229	M.fr.IV,52.
E. pallidifolia, F. v. M., fragm. III, 131 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,236	M.fr.III,131.
E. incrassata, Labillard., Nov. Holl. pl. spec. II, 12, t. 150 (1806)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,231	
E. oleosa, F. v. M. in Nederl. Kruidk. Arch. IV, 127 (1859) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,248	M.fr.II,56;XI,14.
E. goniantha, Turczaninow in Bullet. de Mosc. XX, 163 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,248	
E. falcata, Turczaninow in Bullet. de Mosc. XX, 163 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,248	
E. salmonophloia, F. v. M., fragm. XI, 11 (1878)...	W.A.	—	—	—	—	—	—	—	M.fr.XI,11.
E. leptopoda, Bentham, Fl. Austr. III, 238 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,238	M.fr.XI,11&13.
E. salubris, F. v. M., fragm. X, 54 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.XI,12.
E. angustissima, F. v. M., fragm. IV, 25 (1863)	W.A.	S.A.	—	—	—	—	—	B.fl.III,238	M.fr.IV,25.
E. Doratoxylon, F. v. M., fragm. II, 55 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,249	M.fr.II,55.
E. decurva, F. v. M., fragm. III, 130 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,249	M.fr.III,130.
E. Cooperiana, F. v. M., fragm. XI, 83 (1880)	W.A.	—	—	—	—	—	—	—	M.fr.XI,83.
E. corynocalyx, F. v. M., fragm. II, 43 (1860)	—	S.A.	—	V.	—	—	—	B.fl.III,218	M.fr.VII,43.
E. gomphocephala, De Candolle, prodr. III, 220 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,231	M.fr.II,36.
E. Oldfieldii, F. v. M., fragm. II, 37 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,237	M.fr.II,37.
E. orbifolia, F. v. M., fragm. V, 50 (1865)...	W.A.	—	—	—	—	—	—	B.fl.III,238	M.fr.V,50.
E. diversicolor, F. v. M., fragm. III, 131 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,251	M.fr.III,131.
E. patellaris, F. v. M. in Journ. Linn. Soc. III, 84 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.III,244	
E. tessellaris, F. v. M. in Journ. Linn. Soc. III, 88 (1858) ...	—	—	—	—	—	Q.	N.A.	B.fl.III,251	
E. clavigera, A. Cunningham in Walp. rep. bot. syst. II, 926 (1843)	—	—	—	—	—	—	N.A.	B.fl.III,250	
E. ferruginea, Schauer in Walp. repert. II, 926 (1843)	—	—	—	—	—	—	N.A.	B.fl.III,254	
E. grandifolia, R. Brown in Benth. Fl. Austr. III, 250 (1866) ...	—	—	—	—	—	—	N.A.	B.fl.III,250	
E. aspera, F. v. M. in Journ. Linn. Soc. III, 95 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,250	
E. patens, Bentham, Fl. Austr. III, 247 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,247	
E. Todtiana, F. v. M., in Wing's S. Sc. Record Aug. (1882) ...	W.A.	—	—	—	—	—	—	—	
E. vernicosa, J. Hooker in Lond. Journ. of Bot. VI, 478 (1847)...	—	—	T.	—	—	—	—	B.fl.III,232	
E. Gunnii, J. Hooker in Lond. Journ. Bot. III, 499 (1844) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,246	M.fr.II,62;VII,43;XI,38.
E. resinifera, Smith in White's Journ. Voy. N.S. Wales 231 (1790)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,245	M.fr.II,172;VII,43.
E. saligna, Smith in Transact. Linn. Soc. III, 285 (1797) ...	—	—	—	—	N.S.W.	—	—	B.fl.III,245	M.fr.II,173;VII,44.
E. punctata, De Candolle, prodr. III, 217 (1828)	—	—	—	—	N.S.W.	—	—	B.fl.III,242	M.fr.VII,43.
E. oligantha, Schauer in Walp. rep. II, 926 (1843)	—	—	—	—	—	—	N.A.	B.fl.III,213	
E. alba, Reinwardt in Blume, Bijdr. 1101 (1826)	—	—	—	—	—	Q.	N.A.	B.fl.III,243	
E. pulverulenta, Sims, Bot. Mag. t. 2089 (1819)	—	—	—	V.	N.S.W.	—	—	B.fl.III,224	M.fr.II,70.
E. Stuartiana, F. v. M. in Benth. Fl. Austr. III, 243 (1866) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,243	M.fr.II,62.
E. viminalis, Labillardière, Nov. Holl. pl. spec. II, 12, t. 151 (1806)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,239	M.fr.II,64;VII,44.
E. rostrata, Schlechtendal, Linnaea XX, 655 (1847)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,240	M.fr.VII,44.
E. tereticornis, Smith, Spec. Bot. New Holl. 41 (1793)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,241	
E. rudis, Endlicher in Hueg. enum. 49 (1837)	W.A.	—	—	—	-	—	-	B.fl.III,244	M.fr.XI,14.
E. foecunda, Schauer in Lehm. pl. Preiss. I, 130 (1844)	W.A.	—	—	—	-	—	—	B.fl.III,252	
E. reduncа, Schauer in Lehm. pl. Preiss. I, 127 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,253	M.fr.XI,15.
E. grossa, F. v. M. in Benth. Fl. Austr. III, 232 (1866)... ...	W.A.	—	—	—	-	—	—	B.fl.III,232	M.fr.VII,41.
E. obcordata, Turczaninow in Bull. Acad. Petersb. 416 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.III,234	M.fr.VII,44;VIII,184.
E. occidentalis, Endlicher in Hueg. enum. 49 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,235	M.fr.II,39;VII,43.
E. cornuta, Labillardière, Voy. I, 403, t. 20 (1799)	W.A.	—	—	—	—	—	—	B.fl.III,234	M.fr.II,39;XI,38.
E. gamophylla, F. v. M., fragm. XI, 40 (1878)	—	S.A.	—	—	—	—	N.A.	—	M.fr.XI,40.
E. perfoliata, R. Brown in Benth. Fl. Austr. III, 253 (1866) ...	—	—	—	—	—	—	N.A.	B.fl.III,253	
E. setosa, Schauer in Walpers, rep. bot. syst. II, 926 (1843) ...	—	—	—	—	—	Q.	N.A.	B.fl.III,254	M.fr.III,123;X,106.
E. Torelliana, F. v. M., fragm. X, 106 (1877)	—		—	—	—	Q.	—	—	M.fr.X,106.
E. peltata, Bentham, Fl. Austr. III, 254 (1866)	—	—	—	—	—	Q.	—	B.fl.III,254	
E. latifolia, F. v. M. in Journ. Linn. Soc. III, 94 (1858)... ...	—	—	—	—	—	—	N.A.	B.fl.III,255	
E. ptychocarpa, F. v. M. in Journ. Linn. Soc. III, 90 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.III,255	M.fr.VII,44.
E. Abergiana, F. v. M., fragm. XI, 41 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,41.
E. calophylla, R. Brown in Journ. Geogr. Soc. 20 (1831) ...	W.A.	—	—	—	—	—	-	B.fl.III,255	M.fr.II,35&171. [XI,15.
E. ficifolia, F. v. M., fragm. II, 85 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,256	M.fr.II,85&176;VI,25;
E. corymbosa, Smith. Specim. Bot. New Holl. 43 (1793)... ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,256	M.fr.II,46&174.
E. terminalis, F. v. M. in Journ. Linn. Soc. III, 89 (1858) ...	—	S.A.	—	—	—	Q.	N.A.	B.fl.III,257	
E. dichromophloia, F. v. M. in Journ. Linn. Soc. III, 89 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,257	
E. trachyphloia, F. v. M. in Journ. Linn. Soc. III, 90 (1858) ...	—	—	—	—	—	Q.	—	B.fl.III,221	
E. eximia, Schauer in Walpers, rep. bot. syst. II, 925 (1843) ...	—	—	—	—	N.S.W.	—	—	B.fl.III,258	
E. maculata, Hooker, icones plant. t. 619 (1844)	—	—	—	—	N.S.W.	Q.	-	B.fl.III,258	M.fr.II,47;VII,43.
E. Watsoniana, F. v. M., fragm. X, 98 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,98.
E. odontocarpa, F. v. M. in Journ. Linn. Soc. III, 98 (1858) ...	W.A.	—	—	—	—	-	—	B.fl.III,260	
E. tetrodonta, F. v. M. in Journ. Linn. Soc. III, 97 (1858) ...	—	—	-	-	—	Q.	N.A.	B.fl.III,260	
E. erythrocorys, F. v. M., fragm. II, 33 (1860)	W.A.	-	-	-	—	-	—	B.fl.III,258	M.fr.II,33.
E. tetragona, F. v. M., fragm. VI, 51 (1864)	W.A.	-	-	-	—	-	—	B.fl.III,259	M.fr.II,37.
E. eudesmioides, F. v. M., fragm. II, 35 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,260	M.fr.II,35.

TRISTANIA, R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812). (Lophostemon, Tristaniopsis.)

T. neriifolia, R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812)	—	—	—	—	N.S.W.	—	—	B.fl.III,261	M.fr.IV,56.
T. suaveolens, Smith in Rees, Cycl. XXXVI (1817)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,261	M.fr.I,81.
T. conferta, R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,263	M.fr.I,82.
T. lactiflora, F. v. M., fragm. I, 82 (1858)...	—	—	—	—	—	—	N.A.	B.fl.III,263	M.fr.I,82.

T. exiliflora, F. v. M., fragm. V, 11 (1865)...	—	—	—	—	—	Q.	—	B.fl.III,264	M.fr.IX,190.
T. laurina, R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,264	M.fr.I,81.
T. psidioides, Cunningham in Bot. Regist. XXII, n. 1839 (1837)	—	—	—	—	—	—	N.A.	B.fl.III,264	
T. umbrosa, Cunningham in Bot. Regist. XXII, n. 1839 (1837)...	—	—	—	—	—	—	N.A.	B.fl.III,265	
METROSIDEROS, Banks in Gaertner, de fruct. I, 170, t. 34 (1788). (Syncarpia, Kamptzia, Lysicarpus, Xanthostemon, Nania, Fremya, Cloezia, Spermalepis.)									
M. glomulifera, Smith in Transact. Linn. Soc. III, 269 (1797) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,265	M.fr.I,79.
M. leptopetala, F. v. M., Docum. Intercol. Exhib. 30 (1867) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,266	M.fr.I,79.
M. ternifolia, F. v. M., Docum. Intercol. Exhib. 30 (1867) ...	—	—	—	—	—	Q.	—	B.fl.III,267	
M. eucalyptoides, F. v. M., fragm. I, 243 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,267	M.fr.I,81&243.
M. polymorpha, Gaudichaud, Freyc. Voy. Bot. 99 t. 108 (1826)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,14.
M. nervulosa, C. Moore & F. v. M., fragm. VIII, 15 (1873) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.I,80.
M. chrysantha, F. v. M., fragm. IV, 159 (1864)	—	—	—	—	—	Q.	—	B.fl.III,268	M.fr.IV,159.
M. paradoxa, F. v. M., fragm. I, 243 (1859)	—	—	—	—	—	—	N.A.	B.fl.III,269	M.fr.I,80.
M. tetrapetala, F. v. M., fragm. VII, 41 (1870)	—	—	—	—	—	—	N.A.	—	M.fr.VII,41.
BACKHOUSIA, Hooker and Harvey in Bot. Mag. t. 4133 (1845).									
B. myrtifolia, Hooker & Harvey in Bot. Mag. t. 4133 (1845) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,269	M.fr.I,78;IX,43.
B. angustifolia, F. v. M., fragm. I, 79 (1858)	—	—	—	—	—	Q.	—	B.fl.III,270	M.fr.I,79;IX,43.
B. sciadophora, F. v. M., fragm. II, 26 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,270	M.fr.II,171.
B. citriodora, F. v. M., fragm. I, 78 (1858)	—	—	—	—	—	Q.	—	B.fl.III,270	M.fr.I,78.
OSBORNIA, F. v. Mueller, fragm. III, 30 (1862).									
O. octodonta, F. v. M., fragm. III, 31 (1862)	—	—	—	—	—	Q.	N.A.	B.fl.III,271	M.fr.IX,143.
RHODOMYRTUS, De Candolle, prodr. III, 240 (1828).									
R. psidioides, Bentham, Fl. Austr. III, 272 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,272	M.fr.II,86;IX,142.
R. trineura, F. v. M. in Benth. Fl. Austr. III, 272 (1866) ...	—	—	—	—	—	Q.	—	B.fl.III,272	M.fr.IV,117;IX,142.
R. cymiflora, F. v. M. in Benth. Fl. Austr. III, 273 (1866) ...	—	—	—	—	—	Q.	—	B.fl.III,273	M.fr.V,12;IX,149.
R. macrocarpa, Bentham, Fl. Austr. III, 273 (1866)	—	—	—	—	—	Q.	—	B.fl.III,273	M.fr.IX,142.
MYRTUS, Tournefort, inst. 640, t. 409 (1700), from Plinius.									
M. rhytisperma, F. v. M., fragm. I, 77 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,274	M.fr.IX,147.
M. tenuifolia, Smith in Transact. Linn. Soc. III, 280 (1797) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,274	M.fr.IX,147.
M. gonoclada, F. v. M. in Benth. Fl. Austr. III, 275 (1866) ...	—	—	—	—	—	Q.	—	B.fl.III,275	M.fr.IX,149.
M. lasioclada, F. v. M., fragm. IX, 148 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,148.
M. Hillii, Bentham, Fl. Austr. III, 275 (1866)	—	—	—	—	—	Q.	—	B.fl.III,275	M.fr.IX,149.
M. Becklerii, F. v. M., fragm. II, 85 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,275	M.fr.IX,147.
M. Bidwillii, Bentham, Fl. Austr. III, 275 (1866)	—	—	—	—	—	Q.	—	B.fl.III,275	
M. racemulosa, Bentham, Fl. Austr. III, 276 (1866)	—	—	—	—	—	Q.	—	B.fl.III,276	M.fr.IX,147.
M. acmenioides, F. v. M., fragm. I, 77 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,276	M.fr.IX,147.
M. fragrantissima, F. v. M. in Benth. Fl. Austr. III, 277 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,277	
M. Shepherdi, F. v. M., fragm. IX, 148 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,148.
RHODAMNIA, Jack in Hook. Comp. to the Bot. Mag. I, 153 (1835).									
R. sessiliflora, Bentham, Fl. Austr. III, 275 (1866)	—	—	—	—	—	Q.	—	B.fl.III,277	M.fr.IX,142.
R. trinervia, Blume, Mus. Bot. Lugd. I, 79 (1849)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,278	M.fr.I,76;IX,142.
R. argentea, Bentham, Fl. Austr. III, 276 (1866)...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,278	M.fr.IX,142.
R. Blairiana, F. v. M., fragm. IX, 141 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,141.
FENZLIA, Endlicher, Atacta 19, t. 17 & 18 (1833).									
F. obtusa, Endlicher, Atacta 19, t. 17 (1833)	—	—	—	—	—	Q.	—	B.fl.III,279	M.fr.IX,142.
F. retusa, Endlicher, Atacta 20, t. 18 (1833)	—	—	—	—	—	—	N.A.	B.fl.III,279	
DECASPERMUM, R. & G. Forster, char. gen. 73, t. 37 (1776). (Nelitris.)									
D. paniculatum, Baillon, hist. des plant. 350 (1871)	—	—	—	—	—	Q.	—	B.fl.III,279	M.fr.IV,56;IX,149.
EUGENIA, Micheli, nov. pl. gen. 226, t. 108 (1729). (Plinia, Plumier 1703, Acmene, Jambosa, Syzygium.)									
E. rariflora, Bentham in Hook. Lond. Journ. II, 221 (1843) ...	—	—	—	—	—	Q.	—	B.fl.III,282	M.fr.III,130;V,15;IX,143
E. Smithii, Poiret, Encycl. méthod. Suppl. III, 126 (1813) ...	—	—	—	—	V. N.S.W.	Q.	N.A.	B.fl.III,282	M.fr.IV,59;IX,145.
E. hemilampra, F. v. M., fragm. IX, 145 (1875)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IX,145.
E. Ventenatii, Bentham, Fl. Austr. III, 283 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,283	M.fr.IV,58;IX,145.
E. leptantha, Wight, Illustr. II, 15 (1850)	—	—	—	—	—	Q.	—	B.fl.III,283	M.fr.IX,143.
E. Moorei, F. v. M., fragm. V, 33 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,283	M.fr.IX,143.
E. corynantha, F. v. M., fragm. IX, 144 (1875)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IX,144.
E. cormiflora, F. v. M., fragm. V, 32 (1865)	—	—	—	—	—	Q.	—	B.fl.III,284	M.fr.V,32.
E. Hodgkinsoniae, F. v. M., fragm. IX, 145 (1875)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IX,145.
E. Tierneyana, F. v. M., fragm. V, 14 (1865)	—	—	—	—	—	Q.	—	B.fl.III,284	M.fr.IX,143.
E. grandis, Wight, Illustr. II, 17 (1850)	—	—	—	—	—	Q.	—	B.fl.III,285	M.fr.V,13;IX,143.
E. suborbicularis, Bentham, Fl. Austr. III, 285 (1866)	—	—	—	—	—	Q.	—	B.fl.III,285	M.fr.IX,143.
E. Wilsonii, F. v. M., fragm. V, 12 (1865)...	—	—	—	—	—	Q.	—	B.fl.III,285	M.fr.IX,143.
E. eucalyptoides, F. v. M., fragm. IV, 55 (1863)...	—	—	—	—	—	—	N.A.	B.fl.III,285	M.fr.IX,143.
E. myrtifolia, Sims, Bot. Mag. t. 2230 (1821)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,286	M.fr.I,225;IX,143.
E. angophoroides, F. v. M., fragm. V, 33 (1865)	—	—	—	—	—	Q.	—	B.fl.III,286	M.fr.IX,145.
E. Armstrongii, Bentham, Fl. Austr. III, 286 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,286	
E. oleosa, F. v. M., fragm. V, 15 (1865)	—	—	—	—	—	Q.	—	B.fl.III,287	M.fr.IX,146.
E. Dallachiana, F. v. M. in Benth. Fl. Austr. III, 287 (1866) ...	—	—	—	—	—	Q.	—	B.fl.III,287	M.fr.IX,145.
E. cryptophlebia, F. v. M., fragm. IX, 144 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,144.
ACICALYPTUS, A. Gray, New Gen. Un. St. Exped. 9 (1853). (Calyptranthes partly.)									
A. Fullageri, F. v. M., fragm. VIII, 15 (1873)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,15.
BARRINGTONIA, R. & G. Forster, char. gen. 75, t. 38 (1776). (Stravadium.)									
B. speciosa, R. & G. Forster, char. gen. 76, t. 38 (1776)... ...	—	—	—	—	—	Q.	—	B.fl.III,288	M.fr.IX,118.
B. racemosa, Gaudichaud in Freyc. Voy. Bot. 483, t. 107 (1826)	—	—	—	—	—	Q.	—	—	M.fr.IX,118.
B. acutangula, Gaertner, de fruct. II, 97, t. 101 (1791)	—	—	—	—	—	—	N.A.	B.fl.III,288	M.fr.IX,119.

CAREYA, Roxburgh, pl. Corom. III, 13, t. 217 (1816).
C. australis, F. v. M., fragm. V, 183 (1866) ... ... — — — — — Q. N.A. B.fl.III,289 M.fr.V,183.

SONNERATIA, Linné fil., suppl. plant. 38 (1781).
S. acida, Linné fil., suppl. plant. 38 (1781)... ... ... — — — — — — N.A. B.fl.III,301

MELASTOMACEAE.

A. L. de Jussieu, gen. plant. 328 (1789).

MEMECYLON, Linné, Fl. Zeyl. app. 9 (1747).
M. umbellatum, N. L. Burmann, Fl. Ind. 87 (1768) ... ... — — — — — Q. N.A. B.fl.III,293

OSBECKIA, Linné, spec. plant. I, 345 (1753).
O. Chinensis, Linné, spec. plant. 345 (1753) ... ... — — — — — Q. — B.fl.III,291 M.fr.IV,160.
O. Australiana, Naudin in Ann. sc. nat. trois. sér. XIV, 59 (1850) — — — — — — N.A. B.fl.III,291
O. perangusta, F. v. M., fragm. V, 181 (1866) ... ... — — — — — — N.A. — M.fr.V,181.

OTANTHERA, Blume in Flora, Regensb. IV, 488 (1831).
O. bracteata, Korthals in Verh. Nat. Gesch. der Ned. Bes. 235, t. 51 (1841) ... ... — — — — — Q. — B.fl.III,292

MELASTOMA, J. Burmann, thesaur. Zeyl. 72 (1737).
M. Malabathricum, Linné, spec. plant. 390 (1753) ... ... — — — — N.S.W. Q. N.A. B.fl.III,292

RHAMNACEAE.

A. L. de Jussieu, gen. plant. 376 (1789), from B. de Jussieu (1759).

VENTILAGO, Gaertner, de fruct. I, 223, t. 49 (1788).
V. viminalis, Hooker in Mitch. Trop. Austr. 369 (1848) ... — S.A. — — N.S.W. Q. N.A. B.fl.I,411 M.fr.IX,140.

ZIZYPHUS, Tournefort, inst. 627, t. 403 (1700), from Plinius (Ziziphus).
. Oenoplia, Miller, Gardeners Dictionary n. 3 (1731) ... ... — — — — — — N.A. B.fl.I,412
. Jujuba, Lamarck, Encycl. méth. III, 318 (1789) ... ... — — — — — Q. — B.fl.I,412
. quadrilocularis, F. v. M., fragm. III, 57 (1862) ... ... — — — — — — N.A. B.fl.I,412 M.fr.III,57.

DALLACHYA, F. v. M., fragm. IX, 140 (1875). (Rhamnus partly.)
D. Vitiensis, F. v. M., fragm. IX, 140 (1875) ... ... — — — — — Q. — — M.fr.XI,134.

BERCHEMIA, Necker, elem. bot. II, 122 (1790).
B. corollata, F. v. M., fragm. IX, 141 (1875) ... ... — — — — — Q. — — M.fr.IX,141.

COLUBRINA, L. C. Richard in Ann. des sc. nat. X, 368 (1827).
C. Asiatica, Brogniart in Ann. des sc. nat. X, 369 (1827) ... — — — — — Q. N.A. B.fl.I,413 M.fr.IX,139.

ALPHITONIA, Reisseck in Endl. gen. plant. 1098 (1840).
A. excelsa, Reisseck in Endl. gen. plant. 1098 (1840) ... ... — — — — N.S.W. Q. N.A. B.fl.I,414 M.fr.IX,140.

EMMENOSPERMUM, F. v. M., fragm. III, 63 (1862).
E. alphitonioides, F. v. M., fragm. III, 63 (1862)... ... ... — — — — N.S.W. Q. — B.fl.I,415 M.fr.IX,140.
E. Cunninghamii, Bentham, Fl. Austr. I, 415 (1863) ... ... — — — — — — N.A. B.fl.I,415

POMADERRIS, Labillardière, Nov. Holl. pl. specim. I, 61, t. 86 (1804).
P. lanigera, Sims, Bot. Mag. t. 1823 (1816) ... ... — — T. V. N.S.W. — — B.fl.I,416
P. elliptica, Labillardière, Nov. Holl. pl. spec. I, 61, t. 86 (1804) — — T. V. N.S.W. — — B.fl.I,417 M.fr.IX,139.
P. phillyroides, Sieber in De Cand., prodr. II, 33 (1825)... ... — — — — N.S.W. Q. — B.fl.I,418
P. vacciniifolia, Reisseck & F. v. M. in Linnaea XXIX, 266 (1856) — — — V. N.S.W. — — B.fl.I,418 M.fr.III,71.
P. myrtilloides, Fenzl in Hueg. enum. 22 (1837) ... ... W.A. — — — — — — B.fl.I,419 M.fr.IX,138.
P. grandis, F. v. M., fragm. III, 68 (1862)... ... ... W.A. — — — — — — B.fl.I,417 M.fr.IX,139.
P. Calvertiana, F. v. M., fragm. IX, 138 (1875) ... ... — — — — N.S.W. — — — M.fr.IX,138.
P. ledifolia, Cunningham in Field, N.S.Wales 351 (1825) ... — — — V. N.S.W. — — B.fl.I,419 M.fr.III,69.
P. apetala, Labillardière, Nov. Holl. pl. spec. I, 62, t. 87 (1804) — S.A. T. V. N.S.W. — — B.fl.I,419 M.fr.III,69&73;IX,139.
P. cinerea, Bentham, Fl. Austr. I, 420 (1863) ... ... — — — V. N.S.W. — — B.fl.I,420
P. prunifolia, Cunningham in Hueg. enum. 22 (1837) ... ... — — — V. N.S.W. — — B.fl.I,420 M.fr.III,75;IX,139.
P. ligustrina, Sieber in De Cand., prodr. II, 34 (1825) ... ... — — — — N.S.W. — — B.fl.I,420 M.fr.III,71;IX,139.
P. betulina, Cunningham in Bot. Mag. t. 3212 (1833) ... ... — — — V. N.S.W. — — B.fl.I,421 M.fr.III,76;IX,139.
P. Forrestiana, F. v. M., fragm. IX, 139 (1875) ... ... W.A. — — — — — — — M.fr.IX,139.
P. obcordata, Fenzl in Hueg. enum. 23 (1837) ... ... W.A. S.A. — — — — — B.fl.I,421 M.fr.III,73.
P. racemosa, Hooker, Journ. of Bot. I, 256 (1834) ... ... — S.A. T. V. N.S.W. — — B.fl.I,421 M.fr.III,75&168;IX,139.
P. subrepanda, Reisseck & F. v. M. in Linnaea XXIX, 267 (1856) — — — V. — — — B.fl.I,422 M.fr.III,74.
P. elachophylla, F. v. M., fragm. II, 131 (1861) ... ... — — — V. — — — B.fl.I,422 M.fr.II,166;IX,139.
P. phylicifolia, Loddiges, Bot. Cabinet t. 120 (1818) ... ... — — T. V. — — — B.fl.I,422 M.fr.III,73.
P. intangenda, F. v. M., fragm. X, 52 (1876) ... ... W.A. — — — — — — — M.fr.X,53.

CRYPTANDRA, Smith in Transact. Linn. Soc. IV, 217 (1798). (Trymalium, Spyridium, Stenanthemum, Wichurea, Stenodiscus.)
C. albicans, F. v. M.; Trym., Reisseck in pl. Preiss. II, 280 (1847) W.A. — — — — — — B.fl.I,423
C. Billardieri, F. v. M.; Trym., Fenzl in Hueg. enum. 25 (1837) W.A. — — — — — — B.fl.I,423
C. ledifolia, F. v. M.; Trymalium, Fenzl in Hueg. enum. 24 (1837) W.A. — — — — — — B.fl.I,424
C. Daltoni; Trymalium, F. v. M., fragm. IX, 135 (1875) ... — — — V. — — — — M.fr.IX,135.
C. angustifolia, F. v. M.; Trym., Reisseck in pl. Preiss. II, 284 (1847) W.A. — — — — — — B.fl.I,424 M.fr.IX,135.
C. Wichurae, F. v. M.; Trym., Nees in Lehm. pl. Preiss. II, 281 (1847) W.A. — — — — — — B.fl.I,425 M.fr.IX,135.
C. ericifolia, Smith in Transact. Linn. Soc. X, 294, t. 18 (1811)... — — — N.S.W. Q. — B.fl.I,438 M.fr.IX,137.
C. propinqua, Cunningham in Hueg. enum. 23 (1837) ... ... — S.A. — — N.S.W. — — B.fl.I,442 M.fr.IX,137.
C. hispidula, Reisseck & F. v. M. in Linnaea XXIX, 294 (1856) — S.A. — — — — — B.fl.I,439
C. spyridioides, F. v. M., fragm. III, 68 (1862) ... ... W.A. — — — — — — B.fl.I,439 M.fr.IX,138.
C. scoparia, Reisseck in Lehm. pl. Preiss. II, 283 (1847) W.A. — — — — — — B.fl.I,439
C. spinescens, Sieber in De Cand., prodr. II, 38 (1825) ... ... — — — — N.S.W. — — B.fl.I,439
C. amara, Smith in Transact. Linn. Soc. X, 295, t. 18 (1811) ... — S.A. T. V. N.S.W. Q. — B.fl.I,440 M.fr.III,66&67;IX,137.
C. lanosiflora, F. v. M., fragm. III, 65 (1862) ... ... — — — — N.S.W. — — B.fl.I,440 M.fr.III,65.
C. tomentosa, Lindley in Mitch. Three Exped. II, 178 (1838) ... — S.A. — V. N.S.W. — — B.fl.I,441 M.fr.III,67;IX,138.
C. nutans, Steudel in Lehm. pl. Preiss. I, 186 (1844) ... ... W.A. — — — — — — B.fl.I,441 M.fr.IX,138.

C. glabriflora, Bentham, Fl. Austr. I, 441 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,441	
C. alpina, J. Hooker, Fl. Tasm. I, 75, t. 12 (1860)	—	—	T.	—	—	—	—	B.fl,I,441	
C. Leucopogon, Meissner in Lehm. pl. Preiss. II, 287 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.I,442	
C. parvifolia, Turczaninow in Bull. de Mosc. I, 459 (1858) ...	W.A.	—	—	—	—	—	—	B.fl.I,442	
C. buxifolia, Fenzl in Hueg. enum. 23 (1837)	—	—	—	—	N.S.W.	—	—	B.fl.I,442	M.fr.IX,138.
C. pungens, Steudel in Lehm. pl. Preiss. I, 187 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,443	M.fr.IX,138.
C. mutila, Nees in Lehm. pl. Preiss. II, 289 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,443	
C. longistaminea, F. v. M., fragm. III, 64 (1862)...	—	—	—	—	N.S.W.	—	—	B.fl.I,443	M.fr.IX,138.
C. arbutiflora, Fenzl in Hueg. enum. 26 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,444	M.fr.IX,138.
C. miliaris, Reisseck in Lehm. pl. Preiss. II, 288 (1847)... ...	W.A.	—	—	—	—	—	—	B.fl.I,444	
C. nudiflora, F. v. M., fragm. III, 64 (1862)	W.A.	—	—	—	—	—	—	B.fl.I.444	M.fr.III,64.
C. ulicina, Hooker, Journ. of Bot. I, 257 (1834)	—	—	T.	—	—	—	—	B.fl.I,434	M.fr.IX,137.
C. pomaderroides, Reisseck in Endl. nov. stirp. dec. 29 (1839)...	W.A.	—	—	—	—	—	—	B.fl.I,435	M.fr.IX,137.
C. pumila; Spyridium, F. v. M., fragm. IX, 137 (1875)... ...	W.A.	—	—	—	—	—	—	—	M.fr.IX,137.
C. leucophracta, Schlechtendal, Linnaea XX, 640 (1847) ...	—	S.A.	T.	V.	—	—	—	B.fl.I,435	M.fr.III,77;IX,136.
C. coronata, Reisseck in Lehm. pl. Preiss. II, 288 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.I,436	M.fr.IX,137.
C. humilis, F. v. M.; Stenanthemum, Benth., Fl. Austr. I, 436 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,436	
C. Waterhousii; Spyridium, F. v. M., fragm. III, 83 (1862) ..	—	S.A.	—	—	—	—	—	B.fl.I,436	M.fr.III,83.
C. tridentata, Steudel in Lehm. pl. Preiss. I, 186 (1844)... ...	W.A.	—	—	—	—	—	—	B.fl.I,427	M.fr.IX,137.
C. divaricata, F. v. M.; Spyridium, Bentham, Fl. Austr. I, 427 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,427	
C. obcordata, J. Hooker, Fl. Tasm. I, 71 (1860)	—	—	T.	V.	—	—	—	B.fl.I,427	M.fr.III,80.
C. Lawrencii, J. Hooker, Fl. Tasm. I, 72 (1860)	—	—	T.	—	—	—	—	B.fl.I,430	M.fr.IX,136.
C. Hookeri, F. v. M.; C. parvifolia, J. Hooker, Fl. Tasm. I, 73 (1860)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,428	M.fr.III,79;IX,136.
C. spadicea, F. v. M.; Trymalium, Fenzl in Hueg. enum. 26 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,428	M.fr.IX,137.
C. globulosa, F. v. M.; Pomad., G. Don in Loud. hort. Brit. 84 (1830)	W.A.	—	—	—	—	—	—	B.fl.I,429	M.fr.IX,137.
C. obovata, J. Hooker, Fl. Tasm. I, 74 (1860)	—	S.A.	T.	—	—	—	—	B.fl.I,429	M.fr.IX,136.
C. spathulata; Trymalium, F. v. M. in Trans. Vict. Inst. I, 122 (1855)	—	S.A.	—	—	—	—	—	B.fl.I,430	M.fr.IX,136.
C. cordata, Turczaninow in Bull. de Mosc. XXXI, 459 (1858) ...	W.A.	—	—	—	—	—	—	B.fl.I,430	M.fr.IV,25;IX,137.
C. rotundifolia; Spyridium, F. v. M., fragm. IV, 25 (1863) ...	W.A.	—	—	—	—	—	—	—	M.fr.IV,25.
C. phlebophylla; Trymalium, F. v. M. in Linnaea XXIX, 272 (1856)	—	S.A.	—	—	—	—	—	B.fl.I,430	
C. coactilifolia, F. v. M.; Spyr., Reiss. in Linnaea XXIX, 291 (1856)	—	S.A.	—	—	—	—	—	B.fl.I,431	M.fr.IX,136.
C. complicata; Spyridium, F. v. M., fragm. III, 78 (1862) ...	W.A.	—	—	—	—	—	—	B.fl.I,431	M.fr.III,78.
C. westringifolia, F.v.M.; Trym., Reiss. in pl. Preiss. II, 284 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,431	
C. villosa, Turczaninow in Bull. de Mosc. XXXI, 458 (1858) ...	W.A.	—	—	—	—	—	—	B.fl.I,432	
C. pauciflora, Turczaninow in Bull. de Mosc. XXXI, 458 (1858)	W.A.	—	—	—	—	—	—	B.fl.I,432	
C. halmaturina; Trym., F. v. M. in Linnaea XXIX, 283 (1856)	—	S.A.	—	—	—	—	—	B.fl.I,432	
C. bifida; Trymalium, F. v. M. in Linnaea XXIX, 282 (1856)...	—	S.A.	—	—	—	—	—	B.fl.I,432	M.fr.IX,136.
C. subochreata; Trym., F. v. M. in Trans. Vict. Inst. I, 122 (1855)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,432	M.fr.III,82;IX,136.
C. vexillifera, Hooker, Journ. Bot. I, 257 (1834)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,433	M.fr.III,81;IX,136.
COLLETIA, Commerçon in A. L. de Jussieu, gen. 380 (1789). (Discaria, Tetrapasma.)									
C. pubescens, Brogniart in Annal. des sc. nat. X, 366 (1827) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.I,445	M.fr.III,83;IV,175.
GOUANIA, N. J. Jacquin, stirp. Amer. hist. 263, t. 179 (1763).									
G. Hillii, F. v. M., fragm. IX, 163 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,163.
G. Australiana, F. v. M., fragm. IV, 144 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,144.

ARALIACEAE.

Ventenat, Tabl. III (1799). (Umbelliferarum subordo.)

ASTROTRICHA, De Candolle, Mémoire sur la fam. des Ombellifères 25 (1829).												
A. pterocarpa, Bentham, Fl. Austr. III, 379 (1866)	—	—	—	—	—	Q.	—	B.fl.III,379	M.fr.VII,148.
A. floccosa, De Candolle, Mém. des Ombell. 30, t. 5 (1829) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,380	
A. longifolia, Bentham in Hueg. enum. 55 (1837)...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,380	M.fr.VII,148.
A. ledifolia, De Candolle, Mém. des Ombell. 30, t. 6 (1829) ...	—	—	—	V.	N.S.W.	—	—	B.fl.III,380	M.fr.VII,148.
A. Hamptoni, F. v. M., fragm. VI, 125 (1868)	W.A.	—	—	—	—	—	—	—	M.fr.VI,125&250
POROSPERMUM, F. v. M., fragm. VII, 94 (1870).									
P. Michieanum, F. v. M., fragm. VII, 95 (1870)	—	—	—	—	—	Q.	—	—	M.fr.VII,95.
PANAX, Linné, spec. plant. edit. sec. 1513 (1763). (Nothopanax.)									
P. Gunnii, J. Hooker in Lond. Journ. of Bot. VI, 466 (1847) ...	—	—	T.	—	—	—	—	B.fl.III,381	
P. Murrayi, F. v. M., fragm. II, 106 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,381	M.fr.VII,96.
P. mollis, Bentham, Fl. Austr. III, 382 (1866)	—	—	—	—	—	Q.	—	B.fl.III,382	M.fr.VII,96.
P. Macgillivraei, Bentham, Fl. Austr. III, 382 (1866)	—	—	—	—	—	Q.	—	B.fl.III,382	
P. sambucifolius, Sieber in De Cand. prodr. III, 255 (1825) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,382	M.fr.VII,95.
P. cephalobotrys, F. v. M., fragm. II, 83 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,382	M.fr.II,83;VII,95.
P. elegans, C. Moore & F. v. M. in Trans. phil. Inst. Vict. II, 68 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,383	
P. Cissodendron, C. Moore & F. v. M., fragm. VII, 96 (1870) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,96.
MOTHERWELLIA, F. v. M., fragm. VII, 107 (1871).									
M. haplosciadea, F. v. M., fragm. VII, 107 (1871)	—	—	—	—	—	Q.	—	—	M.fr.VII,107.
CISSODENDRON, Seemann, Journ. of Bot. II, 303 (1864). (Hedera partly, Irvingia, Kissodendron.)									
C. Australianum, Seemann, Journ. of Bot. II, 303 (1864) ...	—	—	—	—	—	Q.	—	B.fl.III,384	M.fr.IV,120;V,18.
HEPTOPLEURUM, Gaertner, de fruct. II, 472, t. 178 (1791). (Paratropia.)									
H. venulosum, Seemann, Journ. of Bot. III, 80 (1865)	—	—	—	—	—	Q.	—	B.fl.III,384	
BRASSAIA, Endlicher, nov. stirp. decad. I, 89 (1839).									
B. actinophylla, Endlicher, nov. stirp. dec. I, 89 (1839)	—	—	—	—	—	Q.	—	B.fl.III,385	
MERYTA, R. & G. Forster, char. gen. 119, t. 60 (1776). (Bothryodendron.)									
M. latifolia, Seemann in Bonplandia, 295 (1862)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.
M. angustifolia, Seemann in Bonplandia, 296 (1862)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.
MACKINLAYA, F. v. M., fragm. IV, 120 (1864).									
M. macrosciadea, F. v. M., fragm. IV, 120 (1864)	—	—	—	—	—	Q.	—	B.fl.III,383	M.fr.II,108&176;IV,120.

UMBELLIFERAE.

Morison, hist. pl. II, lib. 3, sect. 9 (1680).

HYDROCOTYLE, Tournefort, inst. 328, t. 173 (1700).

H. vulgaris, Linné, spec. plant. 234 (1753)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,339	
H. pedunculariis, R. Brown in annal. des sc. phys. VI, 62, t. 61 (1820)					—	—	T.	—	—	—	—	B.fl.III,339	
H. hirta, R. Brown in annal. des sc. phys. VI, 64 (1820)			...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,339	
H. laxiflora, De Candolle, prodr. IV, 61 (1830)	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,340	M.fr.IV,179&180.
H. rhombifolia, F. v. M., fragm. VII, 147 (1871)	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,147.
H. pedicellosa, F. v. M., fragm. IV, 182 (1864)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,341	M.fr.VII,147.
H. tripartita, R. Brown in annal. des sc. phys. VI, 69, t. 61 (1820)				—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,341	M.fr.VII,147.
H. plebeia, R. Brown in annal. des sc. phys. VI, 46, t. 60 (1820)				W.A.	—	—	—	—	—	—	B.fl.III,341	
H. pterocarpa, F. v. M. in Transact. Vict. Inst. 126 (1855)			...	—	—	T.	V.	—	—	—	B.fl.III,342	
H. geranifolia, F. v. M. in Transact. Vict. Inst. 126 (1855)			...	—	—	—	V.	N.S.W.	—	—	B.fl.III,342	
H. medicaginoides, Turczaninow in Bull. Mosc. XXII, 27 (1849)				W.A.	S.A.	—	V.	—	—	—	B.fl.III,342	M.fr.IV,178.
H. muriculata, Turczaninow in Bull. Mosc. 28 (1849)			...	W.A.	—	—	—	—	—	—	B.fl.III,342	
H. callicarpa, Bunge in Lehm. pl. Preiss. I, 283 (1844)			...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,343	
H. scutellifera, Bentham, Fl. Austr. III, 343 (1866)			...	W.A.	—	—	—	—	—	—	B.fl.III,343	
H. hispidula, Bunge in Lehm. pl. Preiss. I, 283 (1844)			...	W.A.	—	—	—	—	—	—	B.fl.III,343	
H. trachycarpa, F. v. M. in Schlecht. Linnaea XXV, 394 (1852)				—	S.A.	—	—	N.S.W.	—	—	B.fl.III,343	
H. pilifera, Turczaninow in Bull. Mosc. XXII, 26 (1849)			...	W.A.	—	—	—	—	—	—	B.fl.III,344	
H. capillaris, F. v. M., fragm. IV, 178 (1864)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,344	M.fr.IV,178.
H. rugulosa, Turczaninow in Bull. Soc. Mosc. XXII, 27 (1849)			...	W.A.	—	—	—	—	—	—	B.fl.III,344	
H. grammatocarpa, F. v. M., fragm. II, 128 (1861)			...	...	—	—	—	—	—	N.A.	B.fl.III,344	M.fr.II,12.
H. diantha, De Candolle, prodr. IV, 63 (1830)			...	W.A.	—	—	—	—	—	—	B.fl.III,345	M.fr.IV,179.
H. lemnoides, Bentham, Fl. Austr. III, 345 (1866)			...	W.A.	—	—	—	—	—	—	B.fl.III,345	
H. alata, R. Brown in annal. des sc. phys. VI, 75, t. 61 (1820)			...	W.A.	—	—	—	—	—	—	B.fl.III,345	
H. tetragonocarpa, Bunge in Lehm. pl. Preiss. I, 284 (1844)			...	W.A.	—	—	—	—	—	—	B.fl.III,345	
H. glochidiata, Bentham, Fl. Austr. III, 346 (1866)			...	W.A.	—	—	—	—	—	—	B.fl.III,346	
H. verticillata, Turczaninow in Bull. Mosc. XXII, 28 (1849)			...	W.A.	—	—	—	—	—	—	B.fl.III,346	M.fr.II,129;VII,147.
H. Asiatica, Linné, spec. plant. 234 (1753)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,346	M.fr.VII,147.

DIDISCUS, De Candolle in Bot. Mag. t. 2857 (1828). (Dimetopia, Pritzelia, Huegelia, Cesatia, Hemicarpus.)

D. pusillus, F. v. M., fragm. IX, 47 (1875)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,348	M.fr.IX,47.
D. cyanopetalus, F. v. M., fragm. IX, 46 (1875)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,348	M.fr.I,231.
D. eriocarpus, F. v. M., fragm. IX, 46 (1875)	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.III,348	M.fr.IX,46.
D. villosus, F. v. M. in Proc. Roy. Soc. Tasm. III, 238 (1857)	...			—	—	—	—	—	—	N.A.	B.fl.III,349	
D. coeruleus, De Candolle in Hook. Bot. Mag. 2875 (1828)			...	W.A.	—	—	—	—	—	—	B.fl.III,349	M.fr.IX,47.
D. pilosus, Bentham in Hueg. enum. 54 (1837)	...	...	..	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,349	
D. glaucifolius, F. v. M. in Schlecht. Linnaea XXV, 395 (1852)				—	S.A.	—	—	N.S.W.	—	—	B.fl.III,350	M.fr.IX,47.
D. glandulosus, F. v. M. in Proc. Roy. Soc. Tasm. III, 238 (1857)				—	—	—	—	—	—	N.A.	B.fl.III,350	
D. incisus, Hooker, Bot. Magaz. 2875 (1828)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,350	
D. procumbens, F. v. M. in Proc. Roy. Soc. Tasm. III, 237 (1857)				—	—	—	—	N.S.W.	Q.	—	B.fl.III,350	M.fr.IX,47.
D. hemicarpus, F. v. M. in Trans. Bot. Soc. Edinb. VII, 491 (1857)				—	—	—	—	—	—	N.A.	B.fl.III,351	M.fr.IX,47.
D. humilis, J. Hooker in icon. pl. t. 304 (1841)	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,351	

TRACHYMENE, Rudge in Transact. Linn. Soc. X, 300 (1810). (Siebera, Fischera, Platysace, Platycarpidium.)

T. compressa, Rudge in Transact. Linn. Soc. X, 300 (1810)	...	W.A.	—	—	—	—	—	—	B.fl.III,352	
T. juncea, Bunge in Lehm. pl. Preiss. I, 286 (1844)	...	...	W.A.	—	—	—	—	—	—	B.fl.III,353	
T. haplosciadea, F. v. M.; Siebera, Benth., Fl. Austr. III, 353 (1866)			W.A.	—	—	—	—	—	—	B.fl.III,353	
T. cirrosa, F. v. M.; Platysace, Bunge in pl. Preiss. I, 285 (1844)			—	S.A.	—	V.	—	—	—	B.fl.III,354	
T. heterophylla, F. v. M., First Gen. Rep. 13 (1853)	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.III,354	
T. tenuissima, F. v. M.; Siebera, Benth., Fl. Austr. III, 354 (1866)			W.A.	—	—	—	—	—	—	B.fl.III,354	
T. dissecta, F. v. M.; Siebera, Bentham, Fl. Austr. III, 354 (1866)			W.A.	—	—	—	—	—	—	B.fl.III,354	
T. commutata, Turczaninow in Bull. Mosc. XXII, 30 (1849)	...		W.A.	—	—	—	—	—	—	B.fl.III,355	
T. effusa, Turczaninow in Bull. Mosc. XXII, 31 (1849)	...	...	W.A.	—	—	—	—	—	—	B.fl.III,355	
T. deflexa, Turczaninow in Bull. Mosc. XXII, 31 (1849)	...	...	W.A.	—	—	—	—	—	—	B.fl.III,355	
T. valida; Platycarpidium, F. v. M. in Kew. Misc. IX, 310 (1857)			—	—	—	—	—	Q.	—	B.fl.III,355	
T. ericoides, Sieber in De Candolle, prodr. IV, 73 (1830)	...		—	—	—	V.	N.S.W.	Q.	—	B.fl.III,356	
T. linearis, Sprengel, spec. umbellif. 7 (1818)	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,356	
T. Billardieri, F. v. M.; Siebera, Benth., Fl. Austr. III, 356 (1866)			—	—	—	V.	N.S.W.	—	—	B.fl.III,356	
T. Stephensonii, Turczaninow in Bull. de Mosc. XX, 170 (1847)			—	—	—	—	N.S.W.	—	—	B.fl.III,357	

XANTHOSIA, Rudge in Transact. Linn. Soc. X, 361 (1810). (Leucolaena, Schoenolaena, Pentapeltis.)

X. juncea, Bentham, Fl. Austr. III, 359 (1866)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. tenuior, Bentham, Fl. Austr. III, 359 (1866)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. peltigera, Steudel, nomencl. bot. 791 (1841)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. hederifolia, Bentham, Fl. Austr. III, 359 (1866)		...	...	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. candida, Steudel, nomencl. bot. 790 (1841)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,259	
X. tridentata, De Candolle, prodr. IV, 75 (1830)	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.III,260	
X. singuliflora, F. v. M., fragm. IV, 184 (1864)	...	...	..	W.A.	—	—	—	—	—	—	B.fl.III,260	M.fr.IV,184.
X. pilosa, Rudge in Transact. Linn. Soc. X. 301, t. 22 (1811)		...		—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,260	
X. ciliata, Hooker, icon. pl. t. 726 (1848)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,261	
X. pusilla, Bunge in Lehm. pl. Preiss. I, 291 (1844)	...	...		W.A.	S.A.	T.	V.	—	—	—	B.fl.III,261	
X. fruticulosa, Bentham, Fl. Austr. III, 361 (1866)	...	...		W.A.	—	—	—	—	—	—	B.fl.III,261	
X. Huegelii, Steudel, nomenclat. bot. 791 (1841)	...	..	...	W.A.	—	—	—	—	—	—	B.fl.III,362	
X. dissecta, J. Hooker in icon. plant. t. 302 (1841)		...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,362	
X. peduncularis, Bentham, Fl. Austr. III, 362 (1866)		...	..	W.A.	—	—	—	—	—	—	B.fl.III,362	
X. vestita, Bentham, Fl. Austr. III, 363 (1866)	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,363	
X. Atkinsoniana, F. v. M., fragm. II, 127 (1861)	...	...	...	W.A.	—	—	—	N.S.W.	—	—	B.fl.III,363	M.fr.II,127.
X. rotundifolia, De Candolle, prodr. IV, 75 (1830)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,363	

AZORELLA, Lamarck, Encycl. méthodiq. I, 344, t. 189 (1783). (Fragosa, Pozoa, Microsciadium, Schizeilema, Oschatzia, Centella partly, Dichopetalum.)

A. Muelleri, Bentham, Fl. Austr. III, 364 (1866)	—	—	—	V.	—	—	—	B.fl.III,364				
A. cuneifolia, F. v. M. in Transact. phil. Inst. Vict. I, 103 (1855)	—	—	—	V.	—	—	—	B.fl.III,365				
A. saxifraga, Bentham, Fl. Austr. III, 365 (1866)	—	—	T.	—	—	—	—	B.fl.III,365				
A. dichopetala, Bentham, Fl. Austr. III, 365 (1866)	—	—	T.	V.	—	—	—	B.fl.III,365				

HUANACA, Cavanilles, icon. VI, 18, t. 528 (1801). (Diplaspis, Pozoopsis.)

H. hydrocotylea, Bentham & J. Hooker, gen. pl. I, 877 (1867) ...	—	—	T.	V.	—	—	—	B.fl.III,366				
H. cordifolia, Bentham & J. Hooker, gen. pl. I, 877 (1867) ...	—	—	T.	—	—	—	—	B.fl.III,366				

ACTINOTUS, Labillardière, Nov. Holl. pl. spec. I, 67, t. 92 (1804). (Hemiphues, Eriocalia, Holotome.)

A. Helianthi, Labillardière, Nov. Holl. pl. spec. I, 67, t. 92 (1804)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,367				
A. leucocephalus, Bentham in Hueg. enum. 56 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,368				
A. minor, De Candolle, prodr. IV, 83 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.III,368				
A. Gibbonsii, F. v. M., fragm. VI, 23 (1867)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,23.			
A. omnifertilis, Bentham, Fl. Austr. III, 368 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,368				
A. rhomboideus, Bentham, Fl. Austr. III, 368 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,368				
A. bellidioides, Bentham, Fl. Austr. III, 369 (1866)	—	—	T.	—	—	—	—	B.fl.III,369				
A. glomeratus, Bentham, Fl. Austr. III, 369 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,369				

ERYNGIUM, Tournefort, inst. 327, t. 173 (1700) from Theophrastos and Dioscorides).

E. rostratum, Cavanilles, icon. pl. VI, 35, t. 552 (1801)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,370				
E. vesiculosum, Labillardière, Nov. Holl. pl. spec. I, 73, t. 98 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,370	M.fr.VII,148.			
E. plantagineum, F. v. M. in Pap. Roy. Soc. Tasm. III, 235 (1857)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,371	M.fr.VII,148.			
E. expansum, F. v. M. in Pap. Roy. Soc. Tasm. III, 236 (1857)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,371	M.fr.VII,148.			

APIUM, Tournefort, inst. 305, t. 160 (1700).

A. prostratum, Labillardière, relat. du voy. I, 141 (1799) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,372	M.fr.VII,148.			
A. leptophyllum, F. v. M. in Benth. Fl. Austr. III, 372 (1866) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,372	M.fr.VII,148.			

SIUM, Tournefort, inst. 308, t. 162 (1700), from C. Bauhin (1623).

S. latifolium, C. Bauhin, pinax 154 (1623)	—	S.A.	—	V.	N.S.W.	Q.	—	—				
S. angustifolium, Linné, spec. plant. ed. sec. 1672 (1763) ...	—	—	—	—	N.S.W.	—	—	—				

SESELI, Rivinus in Rupp. Fl. Jen. 267 (1718).

S. Harveyanum, F. v. M. in Transact. phil. Soc. Vict. I, 104 (1855)	—	—	—	V.	—	—	—	B.fl.III,373				
S. algens, F. v. M. in Transact. phil. Soc. Vict. I, 104 (1855) ...	—	—	—	V.	—	—	—	B.fl.III,373				

CRANTZIA, Nuttall, Gen. of North Amer. plants I, 177 (1818). (Non Scopoli, 1777, Crantziola, F. v. M.)

C. lineata, Nuttall, Gen. of North Amer. pl. I, 178 (1818) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,374				

OENANTHE, Tournefort, inst. 312, t. 166 (1700).

O. Javanica, De Candolle, prodr. IV, 138 (1830)	—	—	—	—	—	Q.	—	—	M.fr.V,182.			

ACIPHYLLA, R. & G. Forster, char. gen. 135, t. 68 (1776). (Anisotome, Calosciadium, Gingidium partly, Ligustici subgenus.)

A. simplicifolia, F. v. M. in Benth. Fl. Austr. III, 375 (1866) ...	—	—	—	V.	N.S.W.	—	—	B.fl.III,375				
A. glacialis, F. v. M. in Benth. Fl. Austr. III, 375 (1866) ...	—	—	—	V.	N.S.W.	—	—	B.fl.III,375				
A. procumbens, F. v. M. in Benth. Fl. Austr. III, 375 (1866) ...	—	—	T.	—	—	—	—	B.fl.III,375				

DAUCUS, Tournefort, inst. 307, t. 161 (1700), from l'Ecluse (1576).

D. brachiatus, Sieber in De Cand., prodr. IV, 214 (1830) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,376				

OREOMYRRHIS, Endlicher, gen. plantar. 787 (1839).

O. andicola, Endlicher, gen. plantar. 787 (1839)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,377	M.fr.VIII,185.			
O. pulvinifica, F. v. M., fragm. VIII, 185 (1874)	—	—	—	V.	N.S.W.	—	—	—	M.fr.VIII,185.			

SYNPETALEAE PERIGYNAE.

F. v. M. in Woolls's plants of the neighb. of Sydney, 27 (1880).

AQUIFOLIACEAE.

De Candolle, Théor. élém. I, 217 (1813).

BYRONIA, Endlicher in den Ann. des Wien. Mus. I, 184 (1836).

B. Arnhemensis, F. v. M., fragm. II, 119 (1861)	—	—	—	—	—	—	N.A.	B.fl.I,397	M.fr.II,119.			

ILEX, Linné, gen. plant. 33 (1737), from C. Bauhin (1623).

I. peduncularis, F. v. M., fragm. VII, 105 (1871)	—	—	—	—	—	Q.	—	—	M.fr.VII,105.			

OLACINEAE.

Mirbel in Bull. de la Soc. philomat. 377 (1813).

XIMENIA, Plumier, nova plant. Amer. genera 6, t. 21 (1703).

X. Americana, Linné, spec. plant. 1193 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,391	M.fr.VI,3;IX,150.			

OLAX, Linné, amoen. acad. 387 (1747). (Spermaxyrum, Lopadocalyx.)

O. phyllanthi, R. Brown, prodr. 358 (1810)	W.A.	—	—	—	—	—	—	B.fl.I,392	M.fr.IX,151.			
O. retusa, F. v. M. in Benth. Fl. Austr. I, 392 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,392	M.fr.IX,151.			
O. stricta, R. Brown, prodr. 358 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,392	M.fr.IX,151.			
O. Benthamiana, Miquel in Lehm. pl. Preiss. I, 228 (1844) ...	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,393	M.fr.IX,151.			
O. aphylla, R. Brown, prodr. 358 (1810)	—	—	—	—	—	—	N.A.	B.fl.I,393	M.fr.IX,151.			

CANSJERA, A. L. de Jussieu, gen. pl. 448 (1789).

C. leptostachya, Benth. in Hook. Lond. Journ. of Bot. II, 231 (1843)	—	—	—	—	—	Q.	—	B.fl.I,394	M.fr.VI,3.			

OPILIA, Roxburgh, pl. Corom. II, 31, t. 158 (1799).

O. amentacea, Roxburgh, pl. Corom. II, 31, t. 158 (1799) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,394	M.fr.VI,4;IX,150.			

PHLEBOCALYMMA, Griffith in Benth. & J. Hook. gen. pl. I, 353 (1862). (Phlebocalymna).

P. lobospora, F. v. M., fragm. IX, 151 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,151.			

PENNANTIA, R. & G. Forster, char. gen. 133, t. 67 (1776).												
P. Cunninghamii, Miers in Ann. of nat. hist. sec. ser. IX, 491 (1852)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,395	M.fr.VI,4;IX,150.
P. Endlicheri, Reisseck in Schlecht. Linnaea XVI, 341, t. 13 (1843)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,150.
APODYTES, E. Meyer in Hook. Journ. of Bot. III, 155 (1841).									
A. brachystylis, F. v. M., fragm. IX, 149 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,149.
VILLARESIA, Ruiz & Pavon, flor. Peruv. et Chil. prodr. 35 (1794).									
V. Moorei, F. v. M. in Benth. Fl. Austr. I, 396 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.I,395	
V. Smythii, F. v. M., fragm. V, 156 (1866)	—	—	—	—	—	Q.	—	—	M.fr.IX,150.
GOMPHANDRA, Wallich, Catal. n. 3718 & 7204 (1832).									
G. Australiana, F. v. M., fragm. VI, 3 (1877)	—	—	—	—	—	Q.	—	—	M.fr.VI,3&253;IX,150.

ELAEAGNEAE.

R. Brown, prodr. I, 350 (1810.)

ELAEAGNUS, Tournefort, coroll. 53, t. 459 (1703) from Camerarius (1586).												
E. latifolia, Linné, spec. plant. 121 (1753)	—	—	—	—	—	Q.	—	M.fr.VI,39	M.fr.IX,190.

SANTALACEAE.

R. Brown, prodr. I, 350 (1810).

THESIUM, Linné, gen. pl. 60 (1737).												
T. australe, R. Brown, prodr. 353 (1810)	—	—	T.	V.	N.S.W.	Q.	—	M.fr.VI,212	M.fr.IX,3.
SANTALUM, Linné, gen. ed. sec., 165 (1742), from C. Bauhin (1623). (Fusanus, Eucarya).									
S. lanceolatum, R. Brown, prodr. 356 (1810)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VI,214	M.fr.I,85;VIII,11;IX,3.
S. ovatum, R. Brown, prodr. 355 (1810)	—	—	—	—	—	—	N.A.	B.fl.VI,214	
S. obtusifolium, R. Brown, prodr, 356 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,215	M.fr.VIII,11.
S. acuminatum, A. de Candolle, prodr. XIV, 684 (1856)... ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,216	M.fr.IX,169;XI,20.
S. persicarium, F. v. M. in Transact. Vict. Inst. 41 (1854) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,216	M.fr.I,86;IX,3;XI,20.
S. cygnorum, Miquel in Lehm. pl. Preiss. I, 615 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,217	M.fr.VIII,11.
S. crassifolium, A. de Candolle, prodr. XIV, 685 (1856)... ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,217	
CHORETRUM, R. Brown, prodr. 354 (1810).									
C. glomeratum, R. Brown, prodr. 354 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,218	
C. chrysanthum, F. v. M. in Transact. phil. Soc. Vict. I, 23 (1854)	—	S.A.	—	V.	N.S.W.	—	—	—	M.fr.VIII,11;XI,138.
C. spicatum, F. v. M., fragm. I, 21 (1858)...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,218	M.fr.VIII,11.
C. lateriflorum, R. Brown, prodr. 354 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,219	
C. Candollei, F. v. M. in Bentham's Fl. Austr. VII, 219 (1873)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,219	
LEPTOMERIA, R. Brown, prodr. 353 (1810).									
L. spinosa, A. de Candolle, prodr. XIV, 678 (1856)	W.A.	—	—	—	—	—	—	B.fl.VI,220	M.fr.VIII,10.
L. Preissiana, A. de Candolle, prodr. XIV, 678 (1856)	W.A.	—	—	—	—	—	—	B.fl.VI,221	M.fr.VIII,10.
L. pauciflora, R. Brown, prodr. 354 (1810)...	W.A.	—	—	—	—	—	—	B.fl.VI,221	M.fr.VIII,10.
L. scrobiculata, R. Brown, prodr. 354 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,221	M.fr.VIII,10.
L. acida, R. Brown, prodr. 353 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,222	M.fr.VIII,10.
L. Billardieri, R. Brown, prodr. 354 (1810)	—	—	T.	—	N.S.W.	Q.	—	B.fl.VI,222	M.fr.VIII,10.
L. aphylla, R. Brown, prodr. 354 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,222	M.fr.VIII,10.
L. glomerata, F. v. M. in J. Hook. Tasm. II, 370 (1860)... ...	—	—	T.	—	—	—	—	B.fl.VI,223	M.fr.VIII,10.
L. squarrulosa, R. Brown, prodr. 354 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,223	M.fr.VIII,11.
L. Cunninghamii, Miquel in Lehm. pl. Preiss. I, 611 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.VI,223	M.fr.VIII,10.
L. empetriformis, Miquel in Lehm. pl. Preiss. I, 610 (1845)' ...	W.A.	—	—	—	—	—	—	B.fl.VI,224	
L. axillaris, R. Brown, prodr. 354 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,224	
L. laxa, Miquel in Lehm. pl. Preiss. I, 612 (1845)...	W.A.	—	—	—	—	—	—	B.fl.VI,224	
L. obovata, Miquel in Lehm. pl. Preiss. I, 613 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,224	
OMPHACOMERIA, Endlicher, gen. pl. 326 (1838).									
O. acerba, A. de Candolle, prodr. XIV, 681 ((1856)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,224	
ANTHOBOLUS, R. Brown, prodr. 357 (1810).									
A. filifolius, R. Brown, prodr. 357 (1810)	—	—	—	—	—	—	N.A.	B.fl.VI,226	M.fr.IX,3.
A. triqueter, R. Brown, prodr. 357 (1810)	—	—	—	—	—	Q.	—	B.fl.VI,226	
A. foveolatus, F. v. M., fragm. I, 212 (1859)	W.A.	—	—	—	—	—	—	B.fl.VI,226	M.fr.II,181;IX,3;XI,20.
A. leptomerioides, F. v. M., fragm. I, 21 (1858)	—	—	—	—	—	Q.	—	B.fl.VI,227	M.fr.IX,3.
A. exocarpoides, F. v. M., fragm. IX, 3 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,3.
EXOCARPOS, Labillardière, Relat. du voy. à la rech. de La Pérouse I, 155 (1798). (Exocarpus.)									
E. latifolia, R. Brown, prodr. 356 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,228	M.fr.VIII,10.
E. odorata, A. de Candolle, prodr. XIV, 689 (1856)	W.A.	—	—	—	—	—	—	B.fl.VI,228	M.fr.VIII,10.
E. cupressiformis, Labillardière, voy. I, 155, t. 14 (1798) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,229	
E. spartea, R. Brown, prodr. 356 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,229	M.fr.VIII,10;XI,20. [20.
E. aphylla, R. Brown, prodr. 357 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,230	M.fr.VIII,10;IX,169;XI,
E. homaloclada, C. Moore & F. v. M., fragm. VIII, 9 (1871) ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,230	M.fr.VIII,9.
E. phyllanthoides, Endlicher, prodr. pl. ins. Norf. 46 (1833) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.
E. stricta, R. Brown, prodr. 357 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,230	M.fr.IX,169.
E. humifusa, R. Brown, prodr. 356 (1810)	—	—	T.	—	—	—	—	B.fl.VI,231	
E. nana, J. Hooker in Lond. Journ. of Bot. VI, 281 (1847) ...	—	—	T.	V.	—	—	-	B.fl.VI,230	

LORANTHACEAE.

A. L. de Jussieu, in Ann. du Mus. XII, 292 (1808).

VISCUM, Tournefort, inst. 609, t. 380 (1700), from Camerarius (1586).												
V. orientale, Willdenow, spec. plant. IV, 737 (1805)	—	—	—	—	—	Q.	—	B.fl.III,396	
V. angulatum, Heyne in De Cand., prodr. IV, 283 (1830) ...	—	—	—	—	—	Q.	—	B.fl.III,396	
V. articulatum, N. L. Burmann, Fl. Ind. 311 (1768)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,396	

NOTOTHIXOS, Oliver in Journ. Linn. Soc. VII, 104 (1865).

N. incanus, Oliver in Journ. Linn. Soc. VII, 104 (1865)... ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,397	M.fr.II,109,176;IV,173.

LORANTHUS, Linné, syst. nat. ed. sec. 22 (1740).

L. celastroides, Sieber in Roem. & Schult. syst. veg. VII, 163 (1829)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,389	M.fr.VI,252.
L. Bidwillii, Bentham, Fl. Austr. III, 390 (1866)...	—	—	—	—	—	Q.	—	B.fl.III,390	
L. myrtifolius, Cunningham in Benth. Fl. Austr. III, 390 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,390	
L. longiflorus, Desrousseaux in Lam. Encycl. méth. IV, 598 (1796)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,390	
L. angustifolius, R. Brown in Benth. Fl. Austr. III, 390 (1866)	—	S.A.	—	—	—	—	—	B.fl.III,390	
L. dictyophlebus, F. v. M., Rep. Burdek. Exped. 14 (1858) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,391	
L. alyxifolius, F. v. M. in Bentham Fl. Austr. III, 391 (1866) ..	—	—	—	—	N.S.W.	Q.	—	B.fl.III,391	
L. odontocalyx, F. v. M. in Bentham Fl. Austr. III, 391 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,391	
L. linearifolius, Hooker in Mitchell, Trop. Austr. 102 (1848) ...	W.A.	S.A.	—	—	—	Q.	—	B.fl.III,391	
L. Exocarpi, Behr in Schlecht. Linnaea XX, 624 (1847)... ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,392	
L. acacioides, Cunningham in Benth. Fl. Austr. III, 392 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,392	
L. signatus, F. v. M. in Bentham Fl. Austr. III, 392 (1866) ...	—	—	—	—	—	Q.	N.A.	B.fl.III,392	
L. maytenifolius, A. Gray, Bot. Wilk. Expl. Exped. I, 739, t. 99 (1854)	—	—	—	—	N.S.W.	—	—	B.fl.III,393	
L. sanguineus, F. v. M., fragm. I, 177 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.III,393	M.fr.IV,171.
L. bifurcatus, Bentham, Fl. Austr. III, 393 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,393	
L. linophyllus, Fenzl in Huegel. enum. 56 (1837)...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,393	
L. pendulus, Sieber in De Candolle, prodr. IV, 294 (1830) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,394	
L. Quandang, Lindley in Mitch. Three Exped. II, 69 (1838) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,395	
L. grandibracteus, F. v. M., Rep. Burdek. Exped. 14 (1858) ...	—	S.A.	—	—	—	Q.	N.A.	B.fl.III,395	

ATKINSONIA, F. v. Mueller, fragm. V, 34 (1865).

A. ligustrina, F. v. M., fragm. V, 34 (1865)	—	—	—	—	N.S.W.	—	—	B.fl.III,388	M.fr.II,130;VI,252.

NUYTSIA, R. Brown in Journ. of the geogr. Soc. I, 17 (1831).

N. floribunda, R. Brown in Journ. geogr. Soc. I, 17 (1831) ...	W.A.	—	—	—	—	—	—	B.fl.III,387	M.fr.VI,252.

PROTEACEAE.

A. L. de Jussieu, gen. pl. 78 (1789).

PETROPHILA, R. Brown in Transact. Linn. Soc. X, 67 (1809).

P. teretifolia, R. Brown in Transact. Linn. Soc. X, 68 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,321	M.fr.VI,245.
P. longifolia, R. Brown, prot. nov. 5 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,322	M.fr.VI,255.
P. media, R. Brown, prot. nov. 5 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,322	M.fr.VI,255.
P. acicularis, R. Brown in Transact. Linn. Soc. X, 69 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,323	M.fr.VI,245.
P. megalostegia, F. v. M., fragm. X, 61 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,61.
P. linearis, R. Brown, prot. nov. 6 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,323	M.fr.VI,243.
P. anceps, R. Brown, prot. nov. 5 (1830)	W.A.	—	—	—	—	—	—	—	M.fr.VI,243;X,47.
P. heterophylla, Lindley, Bot. Regist. XXV, App. XXXV (1839)	W.A.	—	—	—	—	—	—	B.fl.V,324	M.fr.VI,244.
P. biloba, R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,324	M.fr.VI,255.
P. propinqua, R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,325	
P. squamata, R. Brown in Transact. Linn. Soc. X, 70 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,325	M fr.VI,244.
P. colorata, Meissner in Lehm. pl. Preiss. II, 246 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,326	
P. striata, R. Brown, prot. nov. 6 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,326	M.fr.VI,244.
P. divaricata, R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,326	M.fr.VI,344.
P. Serruriae, R. Brown, prot. nov. 6 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,327	M.fr.VI,243.
P. inconspicua, Meissner in Hook. Kew Misc. VII, 68 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,327	M.fr.VI,243.
P. trifida, R. Brown in Transact. Linn. Soc. X, 70 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,328	M.fr.VI,244.
P. carduacea, Meissner in Hook. Kew Misc. IV, 182 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,328	M.fr.VI,244.
P. Shuttleworthiana, Meissner in Lehm. pl. Preiss. II, 246 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,329	M.fr.VI,245.
P. macrostachya, R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,329	M.fr.VI,245.
P. diversifolia, R. Brown in Transact. Linn. Soc. X, 70 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,329	M.fr.VI,244.
P. biternata, Meissner in Hook. Kew Misc. VII, 69 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,330	M.fr.VI,245.
P. plumosa, Meissner in Hook. Kew Misc. VII, 69 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,330	M.fr.VI,244.
P. ericifolia, R. Brown, prot. nov. 5 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,331	M.fr.VI,243.
P. chrysantha, Meissner in Hook. Kew Misc. VII, 68 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,331	M.fr.VI,243.
P. pedunculata, R. Brown in Transact. Linn. Soc. X, 70 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,332	M.fr.VI,242.
P. pulchella, R. Brown in Transact. Linn. Soc. X, 69 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,332	M.fr.VI,242.
P. sessilis, Sieber in Roem. & Schult. mant. 262 (1822)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,332	
P. fastigiata, R. Brown in Transact. Linn. Soc. X, 70 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,333	M.fr.VI,245.
P. seminuda, Lindley, Bot. Regist. XXV, App. XXXIV (1839)	W.A.	—	—	—	—	—	—	B.fl.V,333	M.fr.VI,243.
P. circinata, Kippist in Hook. Kew Misc. VII, 67 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,333	M.fr.VI,244.
P. Drummondii, Meissner in Lehm. pl. Preiss. I, 496 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,334	M.fr.VI,243.
P. crispata, R. Brown, prot. nov. 6 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,334	
P. rigida, R. Brown in Transact. Linn. Soc. X, 69 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,334	M.fr.VI,245.
P. multisecta, F. v. M., fragm. VI, 242 (1868)	—	S.A.	—	—	—	—	—	B.fl.V,335	M.fr.VI,242.
P. conifera, Meissner in Hook. Kew Misc. VII, 67 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,335	M.fr.VI,245.
P. semifurcata, F. v. M. in Bentham Fl. Austr. V, 335 (1870) ...	W.A.	—	—	—	—	—	—	B.fl.V,335	M.fr.X,47.

ISOPOGON, R. Brown in Transact. Linn. Soc. X, 70 (1809).

I. latifolius, R. Brown, prot. nov. 8 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,338	M.fr.VI,237.
I. cuneatus, R. Brown in Transact. Linn. Soc. X, 73 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,339	M.fr.VI,238.
I. linearis, Meissner in Hook. Kew Misc. VII, 69 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,339	M.fr.VI,236.
I. polycephalus, R. Brown in Transact. Linn. Soc. X, 73 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,339	M.fr.VI,236.
I. attenuatus, R. Brown in Transact. Linn. Soc. X, 73 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,340	M.fr.VI,237.
I. sphaerocephalus, Lindl., Bot. Regist. XXV, App. XXXIV (1839)	W.A.	—	—	—	—	—	—	B.fl.V,340	M.fr.VI,236.
I. uncinatus, R. Brown, prot. nov. 8 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,341	M.fr.VI,237.
I. buxifolius, R. Brown in Transact. Linn. Soc. X, 74 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,341	M.fr.VI,237.
I. axillaris, R. Brown in Transact. Linn. Soc. X, 74 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,341	M.fr.VI,237.
I. tridens, F. v. M., fragm. VI, 239 (1868)...	W.A.	—	—	—	—	—	—	B.fl.V,342	M.fr.VI,239.

I. Baxteri, R. Brown, prot. nov. 9 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,342	M.fr.VI,240.
I. roseus, Lindley, Bot. Regist. Misc. n. 37 (1842)	W.A.	—	—	—	—	—	—	B.fl.V,343	M.fr.VI,240
I. adenanthoides, Meissner in Hook. Kew Misc. VII, 69 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,343	M.fr.VI,241.
I. trilobus, R. Brown in Transact. Linn. Soc. X, 72 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,343	M.fr.VI,239.
I. tripartitus, R. Brown, prot. nov. 8 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,344	
I. longifolius, R. Brown in Transact. Linn. Soc. X, 73 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,344	M.fr.VI,237.
I. Drummondii, Bentham, Fl. Austr. V, 344 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,344	M.fr.VI,241.
I. heterophyllus, Meissner in Lehm. pl. Preiss. I, 504 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,345	M.fr.VI,241.
I. villosus, Meissner in De Candolle, prodr. XIV, 277 ( ) ...	—	—	—	—	—	—	—	—	M.fr.VI,241.
I. teretifolius, R. Brown in Transact. Linn. Soc. X, 71 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,345	M.fr.VI,241.
I. anethifolius, R. Brown in J. Knight, proteac. 94 (1809) ...	W.A.	—	—	—	N.S.W.	—	—	B.fl.V,346	M.fr.VI,238.
I. petiolaris, Cunningham in R. Br., prot. nov. 8 (1830)... ...	—	—	—	—	N.S.W.	—	—	B.fl.V,346	
I. anemonifolius, R. Brown in J. Knight, prot. 93 (1809) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,347	M.fr.VI,238;X,90.
I. ceratophyllus, R. Brown in Transact. Linn. Soc. X, 72 (1809)	—	S.A.	T.	—	N.S.W.	—	—	B.fl.V,347	M.fr.VI,238.
I. asper, R. Brown, prot. nov. 8 (1830)	W.A.	S.A.	T.	V.	—	—	—	B.fl.V,348	M.fr.VI,240.
I. crithmifolius, F. v. M., fragm. VI, 239 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,348	M.fr.VI,239.
I. formosus, R. Brown in Transact. Linn. Soc. X, 72 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,349	M.fr.VI,240.
I. divergens, R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,349	M.fr.VI,241.
I. scabriusculus, Meissner in Hook. Kew Misc. IV, 182 (1852)...	W.A.	—	—	—	—	—	—	B.fl.V,349	M.fr.VI,240.
ADENANTHOS, Labillardière, Nov. Holl. pl. specim. I, 28, t. 36 (1804).									
A. Detmoldi, F. v. M., fragm. VIII, 149 (1874)	W.A.	—	—	—	—	—	—	—	M.fr.VIII,149.
A. barbigerus, Lindley, Bot. Regist. XXV, App. XXXVI (1839)	W.A.	—	—	—	—	—	—	B.fl.V,351	M.fr.VI,205;VIII,149
A. obovatus, Labillardière, Nov. Holl. pl. spec. I, 29, t. 37 (1804)	W.A.	—	—	—	—	—	—	B.fl.V,352	
A. cuneatus, Labillardière, Nov. Holl. pl. spec. I, 28, t. 36 (1804)	W.A.	—	—	—	—	—	—	B.fl.V,352	
A. Cunninghami, Meissner in Lehm. pl. Preiss. I, 513 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,352	
A. pungens, Meissner in Lehm. pl. Preiss. I, 515 (1845)... ...	W.A.	—	—	—	—	—	—	B.fl.V,353	
A. venosus, Meissner in Hook. Kew Misc. IV, 183 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,353	
A. Dobsoni, F. v. M., fragm. VI, 204 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,353	M.fr.VI,204.
A. linearis, Meissner in Hook. Kew Misc. IV, 183 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,354	
A. sericeus, Labillardière, Nov. Holl. pl. specim. I, 29, t. 38 (1804)	W.A.	S.A.	—	—	—	—	—	B.fl.V,354	
A. Meissneri, Lehmann, pl. Preiss. I, 515 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,354	
A. filifolius, Bentham, Fl. Austr. V, 355 (1873)	W.A.	—	—	—	—	—	—	B.fl.V,355	
A. terminalis, R. Brown in Transact. Linn. Soc. X, 152 (1809)...	—	S.A.	—	V.	—	—	—	B.fl.V,355	
A. flavidiflorus, F. v. M., fragm. I, 157 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,355	M.fr.I,157.
A. apiculatus, R. Brown, prot. nov. 9 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,356	
SIMSIA, R. Brown in Transact. Linn. Soc. X, 152 (1809). (Stirlingia.)									
S. simplex, F.v.M.; Stirl., Lindl. Bot. Reg. XXV, App. XXX (1839)	W.A.	—	—	—	—	—	—	B.fl.V,357	
S. abrotanoides, F.v.M.; Stirlingia, Meissn. in pl. Preiss. I, 517 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,357	
S. teretifolia, F. v. M.; Stirlingia, Meissn. in pl. Preiss. 515 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,357	
S. tenuifolia, R. Brown in Transact. Linn. Soc. X, 152 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,358	
S. latifolia, R. Brown, prot. nov. 9 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,358	
SYNAPHEA, R. Brown in Transact. Linn. Soc. X, 155 (1809).									
S. polymorpha, R. Brown in Transact. Linn. Soc. X, 156 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,360	
S. dilatata, R. Brown in Transact. Linn. Soc. X, 156 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,360	
S. favosa, R. Brown in Transact. Linn. Soc. X, 156 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,360	
S. Preissii, Meissner in Lehm. pl. Preiss. I, 529 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,361	
S. acutiloba, Meissner in Lehm. pl. Preiss. I, 528 (1845)... ...	W.A.	—	—	—	—	—	—	B.fl.V,361	
S. petiolaris, R. Brown in Transact. Linn. Soc. X, 156 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,361	
S. decorticans, Lindley, Bot. Regist. XXV, App. XXXII (1839)	W.A.	—	—	—	—	—	—	B.fl.V,362	
S. pinnata, Lindley, Bot. Regist. XXV, App. XXXII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.V,362	
CONOSPERMUM, Smith in Transact. Linn. Soc. IV, 213 (1798).									
C. capitatum, R. Brown in Transact. Linn. Soc. X, 155 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,365	
C. petiolare, R. Brown, prot. nov. 11 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,365	
C. teretifolium, R. Brown in Transact. Linn. Soc. X, 155 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,365	M.fr.VI,224.
C. flexuosum, R. Brown, prot. nov. 11 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,366	M.fr.VI,224.
C. acerosum, Lindley, Bot. Regist. XXV, App. XXX (1839) ...	W.A.	—	—	—	—	—	—	B.fl.V,366	
C. amoenum, Meissner in Lehm. pl. Preiss. I, 522 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,366	M.fr.I,157;VI,224.
C. nervosum, Meissner in Hook. Kew Misc. VII, 71 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,367	M.fr.VI,224.
C. diffusum, Bentham, Fl. Austr. V, 367 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,367	
C. glumaceum, Lindley, Bot. Regist. XXV, App. XXX (1839)...	W.A.	—	—	—	—	—	—	B.fl.V,367	
C. ephedroides, Kippist in Hook. Kew Misc. VII, 70 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,368	
C. Toddii, F. v. M., fragm. X, 20 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,20,90.
C. polycephalum, Meissner in Lehm. pl. Preiss. II, 249 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,368	
C. coeruleum, R. Brown in Transact. Linn. Soc. X, 154 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,368	M.fr.VI,224.
C. debile, Kippist in Hook. Kew Misc. VII, 70 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,369	
C. scaposum, Bentham, Fl. Austr. V, 369 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,369	
C. Huegelii, R. Brown in Endlicher, nov. stirp. dec. 58 (1839)...	W.A.	—	—	—	—	—	—	B.fl.V,369	
C. densiflorum, Lindley, Bot. Regist. XXV, App. XXXII (1839)	W.A.	—	—	—	—	—	—	B.fl.V,369	
C. Brownii, Meissner in Lehm. pl. Preiss. II, 248 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,370	
C. longifolium, Smith, Exot. Bot. II, 45, t. 82 (1805)	—	—	—	—	N.S.W.	—	—	B.fl.V,370	
C. tenuifolium, R. Brown in Transact. Linn. Soc. X, 154 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,370	M.fr.VI,223.
C. Mitchellii, Meissner in De Candolle, prodr. XIV, 320 (1856)	—	—	—	V.	—	—	—	B.fl.V,371	
C. sphacelatum, Hooker in Mitchell, Trop. Austr. 342 (1848) ...	—	—	—	—	—	Q.	N.A.	B.fl.V,371	M.fr.VI,223.
C. patens, Schlechtendal, Linnaea XX, 587 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,371	M.fr.VI,223.
C. taxifolium, Smith in Rees's Cycl. IX (1808)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.V,372	M.fr.VI,224.
C. ericifolium, Smith in Rees's Cycl. IX (1808)	—	—	—	—	N.S.W.	—	—	B.fl.V,372	
C. ellipticum, Smith in Rees's Cycl. IX (1808)	—	—	—	—	N.S.W.	—	—	B.fl.V,372	
C. distichum, R. Brown in Transact. Linn. Soc. X, 155 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,373	M.fr.VI,224.
C. floribundum, Bentham, Fl. Austr. V, 373 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,373	

C. incurvum, Lindley, Bot. Regist. XXV, App. XXX (1839) ...	W.A.	—	—	—	—	—	—	B.fl.V,373	
C. brachyphyllum, Lindley, Bot. Regist. XXV, App. XXXI (1839)	W.A.	—	—	—	—	—	—	B.fl.V,374	
C. stoechadis, Endlicher in Ann. Wien. Mus. II, 208 (1838) ...	W.A.	—	—	—	N.S.W.	—	—	B.fl.V,374	M.fr.XI,20.
C. triplinervium, R. Brown, prot. nov. 11 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,375	
C. bracteosum, Meissner in Lehm. pl. Preiss. I, 518 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,375	M.fr.VI,224.
C. crassinervium, Meissner in Hook. Kew Misc. IV, 184 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,375	
FRANKLANDIA, R. Brown in Transact. Linn. Soc. X, 157 (1809).									
F. fucifolia, R. Brown in Transact. Linn. Soc. X, 157 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,376	M.fr.VI,223;VII,133.
F. triaristata, Bentham, Fl. Austr. V, 377 (1870)...	W.A.	—	—	—	—	—	—	B.fl.V,377	
SYMPHYONEMA, R. Brown in Transact. Linn. Soc. X, 157 (1809). (Symphionema.)									
S. montanum, R. Brown in Transact. Linn. Soc. X, 158 (1809)...	—	—	—	—	N.S.W.	—	—	B.fl.V,377	M.fr.VI,223.
S. paludosum, R. Brown in Transact. Linn. Soc. X, 158 (1809)...	—	—	—	—	N.S.W.	—	—	B.fl.V,378	M.fr.VI,223.
BELLENDENA, R. Brown in Transact. Linn. Soc. X, 156 (1809).									
B. montana, R. Brown in Transact. Linn. Soc. X, 166 (1809) ...	—	—	T.	—	—	—	—	B.fl.V,378	
AGASTACHYS, R. Brown in Transact. Linn. Soc. X, 158 (1809).									
A. odorata, R. Brown in Transact. Linn. Soc. X, 158 (1809) ...	—	—	T.	—	—	—	—	B.fl.V,379	M.fr.X,90.
CENARRHENES, Labillardière, Nov. Holl. pl. spec. I, 36, t. 50 (1804).									
C. nitida, Labillardière, Nov. Holl. pl. spec. I, 36, t. 50 (1804)...	—	—	T.	—	—	—	—	B.fl.V,380	
PERSOONIA, Smith in Transact. Linn. Soc. IV, 215 (1798). (Linkia, 1797, Pentadactylon.)									
P. hakeiformis, Meissner in Hook. Kew Misc. IV, 185 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,383	M.fr.VI,223.
P. teretifolia, R. Brown in Transact. Linn. Soc. X, 160 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,383	M.fr.VI,223.
P. saccata, R. Brown, prot. nov. 12 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,384	M.fr.VI,223.
P. Saundersiana, Kippist in Hook. Kew Misc. VII, 72 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,384	
P. diadena, F. v. M., fragm. X, 46 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,46.
P. comata, Meissner in Hooker's Kew Misc. VII, 71 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,385	
P. brachystylis, F. v. M., fragm. VI, 221 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,385	
P. falcata, R. Brown in Transact. Linn. Soc. X, 162 (1809) ...	—	—	—	—	—	Q.	N.A.	B.fl.V,385	M.fr.VI,222.
P. trinervis, Meissner in Hook. Kew Misc. IV, 185 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,386	
P. tortifolia, Meissner in Hook. Kew Misc. IV, 185 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,386	
P. angustiflora, Bentham, Fl. Austr. V, 386 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,386	
P. rudis, Meissner in Hook. Kew Misc. IV, 185 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,387	
P. microcarpa, R. Brown in Transact. Linn. Soc. X, 160 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,387	M.fr.VI,222.
P. sulcata, Meissner in Hook. Kew Misc. IV, 185 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,387	
P. acicularis, F. v. M., fragm. VI, 220 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,388	M.fr.VI,220.
P. scabrella, Meissner in Hooker's Kew Misc. VII, 72 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,388	
P. dillwynioides, Meissner in Hook. Kew Misc. IV, 185 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,388	M.fr.VI,222.
P. striata, R. Brown, prot. nov. 13 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,388	M.fr.VI,223.
P. quinquenervis, Hooker, icon. plant. t. 425 (1842)	W.A.	—	—	—	—	—	—	B.fl.V,389	
P. rufiflora, Meissner in Hooker's Kew Misc. VII, 72 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,389	
P. scabra, R. Brown in Transact. Linn. Soc. X, 162 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,389	
P. graminea, R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,390	M.fr.VI,222.
P. longifolia, R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,390	M.fr.VI,222.
P. articulata, R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,390	
P. elliptica, R. Brown in Transact. Linn. Soc. X, 164 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,391	M.fr.VI,222.
P. ferruginea, Smith, Exot. Bot. II, 47, t. 83 (1805)	—	—	—	—	N.S.W.	—	—	B.fl.V,391	
P. media, R. Brown, prot. nov. 16 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,391	
P. cornifolia, Cunningham in R. Br., prot. nov. 16 (1830) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,392	
P. marginata, R. Brown, prot. nov. 16 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,492	
P. sericea, Cunningham in R. Br., prot. nov. 14 (1830) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,393	
P. Mitchellii, Meissner in Hook. Kew Misc. VII, 73 (1809) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,393	
P. fastigiata, R. Brown, prot. nov. 13 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,393	
P. hirsuta, Persoon, synops. plant. I, 118 (1805)	—	—	—	—	N.S.W.	—	—	B.fl.V,394	M.fr.VI,222.
P. Chamaepithys, Cunningham in Field, N.S. Wales, 329 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.V,394	M.fr.VI,223.
P. arborea, F. v. M., fragm. V, 37 (1865)	—	—	—	V.	—	—	—	B.fl.V,394	M.fr.V,37;VI,221.
P. salicina, Persoon, synops. plant. I, 118 (1805)...	—	—	—	V.	N.S.W.	—	—	B.fl.V,395	
P. prostrata, R. Brown in Transact. Linn. Soc. X, 163 (1809) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,395	
P. lanceolata, Andrews, Bot. Reposit. t. 74 (1800)	—	—	—	—	N.S.W.	—	—	B.fl.V,395	M.fr.VI,222.
P. confertiflora, Bentham, Fl. Austr. V, 396 (1870)	—	—	—	V.	—	—	—	B.fl.V,396	
P. lucida, R. Brown in Transact. Linn. Soc. X, 161 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,396	
P. linearis, Andrews, Bot. Reposit. t. 77 (1800)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,397	M.fr.VI,222.
P. pinifolia, R. Brown in Transact. Linn. Soc. X, 160 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,397	M.fr.VI,222;X,47.
P. Caleyi, R. Brown, prot. nov. 13 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,398	
P. ledifolia, Cunningham in De Cand. prodr. XIV, 339 (1856) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,398	
P. revoluta, Sieber in Roem. & Schult. mant. 272 (1822) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,398	M.fr.VI,221.
P. Gunnii, J. Hooker in Lond. Journ. Bot. VI, 283 (1847) ..	—	—	T.	—	—	—	—	B.fl.V,399	
P. mollis, R. Brown in Transact. Linn. Soc. X, 161 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,399	M.fr.VI,221. [47.
P. rigida, R. Brown, prot. nov. 14 (1830)	—	—	—	V.	N.S.W.	—	—	B.fl.V,399	M.fr.VI,221;VIII,149,X,
P. curvifolia, R. Brown, prot. nov. 13 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,400	
P. oblongata, Cunningham in R. Br. prot. nov. 14 (1830) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,400	
P. Cunninghamii, R. Brown, prot. nov. 15 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,400	
P. myrtilloides, Sieber in Roem. & Schult. mant. 272 (1822) ...	—	—	—	V.	N.S.W.	—	—	B.fl.V,401	M.fr.VI,221.
P. oxycoccoides, Sieber in Roem. & Schult. mant. 270 (1822) ...	—	—	—	—	N.S.W.	-	—	B.fl.V,401	M.fr.VI,221.
P. nutans, R. Brown in Transact. Linn. Soc. X, 162 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,401	M.fr.VI,222.
P. angulata, R. Brown, prot. nov. 14 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,402	
P. virgata, R. Brown in Transact. Linn. Soc. X, 161 (1809) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,402	
P. Chamaepeuce, Lhotsky in De Cand. prodr. XIV, 336 (1856)...	—	—	—	V.	N.S.W.	—	—	B.fl.V,402	M.fr.VIII,149;X,47.
P. juniperina, Labillardière, Nov. Holl. pl. spec. I, 33, t. 45 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,403	
P. tenuifolia, R. Brown, prot. nov. 12 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,403	
P. acerosa, Sieber in Roem. & Schult. mant. 269 (1822)	—	—	—	—	N.S.W.	—	—	B.fl.V,403	M.fr.VI,222.

[VII,59.

MACADAMIA, F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857). (Panopsis partly.)

M. ternifolia, F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,406	M.fr.II,91;VI,191;224,

HELICIA, Loureiro, Fl. Cochinch. I, 83 (1790).

H. Youngiana, C. Moore & F. v. M., fragm. IV, 84 (1864) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,406	M.fr.V,186.
H. praealta, F. v. M., fragm. III, 37 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,404	M.fr.IV,174;VI,224.
H. Australasica, F. v. M. in Hook. Kew Misc. IX, 27 (1857) ...	—	—	—	—	—	—	N.A.	B.fl.V,405	M.fr.II,91.
H. glabriflora, F. v. M., fragm. II, 91 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,405	M.fr.V,38,186. [224.
H. ferruginea, F. v. M., fragm. III, 37 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,405	M.fr.IV,174;V,186;VI,

ROUPALA, Aublet, Hist. des pl. de la Guian. I, 83, t. 32 (1775). (Rhopala, Bleasdalea. Adenostephanus partly.)

R. Bleasdalei, F. v. M. in proceed. Roy. Soc. N.S. Wales, 28 (1881)	—	—	—	—	—	Q.	—	B.fl.V,417	M.fr.V,90.

XYLOMELUM, Smith in Transact. Linn. Soc. IV, 214 (1798).

X. pyriforme, Smith in J. Knight, prot. 105 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,408	M.fr.VI,220.
X. occidentale, R. Brown, prot. nov. 31 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,408	
X. salicinum, Cunningham in R. Br. prot. 31 (1830)	—	—	—	—	—	Q.	—	B.fl.V,408	M.fr.IV,107,174;VI,220.
X. angustifolium, Kippist in Hook. Kew Misc. IV., 209 (1852)...	W.A.	—	—	—	—	—	—	B.fl.V,409	M.fr.VI,220.

LAMBERTIA, Smith in Transact. Linn. Soc. IV, 214 (1798).

L. uniflora, R. Brown in Transact. Linn. Soc. X, 188 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,413	M.fr.VI,248.
L. rariflora, Meissner in Lehm. pl. Preiss. II, 263 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,414	
L. inermis, R. Brown in Transact. Linn. Soc. X, 188 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,414	M.fr.VI,248.
L. ericifolia, R. Brown, prot. nov. 30 (1830)	W.A	—	—	—	—	—	—	B.fl.V,414	M.fr.VI,248.
L. multiflora, Lindley, Bot. Regist. XXV, App. XXXII (1839)	W.A.	—	—	—	—	—	—	B.fl.V,415	
L. formosa, Smith in Transact. Linn. Soc. IV, 214, t. 20 (1798)	W.A.	—	—	—	N.S.W.	—	—	B.fl.V,415	M.fr.VII,133.
L. echinata, R. Brown in Transact. Linn. Soc. X, 189 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,416	M.fr.VI,248.
L. ilicifolia, Hooker, icon. plant. t. 553 (1843)	W.A.	—	—	—	—	—	—	B.fl.V,416	

ORITES, R. Brown in Transact. Linn. Soc. X, 189 (1809). (Oritina.)

O. excelsa, R. Brown, prot. nov. 32 (1832)...	—	—	—	—	N.S.W.	—	—	B.fl.V,411	M.fr.V,153;VI,223.
O. diversifolia, R. Brown in Transact. Linn. Soc. X, 190 (1809)	—	—	T.	—	—	—	—	B.fl.V,411	M.fr.V,153.
O. Milligani, Meissner in Hook. Kew Misc. IV, 209 (1852) ...	—	—	T.	—	—	—	—	B.fl.V,411	[133.
O. lancifolia, F. v. M. in Transact. phil. Soc. Vict. I, 108 (1855)	—	—	—	V.	—	—	—	B.fl.V,412	M.fr.V,153;VI,223;VII,
O. revoluta, R. Brown in Transact. Linn. Soc. X, 19 (1809) ...	—	—	T.	—	—	—	—	B.fl.V,412	
O. acicularis, R. Brown, prot. nov. 32 (1830)	—	—	T.	—	—	—	—	B.fl.V,412	

STRANGEA, Meissner in Hooker's Kew Misc. VII, 66 (1855). (Molloya.)

S. linearis, Meissner in Hook. Kew Misc. VII, 66 (1855) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,453	M.fr.VI,223;VII,132.
S. cynanchocarpa, F. v. M., fragm. VII, 132 (1871)	W.A.	—	—	—	—	—	—	—	M.fr.VII,132.

GREVILLEA, R. Brown in Transact. Linn. Soc. X, 168 (1809). (Anadenia, Lyssanthe, Stylurus, Manglesia.)

G. Pinaster, Meissner in Hook. Kew Misc. VII, 76 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,427	M.fr.VI,212.
G. obtusifolia, Meissner in Hook. Kew Misc. IV, 187 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,427	
G. sparsiflora, F. v. M., fragm. VI, 206 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,428	M.fr.VI,206.
G. macrostylis, F. v. M., fragm. I, 137 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,428	M.fr.VI,211.
G. tripartita, Meissner in Hook. Kew Misc. IV, 186 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,428	M.fr.VI,213.
G. platypoda, F. v. M., fragm. VI, 205 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,428	M.fr.VI,205.
G. patentiloba, F. v. M., fragm. I, 137 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,429	M.fr.I,137.
G. pectinata, R. Brown, prot. nov. 23 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,429	M.fr.VI,212.
G. plurijuga, F. v. M., fragm. IV, 84 (1864)	W.A.	—	—	—	—	—	—	B.fl.V,430	M.fr.VI,212.
G. nudiflora, Meissner in Hook. Kew Misc. IV, 186 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,430	M.fr.I,35;VI,212.
G. stenomera, F. v. M., fragm. IV, 85 (1864)	W.A.	—	—	—	—	—	—	B.fl.V,430	M.fr.IV,85.
G. Thelemanniana, Endlicher, nov. stirp. dec. 6 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,431	M.fr.VI,212.
G. concinna, R. Brown in Transact. Linn. Soc. X, 172 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,431	M.fr.VI,212.
G. Hookeriana, Meissner in Lehm. pl. Preiss. I, 546 (1845) . .	W.A.	—	—	—	—	—	—	B.fl.V,432	M.fr.VI,212.
G. Baxteri, R. Brown, prot. nov. 22 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,432	
G. eriobotrya, F. v. M., fragm. X, 44 (1876)	W.A.	—	—	—	—	—	—	—	B.fl.X,44.
G. pterosperma, F. v. M. in Transact. phil. Soc. Vict. I, 22 (1854)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,432	M.fr.VI,210.
G. stenobotrya, F. v. M., fragm. IX, 3 (1875)	—	S.A.	—	—	—	—	N.A.	—	M.fr.IX,3.
G. eriostachya, Lindley, Bot. Regist. XXV, App. XXXVI (1839)	—	—	—	—	—	—	N.A.	B.fl.V,433	M.fr.VI,208&213.
G. thyrsoides, Meissner in Hooker's Kew Misc. VII, 77 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,433	
G. Chrysodendron, R. Brown in Transact. Linn. Soc. X, 176 (1809)	—	—	—	—	—	Q.	N.A.	B.fl.V,434	M.fr.VI,210. [46.
G. Banksii, R. Brown in Transact. Linn. Soc. X, 176 (1809) ...	—	—	—	—	—	Q.	—	B.fl.V,434	M.fr.VI,210;VIII,150;X,
G. Caleyi, R. Brown, prot. nov. 22 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,435	
G. asplenifolia, R. Brown in J. Knight, prot. 120 (1869)... ...	—	—	—	—	N.S.W.	—	—	B.fl.V,435	M.fr.VI,212.
G. cirsiifolia, Meissner in Lehm. pl. Preiss. II, 253 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,436	
G. laurifolia, Sieber in Roem. & Schult. mant. 279 (1822) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,436	[X,46.
G. Barklyana, F. v. M., gen. report 14 & 18 (1861)	—	—	—	V.	—	—	—	B.fl.V,436	M.fr.VII,133;VIII,150;
G. repens, F. v. M. in Schlecht. Linnaea XXVI, 355 (1853) ...	—	—	—	V.	—	—	—	B.fl.V,437	
G. Aquifolium, Lindley in Mitchell, Three Exped. II, 178 (1838)	—	—	—	V.	—	—	—	B.fl.V,437	
G. ilicifolia, R. Brown, prot. nov. 21 (1830)	—	S.A.	—	V.	—	—	—	B.fl.V,437	M.fr.VI,212.
G. Gaudichaudii, R. Brown in Freyc. Voy. Bot. 443, t. 46 (1826)	—	—	—	—	N.S.W.	—	—	B.fl.V,438	
G. acanthifolia, Cunningham in Field, N.S.Wales, 328 (1825) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,438	M.fr.VI,212;VIII,150.
G. bipinnatifida, R. Brown, prot. nov. 23 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,439	M.fr.VI,212.
G. armigera, Meissner in Hook. Kew Misc. IV, 186 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,439	
G. asparagoides, Meissner in Hook. Kew Misc. IV, 186 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,439	M.fr.IX,123.
G. Treueriana, F. v. M., fragm. IX, 123 (1875)	W.A.	S.A.	—	—	—	—	—	—	M.fr.IX,123.
G. floribunda, R. Brown, prot. nov. 19 (1830)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,440	
G. cinerea, R. Brown in Transact. Linn. Soc. X, 173 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,440	
G. alpina, Lindley in Mitchell, Three Exped. II, 179 (1838) ...	—	—	—	V.	—	—	—	B.fl.V,441	
G. montana, R. Brown in Transact. Linn. Soc. X, 172 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,441	
G. obtusiflora, R. Brown, prot. nov. 19 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,442	
G. arenaria, R. Brown in Transact. Linn. Soc. X, 172 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,442	
G. mucronulata, R. Brown in Transact. Linn. Soc. X, 173 (1809)	—	—	—	—	N.S.W.		—	B.fl.V,443	

G. Baueri, R. Brown in Transact. Linn. Soc. X, 173 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,443		
G. lanigera, Cunningham in R. Br., prot. nov. 20 (1830)... ...	—	—	—	V.	N.S.W.	—	—	B.fl.V,444		
G. rosmarinifolia, Cunningham in Field, N.S.Wales, 328 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.V,445		
G. Goodii, R. Brown in Transact. Linn. Soc. X, 174 (1809) ...	—	—	—	—	—	Q.	N.A.	B.fl.V,446	M.fr.VI,211.	
G. venusta, R. Brown in Transact. Linn. Soc. X, 175 (1809) ...	—	—	—	—	—	Q.	—	B.fl.V,446		
G. longistyla, Hooker in Mitchell, Trop. Austr. 343 (1848) ...	—	—	—	—	—	Q.	N.A.	B.fl.V,446	M.fr.VI,211.	
G. juncifolia, Hooker in Mitchell, Trop. Austr. 341 (1848) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,447	M.fr.VI,209;VIII,150.	
G. Wilsoni, Cunningham in Wilson's Voy. 273 (1835)	W.A.	—	—	—	—	—	—	B.fl.V,447	M.fr.VI,212;X,45.	
G. erectiloba, F. v. M., fragm. X, 44 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,44.	
G. lavandulacea, Schlechtendal, Linnaea XX, 586 (1847) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,448		
G. insignis, Kippist in De Cand., prodr. XIV, 379 (1856) ...	W.A.	—	—	—	—	—	—	B.fl.V,448	M.fr.VIII,150;X,45.	
G. Brownii, Meissner in Lehm. pl. Preiss. I, 537 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,449	M.fr.VI,213.	
G. fasciculata, R. Brown, prot. nov. 20 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,449		
G. aspera, R. Brown in Transact. Linn. Soc. X, 172 (1809) ...	W.A.	S.A.	—	—	—	—	—	B.fl.V,450	M.fr.VI,211.	
G. brachystylis, Meissner in Lehm. pl. Preiss. I, 538 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,450		
G. saccata, Bentham, Fl. Austr. V, 450 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,450		
G. Drummondii, Meissner in Lehm. pl. Preiss. I, 536 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,451		
G. disjuncta, F. v. M., fragm. VI, 206 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,451	M.fr.VI,206.	
G. haplantha, F. v. M. in Bentham Fl. Austr. V, 451 (1870) ...	W.A.	—	—	—	—	—	—	B.fl.V,452		
G. pinifolia, Meissner in Hook. Kew Misc. 186 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,452		
G. acuaria, F. v. M. in Bentham Fl. Austr. V, 452 (1870) ...	W.A.	—	—	—	—	—	—	B.fl.V,452		
G. singuliflora, F. v. M., fragm. VI, 92 (1868)	—	—	—	—	—	Q.	—	B.fl.V,452	M.fr.VI,92.	
G. pauciflora, R. Brown in Transact. Linn. Soc. X, 171 (1809)...	W.A.	S.A.	—	—	—	—	—	B.fl.V,453	M.fr.I,135;VI,206,208.	
G. quercifolia, R. Brown, prot. nov. 23 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,454	M.fr.VI,213.	
G. angulata, R. Brown, prot. nov. 24 (1830)	—	S.A.	—	—	—	—	N.A.	B.fl.V,455		
G. Wickhami, Meissner in Hook. Kew Misc. 187 (1852) ...	—	S.A.	—	—	—	—	N.A.	B.fl.V,455		
G. agrifolia, Cunningham in R. Br., prot. nov. 24 (1830) ...	—	—	—	—	—	—	N.A.	B.fl.V,455		
G. Cunninghamii, R. Brown, prot. nov. 23 (1830)...	—	—	—	—	—	—	N.A.	B.fl.V,456		
G. pungens, R. Brown in Transact. Linn. Soc. X, 175 (1809) ...	—	—	—	—	—	—	N.A.	B.fl.V,456		
G. Huegelii, Meissner in Lehm. pl. Preiss. I, 543 (1845)... ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,457	M.fr.VI.212;X,46.	
G. dimidiata, F. v. M., fragm. III, 146 (1863)	—	—	—	—	—	—	N.A.	B.fl.V,457	M.fr.III,146.	
G. heliosperma, R. Brown in Transact. Linn. Soc. X, 176 (1809)	—	—	—	—	—	—	N.A.	B.fl.V,457	M.fr.VI,210.	
G. refracta, R. Brown in Transact. Linn. Soc. X, 176 (1809) ...	—	—	—	—	—	—	N.A.	B.fl.V,458		
G. Dryandri, R. Brown in Transact. Linn. Soc. X, 175 (1809) ...	—	—	—	—	—	Q.	N.A.	B.fl.V,458	M.fr.VI,209.	
G. polystachya, R. Brown in Transact. Linn. Soc. X, 177 (1809)	—	—	—	—	—	Q.	N.A.	B.fl.V,459	M.fr.VI,210.	
G. robusta Cunningham in R. Br., prot. nov. 24 (1830)... ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,459	M.fr.VI.210;VII.133.[XI.20.	
G. annulifera, F. v. M., fragm. IV, 85 (1864)	W.A.	—	—	—	—	—	—	B.fl.V,460	M.fr.IV,85;VI,212;VII,132;	
G. leucopteris, Meissner in Hook. Kew Misc. VII, 76 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,460	M.fr.III,145;IV,85,176;	
G. Leucadendron, Cunningham in R. Br., prot. nov. 25 (1830)...	—	—	—	—	—	—	N.A.	B.fl.V,461	M.fr.I,136-7. [XI,20	
G. pyramidalis, Cunningham in R. Br., prot. nov. 25 (1830) ...	—	—	—	—	-	—	N.A.	B.fl.V,462		
G. striata, R. Brown in Transact. Linn. Soc. X, 177 (1809) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,462		
G. mimosoides, R. Brown in Transact. Linn. Soc. X, 177 (1809)	—	—	—	—	—	—	N.A.	B.fl.V,462		
G. Hilliana, F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,463		
G. gibbosa, R. Brown in Transact. Linn. Soc. X, 177 (1809) ...	—	—	—	—	—	Q.	N.A.	B.fl.V,463	M.fr.VI,210.	
G. buxifolia, R. Brown in Transact. Linn. Soc. X, 174 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,464		
G. phylicoides, R. Brown in Transact. Linn. Soc. X, 174 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,464	M.fr.VI,213.	
G. sphacelata, R. Brown in Transact. Linn. Soc. X, 174 (1809)...	—	—	—	—	N.S.W.	—	—	B.fl.V,464		
G. occidentalis, R. Brown in Transact. Linn. Soc. X, 173 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,465		
G. acerosa, F. v. M., fragm. I, 136 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,465	M.fr.I,136.	
G. umbellulata, Meissner in Lehm. pl. Preiss. II, 252 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,465		
G. oxystigma, Meissner in Lehm. pl. Preiss. I, 540 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,466		
G. Candolleana, Meissner in Lehm. pl. Preiss. I, 541 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,466	M.fr.VI,213.	
G. scabra, Meissner in Lehm. pl. Preiss. I, 541 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,466		
G. Miqueliana, F. v. M. in Transact. Vict. Inst. 132 (1855) ...	—	—	—	V.	—	—	—	B.fl.V,467		
G. Victoriae, F. v. M. in Transact. phil. Soc. Vict. I, 107 (1865)	—	—	—	V.	N.S.W.	—	—	B.fl.V,468	M.fr.VIII,142,150.	
G. punicea, R. Brown in Transact. Linn. Soc. X, 169 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,468	M.fr.XI,123.	
G. oleoides, Sieber in Roem. & Schult. mant. 277 (1822)... ...	—	—	—	V.	N.S.W.	—	—	B.fl.V,468		
G. trinervis, R. Brown, prot. nov. 18 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,469	M.fr.VI,211.	
G. juniperina, R. Brown in Transact. Linn. Soc. X, 171 (1809)...	—	—	—	—	N.S.W.	—	—	B.fl.V,469		
G. sericea, R. Brown in Transact. Linn. Soc. X, 170 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,470		
G. capitellata, Meissner in Hook. Kew Misc. IV, 187 (1852) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,470		
G. leiophylla, F. v. M. in Bentham, Fl. Austr. V, 471 (1870) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,471		
G. linearis, R. Brown in Transact. Linn. Soc. X, 170 (1809) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,471		
G. confertifolia, F. v. M. in Transact. phil. Soc. Vict. I, 22 (1854)	—	—	—	V.	—	—	—	B.fl.V,471	M.fr.VI,211.	
G. parviflora, R. Brown in Transact. Linn. Soc. X, 171 (1809) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,472		
G. australis, R. Brown in Transact. Linn. Soc. X, 171 (1809) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.V,472	M.fr.VII,133.	
G. commutata, F. v. M., fragm. VI, 207 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,473	M.fr.VII,207.	
G. pinnatisecta, F. v. M. in Bentham, Fl. Austr. V, 473 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,473		
G. argyrophylla, Meissner in Hook. Kew Misc. VII, 75 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,474	M.fr.VI,411.	
G. brachystachys, Meissner in Lehm. pl. Preiss. II, 254 (1847) ..	W.A.	—	—	—	—	—	—	B.fl.V,474	M.fr.VI,211.	
G. Endlicheriana, Meissner in Lehm. pl. Preiss. I, 546 (1845) ...	W.A.	—	—	—	-	—	—	B.fl.V,474	M.fr.VI,211.	
G. manglesoides, Meissner in Lehm. pl. Preiss. I, 547 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,475		
G. diversifolia, Meissner in Lehm. pl. Preiss. I, 547 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,475		
G. filifolia, Meissner in Lehm. pl. Preiss. I, 547 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,475		
G. hakeoides, Meissner in Lehm. pl. Preiss. II, 252 (1847) ...	W.A.	—	—	—	-	—	-	B.fl.V,476		
G. teretifolia, Meissner in Lehm. pl. Preiss. II, 255 (1847) ...	W.A.	—	—	-		—	—	B.fl.V,476		
G. eryngioides, Bentham, Fl. Austr. V, 476 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,476	M.fr.X,46.	
G. bracteosa, Meissner in Lehm. pl. Preiss. II, 254 (1847) ...	W.A.	—	—	—	—	-	—	B.fl.V,477		
G. crithmifolia, R. Brown, prot. nov. 23 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,477		
G. trachytheca, F. v. M., fragm. VI, 207 (1868)	W.A.	—		-	—		—	B.fl.V,477	M.fr.VI,207.	
G. triternata, R. Brown, prot. nov. 21 (1830)	—	—	—	V.	N.S.W.	—	—	B.fl.V,478		

G. ramosissima, Meissner in De Candolle, prodr. XIV, 388 ...	—	—	—	—	N.S.W.	—	—	B.fl.V,478	M.fr.VIII,150;X,46.
G. monticola, Meissner in Lehm. pl. Preiss. II, 259 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,478	M.fr.VI,209.
G. Muelleri, Bentham, Fl. Austr. V, 479 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,479	
G. trifida, Meissner in Lehm. pl. Preiss. I, 553 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,479	M.fr.VI,209.
G. Synapheae, R. Brown, prot. nov. 23 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,480	
G. flexuosa, Meissner in Lehm. pl. Preiss. I, 553 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,480	
G. leptobotrya, Meissner in Lehm. pl. Preiss. II, 256 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,480	M.fr.VI,209.
G. brevicuspis, Meissner in Lehm. pl. Preiss. II, 256 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,481	
G. intricata, Meissner in Hook. Kew Misc. VII, 74 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,481	
G. didymobotrya, Meissner in Hook. Kew Misc. IV, 186 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,481	M.fr.I,136;VI,208.
G. polybotrya, Meissner in Hook. Kew Misc. IV, 185 (1852) ...	W.A.	—	—	—	—	—	N.A.	B.fl.V,482	M.fr.IV,129;VI,208.
G. nematophylla, F. v. M., fragm. I, 136 (1859)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,482	M.fr.I,136;VI,208.
G. anethifolia, R. Brown, prot. nov. 21 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,482	M.fr.VI,209.
G. paradoxa, F. v. M., fragm. VI, 246 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,483	M.fr.VI,246.
G. petrophiloides, Meissner in Lehm. pl. Preiss. II, 257 (1847)...	W.A.	—	—	—	—	—	—	B.fl.V,483	M.fr.VI,209; X,46.
G. tenuiflora, Meissner in Lehm. pl. Preiss. I, 554 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,483	
G. pulchella, Meissner in Lehm. pl. Preiss. I, 553 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,484	M.fr.VI,209.
G. rudis, Meissner in Hook. Kew Misc. VII, 73 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,484	
G. apiciloba, F. v. M., fragm. X, 45 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,45.
G. Shuttleworthiana, Meissner in Lehm. pl. Preiss. II, 258 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,484	
G. integrifolia, Meissner in De Candolle, prodr. XIV, 385 (1856)	W.A.	—	—	—	—	—	—	B.fl.V,485	
G. stenocarpa, F. v. M. in Bentham Fl. Austr. V, 485 (1870) ...	W.A.	—	—	—	—	—	—	B.fl.V,485	
G. acrobotrya, Meissner in Hooker's Kew Misc. VII, 74 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,486	M.fr.VI,213.
G. glabrata, Meissner in Lehm. pl. Preiss. I, 549 (1845)... ...	W.A.	—	—	—	—	—	—	B.fl.V,486	
G. ornithopoda, Meissner in Lehm. pl. Preiss. II, 256 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,486	
G. paniculata, Meissner in Lehm. pl. Preiss. I, 550 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,487	
G. biternata, Meissner in Lehm. pl. Preiss. I, 549 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,487	M.fr.VI,213.
G. triloba, Meissner in Hooker's Kew Misc. VII, 74 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,487	M.fr.VI,213.
G. amplexans, F. v. M. in Bentham Fl. Austr. V, 488 (1870) ...	W.A.	—	—	—	—	—	—	B.fl.V,488	
G. vestita, Meissner in Lehm. pl. Preiss. I, 548 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,488	M.fr.VI,213.
G. tridentifera, Meissner in Lehm. pl. Preiss. I, 547 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,488	
G. erinacea, Meissner in Hooker's Kew Misc. VII, 74 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,489	M.fr.X,45.
HAKEA, Schrader, sert. Hannov. I, fasc. 3, 27, t. 17 (1797). (Conchium.)									
H. chordophylla, F. v. M. in Hooker's Kew Misc. IX, 23 (1857)	—	—	—	—	—	—	N.A.	B.fl.V,495	M.fr.VI,190.
H. Cunninghamii, R. Brown, prot. nov. 26 (1830)	—	—	—	—	—	—	N.A.	B.fl.V,495	M.fr.VI,190.
H. lorea, R. Brown, prot. nov. 25 (1830)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.V,495	M.fr.VI,189;VII,133.
H. Fraseri, R. Brown, prot. nov. 26 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,496	
H. macrocarpa, Cunningham in R. Brown, prot. nov. 30 (1830)	—	—	—	—	—	—	N.A.	B.fl.V,496	M.fr.VI,190,210.
H. arborescens, R. Brown in Transact. Linn. Soc. X, 187 (1809)	—	—	—	—	—	Q.	N.A.	B.fl.V,497	M.fr.VI,190.
H. stenophylla, Cunningham in De Cand. prodr. XIV, 417 (1856)	—	—	—	—	—	—	N.A.	B.fl.V,497	
H. trineura, F. v. M., fragm. III, 146 (1863)	—	—	—	—	—	Q.	—	B.fl.V,497	M.fr.VI,216.
H. cyclocarpa, Lindley, Bot. Regist. XXV, App. XXXVI (1839)	W.A.	—	—	—	—	—	—	B.fl.V,498	
H. crassifolia, Meissner in Lehm. pl. Preiss. I, 570 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,498	M.fr.VI,216.
H. pandanocarpa, R. Brown, prot. nov. 29 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,499	M.fr.VI,216.
H. Roei, Bentham, Fl. Austr. V, 499 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,499	
H. adnata, R. Brown, prot. nov. 26 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,499	
H. obliqua, R. Brown in Transact. Linn. Soc. X, 180 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,500	M.fr.VI,218.
H. Hookeriana, Meissner in Hook. Kew Misc. IV, 208 (1852) ..	W.A.	—	—	—	—	—	—	B.fl.V,500	M.fr.VI,216.
H. incrassata, R. Brown, prot. nov. 29 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,500	
H. flabellifolia, Meissner in Hooker's Kew Misc. VII, 116 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,501	
H. Brownii, Meissner in Lehm. pl. Preiss. I, 569 (1845)... ...	W.A.	—	—	—	—	—	—	B.fl.V,501	
H. Baxteri, R. Brown, prot. nov. 28 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,501	
H. ceratophylla, R. Brown in Transact. Linn. Soc. X, 184 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,501	M.fr.IV,49;VI,217.
H. lasiantha, R. Brown, prot. nov. 29 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,502	M.fr.VI,215.
H. eriantha, R. Brown, prot. nov. 29 (1830)	—	—	—	V.	N.S.W.	—	—	B.fl.V,502	M.fr.VI,217.
H. megalosperma, Meissner in Hook. Kew Misc. VII, 117 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,503	
H. clavata, Labillardière, Nov. Holl. pl. spec. I, 31, t. 41 (1804)	W.A.	—	—	—	—	—	—	B.fl.V,503	M.fr.VI,216.
H. orthorrhyncha, F. v. M., fragm. V, 214 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,503	M.fr.VI,214.
H. Candolleana, Meissner in Lehm. pl. Preiss. II, 262 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,504	M.fr.VI,215.
H. trifurcata, R. Brown in Transact. Linn. Soc. X, 183 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,504	M.fr.VI,220.
H. erinacea, Meissner in Lehm. pl. Preiss. I, 559 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,505	M.fr.VI,220.
H. platysperma, Hooker, icones pl. t. 433 (1842)	W.A.	—	—	—	—	—	—	B.fl.V,505	
H. brachyptera, Meissner in Hook. Kew Misc. IV, 208 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,505	M.fr.VI,217.
H. Kippistiana, Meissner in Hooker's Kew Misc. VII, 115 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,506	
H. Preissii, Meissner in Lehm. pl. Preiss. I, 557 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,506	M.fr.VI,219.
H. pugioniformis, Cavanilles, Annal. hist. nat. I, 213, t. 11 (1800)	—	—	T.	V.	N.S.W.	—	—	B.fl.V,506	M.fr.VI,217.
H. Pampliniana, Kippist in Hook. Kew Misc. VII, 115 (1855)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,507	
H. vittata, R. Brown in Transact. Linn. Soc. X, 182 (1809) ...	—	S.A.	—	—	—	—	—	B.fl.V,507	
H. rostrata, F. v. M. in Schlecht. Linnaea XXVI, 259 (1853) .	—	S.A.	—	V.	—	—	—	B.fl.V,508	M.fr.VI,218;VII,133.
H. rugosa, R. Brown in Transact. Linn. Soc. X, 179 (1809) ...	—	S.A.	—	V.	—	—	—	B.fl.V,508	M.fr.VI,218.
H. Epiglottis, Labillardière, Nov. Holl. pl. spec. I, 30, t. 40 (1804)	—	—	T.	—	—	—	—	B.fl.V,508	
H. amplexicaulis, R. Brown in Transact. Linn. Soc. X, 184 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,509	M.fr.VI,217;VII,133.
H. glabella, R. Brown, prot. nov. 28 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,509	
H. auriculata, Meissner in Hook. Kew Misc. VII, 116 (1855)...	W.A.	—	—	—	—	—	—	B.fl.V,510	M.fr.VI,217.
H. cristata, R. Brown, prot. nov. 28 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,510	
H. linearis, R. Brown in Transact. Linn. Soc. X, 183 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,511	
H. stenocarpoides, F. v. M. in Bentham Fl. V, 511 (1870) ...	W.A.	—	—	—	—	—	—	B.fl.V,511	
H. ruscifolia, Labillardière, Nov. Holl. pl. spec. I, 30, t. 39 (1804)	W.A.	—	—	—	—	—	—	B.fl.V,511	
H. saligna, R. Brown in J. Knight, prot. 108 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,512	M.fr.VI,217.
H. verrucosa, F. v. M., fragm. V, 25 (1865)	W.A.	—	—	—	—	—	—	B.fl.V,512	M.fr.V,25;VI,218.

H. rhombales, F. v. M., fragm. X, 90 (1876)	—	—	—	—	—	—	N.A.	—		M.fr.X,90.
H. purpurea, Hooker in Mitch. Trop. Austr. 348 (1848)... ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,513		
H. gibbosa, Cavanilles, Anal. hist. nat. I, 215 (1800)	—	—	—	—	N.S.W.	—	—	B.fl.V,513		M.fr.VI,218.
H. propinqua, Cunningham in Field N.S. Wales, 327 (1825) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,513		M.fr.V,26.
H. nodosa, R. Brown in Transact. Linn. Soc. X, 179 (1809) ...	—	—	T.	V.	—	—	—	B.fl.V,514		M.fr,V,26;VI,217.
H. acicularis, R. Brown in Transact. Linn. Soc. X, 181 (1809)...	—	—	T.	V.	N.S.W.	—	—	B.fl.V,514		M.fr.VI,218.
H. leucoptera, R. Brown in Transact. Linn. Soc. X, 180 (1809)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,515		M.fr.VI,219;VII,133.
H. cycloptera, R. Brown in Transact. Linn. Soc. X, 182 (1809)	—	S.A.	—	—	—	—	—	B.fl.V,515		M.fr.VI,217.
H. microcarpa, R. Brown in Transact. Linn. Soc. X, 182 (1809)	—	—	T.	V.	N.S.W.	—	—	B.fl.V,516		M.fr.VI,217.
H. recurva, Meissner in Hook. Kew Misc. IV, 207 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,516		M.fr.VI,218.
H. circumalata, Meissner in Hook. Kew Misc. VII, 114 (1855)...	W.A.	—	—	—	—	—	—	B.fl.V,516		M.fr.VI,219.
H. commutata, F. v. M., fragm. V, 26 (1865)	W.A.	—	—	—	—	—	—	B.fl.V,517		M.fr.VI,219.
H. strumosa, Meissner in Hook. Kew Misc. IV, 208 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,517		M.fr.VI,219.
H. multilineata, Meissner in Lehm. pl. Preiss. II, 261 (1847) ...	W.A.	S.A.	—	—	—	—	—	B.fl.V,518		M.fr.VI,215.
H. laurina, R. Brown, prot. nov. 29 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,518		M.fr.IV,130,216.
H. obtusa, Meissner in Hook. Kew Misc. IV, 209 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,519		
H. cinerea, R. Brown in Transact. Linn. Soc. X, 186 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,519		
H. corymbosa, R. Brown, prot. nov. 28 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,519		M.fr.VI,216.
H. undulata, R. Brown in Transact. Linn. Soc. X, 185 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,520		M.fr.VI,216.
H. petiolaris, Meissner in Lehm. pl. Preiss. I, 577 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,520		
H. neurophylla, Meissner in Hook. Kew Misc. VII, 117 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,521		
H. loranthifolia, Meissner in Lehm. pl. Preiss. I, 574 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,521		
H. cucullata, R. Brown, prot. nov. 30 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,521		M.fr.VI,216.
H. ferruginea, Sweet, Fl. Austral. t. 45 (1828)	W.A.	—	—	—	—	—	—	B.fl.V,522		M.fr.VI,216.
H. smilacifolia, Meissner in Lehm. pl. Preiss. I, 567 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,522		M.fr.VI,217.
H. elliptica, R. Brown in Transact. Linn. Soc. X, 187 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,523		M.fr.VI,216.
H. ambigua, Meissner in Lehm. pl. Preiss. II, 260 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,523		
H. plurinervia, F. v. M. in Benth. Fl. Austr. V, 523 (1870) ...	—	—	—	—	—	Q.	—	B.fl.V,523		
H. dactyloides, Cavanilles, Anal. hist. nat. I, 215, t. 12 (1800)...	—	—	—	—	N.S.W.	—	—	B.fl.V,524		M.fr.VI,215.
H. ulicina, R. Brown, prot. nov. 29 (1830)...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,524		
H. falcata, R. Brown, prot. nov. 29 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,524		
H. pycnoneura, Meissner in Hook. Kew Misc. VII, 117 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,525		M.fr.VI,215.
H. stenocarpa, R. Brown, prot. nov. 29 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,525		
H. marginata, R. Brown in Transact. Linn. Soc. X, 185 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,556		M.fr.VI,216.
H. myrtoides, Meissner in Lehm. pl. Preiss. I, 577 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,526		
H. costata, Meissner in Lehm. pl. Preiss. I, 575 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,526		
H. oleifolia, R. Brown in Transact. Linn. Soc. X, 185 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,527		M.fr.VI,215.
H. florida, R. Brown in Transact. Linn. Soc. X, 183 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,527		
H. varia, R. Brown in Transact. Linn. Soc. X, 183 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,527		M.fr.VI,219.
H. sulcata, R. Brown in Transact. Linn. Soc. X, 180 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,528		M.fr.VI,219.
H. Meissneriana, Kippist in Hook. Kew Misc. VII, 114 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,529		M.fr.VI,219.
H. subsulcata, Meissner in Lehm. pl. Preiss. I, 555 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,529		M.fr.VI,219.
H. Lehmanniana, Meissner in Lehm. pl. Preiss. I, 557 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,529		M.fr.VI,218.
H. flexilis, F. v. M. in Schlecht. Linnaea XXVI, 359 (1853) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,530		M.fr.VI,216.
H. nitida, R. Brown in Transact. Linn. Soc. X, 184 (1809) ...	W.A.	—	—	—	—		—	B.fl.V,530		M.fr.V,72;VI,217.
H. Oldfieldii, Bentham, Fl. Austr. V, 530 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,530		
H. suaveolens, R. Brown in Transact. Linn. Soc. X, 182 (1809)	W.A.	—	—	—			—	B.fl.V,531		M.fr.VI,220.
H. lissocarpha, R. Brown, prot. nov. 27 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,531		M.fr.VI,219.
H. bipinnatifida, R. Brown, prot. nov. 28 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,532		
CANARVONIA, F. v. M., fragm. VI, 81 (1867).										
C. aralifolia, F. v. M., fragm. VI, t. 55 (1867)	—	—	—	—		Q.	—	B.fl.V,410		M.fr.VI,81,247;VII,59.
BUCKINGHAMIA, F. v. M., fragm. VI, 248 (1868).										
B. celsissima, F. v. M., fragm. VI, 248 (1868)	—	—	—	—		Q.	—	B.fl.V,532		M.fr.VI,248.
DARLINGIA, F. v. M., fragm. V, 152 (1866).										
D. spectatissima, F. v. M., fragm. V, 152 (1866)	—	—	—	—	—	Q.	—	B.fl.V,533		M.fr.X,90.
CARDWELLIA, F. v. M., fragm. V, 23 (1865).										
C. sublimis, F. v. M., fragm. V, 24 (1865)...	—	—	—	—	—	Q.	—	B.fl.V,538		M.fr.V,24.
STENOCARPUS, R. Brown in Transact. Linn. Soc. X, 201 (1809). (Cybele, Agnostus.)										
S. sinuatus, Endlicher, gen. pl. suppl. IV, 88 (1847)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,539		M.fr.VI,224. [VII,133.
S. salignus, R. Brown in Transact. Linn. Soc. X, 202 (1809) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,539		M.fr.I,135,244;VI,224;
S. Cunninghamii, R. Brown, prot. nov. 34 (1830)...	—	—	—	—	—	—	N.A.	B.fl.V,540		
LOMATIA, R. Brown in Transact. Linn. Soc. X, 199 (1809). (Tricondylus.)										
L. fraxinifolia, F. v. M. in Benth. Fl. Austr. V, 536 (1870) ...	—	—	—	—	—	Q.	—	B.fl.V,536		
L. ilicifolia, R. Brown in Transact. Linn. Soc. X, 200 (1809) ...	—	—	—	V.	N.S.W.	—	—	B.fl.V,536		M.fr.VII,133.
L. longifolia, R. Brown in Transact. Linn. Soc. X, 200 (1809) ...	—	—	—	V.	N.S.W.	—	—	B.fl.V,537		
L. silaifolia, R. Brown in Transact. Linn. Soc. X, 199 (1809) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,537		M.fr.V,153;VI,191.
L. tinctoria, R. Brown in Transact. Linn. Soc. X, 199 (1809) ...	—	—	T.	—	—	—	—	B.fl.V,537		M.fr.VI,224.
L. polymorpha, R. Brown in Transact. Linn. Soc. X, 200 (1809)	—	—	T.	—	—	—	—	B.fl.V,538		
EMBOTHRIUM, R. & G. Forster, charact. gen. 15, t. 8 (1776).										
E. Wickhami, F. v. M., fragm. VIII, 164 (1874)...	—	—	—	—	—	Q.	—	—		M.fr.VIII,164;IX,194.
TELOPEA, R. Brown in Transact. Linn. Soc. X, 197 (1809).										
T. speciosissima, R. Brown in Transact. Linn. Soc. X, 198 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,534		M.fr.X,90.
T. oreades, F. v. M., Annual Report 18 (1861)	—	—	—	V.	—	—	—	B.fl.V,534		M.fr.II,170;VII,133.
T. truncata, R. Brown in Transact. Linn. Soc. X, 198 (1809) ...	—	—	T.	—	—	—	—	B.fl.V,535		M.fr.II,171.
BANKSIA, Linné, fil. suppl. 15 et 126 (1781).										
B. pulchella, R. Brown in Transact. Linn. Soc. X, 202 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,544		M.fr.VII,54.
B. Meissneri, Lehmann, plant. Preiss. I, 582 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,545		

B. nutans, R. Brown in Transact. Linn. Soc. X, 203 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,545	M.fr.IV,108;VII,54.
B. sphaerocarpa, R. Brown in Transact. Linn. Soc. X, 203 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,546	M.fr.VII,54.
B. tricuspis, Meissner in Hook. Kew Misc. VII, 118 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,546	M.fr.VII,54.
B. occidentalis, R. Brown in Transact. Linn. Soc. X, 204 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,546	M.fr.VII,54.
B. littoralis, R. Brown in Transact. Linn. Soc. X, 204 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,547	M.fr.VII,55.
B. ericifolia, Linné, fil. suppl. 127 (1781)	—	—	—	—	N.S.W.	—	—	B.fl.V,547	M.fr.VII,54.
B. spinulosa, Smith, Specim. Bot. New Holl. 13, t. 4 (1793) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,547	
B. collina, R. Brown in Transact. Linn. Soc. X, 204 (1809) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,548	M.fr.VII,54.
B. verticillata, R. Brown in Transact. Linn. Soc. X, 207 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,548	
B. dryandroides, Baxter in Sweet, Fl. Austral. t. 56 (1828) ...	W.A.	—	—	—	—	—	—	B.fl.V,549	M.fr.VII,58.
B. Brownii, Baxter in R. Br. prot. nov. 37 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,549	M.fr.VII,58.
B. attenuata, R. Brown in Transact. Linn. Soc. X, 209 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,549	M.fr.VII,55.
B. media, R. Brown, prot. nov. 35 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,550	M.fr.IV,109.
B. Solandri, R. Brown, prot. nov. 35 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,550	M.fr.VII,58.
B. Goodii, R. Brown, prot. nov. 36 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,550	M.fr.VII,58.
B. petiolaris, F. v. M., fragm. IV, 109 (1864)	W.A.	—	—	—	—	—	—	B.fl.V,551	M.fr.IV,109.
B. repens, Labillardière, Voy. I, 411, t. 23 (1798)	W.A.	—	—	—	—	—	—	B.fl.V,551	M.fr.IV,108,177;VII,58.
B. prostrata, R. Brown, prot. nov. 36 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,551	M.fr.VII,57.
B. grandis, Willdenow, spec. plant. I, 535 (1797)	W.A.	—	—	—	—	—	—	B.fl.V,552	M.fr.VII,57.
B. quercifolia, R. Brown in Transact. Linn. Soc. X, 210 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,552	M.fr.VII,57.
B. Baueri, R. Brown, prot. nov. 35 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,552	M.fr.IV,107;VII,56.
B. marginata, Cavanilles, Anal. hist. nat. I, 227, t. 13 (1800) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,553	M.fr.VII,55,133;X,90.
B. integrifolia, Linné, fil. suppl. 127 (1781)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,554	M.fr.VII,55.
B. dentata, Linné, fil. suppl. 127 (1781)	—	—	—	—	—	Q.	N.A.	B.fl.V,555	M.fr.VII,57.
B. latifolia, R. Brown in Transact. Linn. Soc. X, 208 (1809) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,555	M.fr.VII,56.
B. serrata, Linné, fil. suppl. 126 (1781)	—	—	T.	V.	N.S.W.	—	—	B.fl.V,556	M.fr.VII,56.
B. aemula, R. Brown in Transact. Linn. Soc. X, 210 (1809) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,556	M.fr.VII,56.
B. ornata, F. v. M. in Schlecht. Linnaea XXVI, 352 (1853) ...	—	S.A.	—	V.	—	—	—	B.fl.V,557	M.fr.VII,56.
B. coccinea, R. Brown in Transact. Linn. Soc. X, 207 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,557	M.fr.VII,56.
B. Sceptrum, Meissner in Hook. Kew Misc. VII, 120 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,557	M.fr.VII,56.
B. Menziesii, R. Brown, prot. nov. 36 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,558	
B. laevigata, Meissner in Hook. Kew Misc. IV, 210 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,558	M.fr.VII,56.
B. Hookeriana, Meissner in Hook. Kew Misc. VII, 119 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,558	
B. prionotes, Lindley, Bot. Regist. XXV, App. XXXIV (1839)	W.A.	—	—	—	—	—	—	B.fl.V,558	M.fr.VII,56.
B. Victoriae, Meissner in Hook. Kew Misc. VII, 119 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,559	M.fr.VII,58.
B. speciosa, R. Brown in Transact. Linn. Soc. X, 210 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,559	M.fr.VII,58.
B. Baxteri, R. Brown, prot. nov. 36 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,559	M.fr.VII,58.
B. marcescens, R. Brown in Transact. Linn. Soc. X, 208 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,560	M.fr.VII,56.
B. Lemanniana, Meissner in Hook. Kew Misc. 210 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,560	M.fr.VII,57.
B. Caleyi, R. Brown, prot. nov. 35 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,560	M.fr.VII,57.
B. Lindleyana, Meissner in Hook. Kew Misc. VII, 120 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,561	M.fr.VII,55.
B. elegans, Meissner in Hook. Kew Misc. VII, 119 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,561	
B. Candolleana, Meissner in Hook. Kew Misc. VII, 118 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,561	M.fr.VII,58.
B. ilicifolia, R. Brown in Transact. Linn. Soc. X, 211 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,561	M.fr.VII,58.
DRYANDRA, R. Brown in Transact. Linn. Soc. X, 211 (1809). (Josephia, Hemiclidia.)									
D. quercifolia, Meissner in Hook. Kew Misc. IV, 210 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,566	M.fr.VII,50.
D. praemorsa, Meissner in Lehm. pl. Preiss. II, 265 (1847) .	W.A.	—	—	—	—	—	—	B.fl.V,566	M.fr.VII,50.
D. cuneata, R. Brown in Transact. Linn. Soc. X, 212 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,566	M.fr.VII,50.
D. falcata, R. Brown in Transact. Linn. Soc. X, 213 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,567	
D. armata, R. Brown in Transact. Linn. Soc. X, 212 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,567	M.fr.VII,50.
D. longifolia, R. Brown in Transact. Linn. Soc. X, 215 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,568	
D. Fraseri, R. Brown, prot. nov. 39 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,568	M.fr.VII,52.
D. floribunda, R. Brown in Transact. Linn. Soc. X, 212 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,569	M.fr.VI,92;VII,50.
D. carduacea, Lindley, Bot. Regist. XXV, App. XXXIII (1839)	W.A.	—	—	—	—	—	—	B.fl.V,569	M.fr.VII,51.
D. carlinoides, Meissner in Lehm. pl. Preiss. II, 267 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.V,569	M.fr.VII,53.
D. polycephala, Bentham, Fl. Austr. V, 570 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,570	
D. Kippistiana, Meissner in Hook. Kew Misc. VII, 122 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,570	M.fr.VII,52.
D. squarrosa, R. Brown, prot. nov. 38 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,571	M.fr.VII,52.
D. Serra, R. Brown, prot. nov. 38 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,571	M.fr.VII,51.
D. concinna, R. Brown, prot. nov. 38 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,571	M.fr.VII,52.
D. foliolata, R. Brown, prot. nov. 38 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,572	M.fr.VII,51.
D. stupposa, Lindley, Bot. Regist. XXV, App. XXXIII (1839)	W.A.	—	—	—	—	—	—	B.fl.V,572	M.fr.VII,51.
D. nobilis, Lindley, Bot. Regist. XXV, App. XXXIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.V,572	
D. mucronulata, R. Brown in Transact. Linn. Soc. X, 213 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,573	
D. formosa, R. Brown in Transact. Linn. Soc. X, 213, t. 3 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,573	M.fr.VII,51.
D. Baxteri, R. Brown, prot. nov. 38 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,573	M.fr.VII,51.
D. nivea, R. Brown in Transact. Linn. Soc. X, 214 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,574	M.fr.VII,52.
D. Arctotidis, R. Brown, prot. nov. 39 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,574	
D. nana, Meissner in Hook. Kew Misc. VII, 121 (1855)... ...	W.A.	—	—	—	—	—	—	B.fl.V,575	M.fr.VII,52.
D. Preissii, Meissner in Lehm. pl. Preiss. I, 599 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,575	M.fr.VII,54.
D. sclerophylla, Meissner in Hook. Kew Misc. VII, 123 (1855)...	W.A.	—	—	—	—	—	—	B.fl.V,576	M.fr.VII,52.
D. pulchella, Meissner in Hook. Kew Misc. IV, 211 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,576	
D. plumosa, R. Brown in Transact. Linn. Soc. X, 214 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,576	M.fr.VII,51.
D. senecionifolia, R. Brown, prot. nov. 39 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,53.
D. vestita, Kippist in Hook. Kew Misc. VII, 121 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,53.
D. cirsioides, Meissner in Hook. Kew Misc. IV, 211 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,52.
D. Hewardiana, Meissner in Hook. Kew Misc. IV, 210 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,53.
D. patens, Bentham, Fl. Austr. V, 578 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,578	
D. conferta, Bentham, Fl. Austr. V, 578 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,578	
D. horrida, Meissner in Hook. Kew Misc. IV, 211 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,579	M.fr.VII,52.

D. serratuloides, Meissner in Hook. Kew Misc. VII, 123 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,579	M.fr.VII,52.
D. comosa, Meissner in Hook. Kew Misc. IV, 211 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,579	M.fr.VII,53.
D. Shuttleworthiana, Meissner in Hook. Kew Misc. VII, 122 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,580	M.fr.VII,52.
D. speciosa, Meissner in Hook. Kew Misc. IV, 211 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,580	
D. tridentata, Meissner in Hook. Kew Misc. VII, 120 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,580	M.fr.VII,53.
D. tenuifolia, R. Brown in Transact. Linn. Soc. X, 215 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,581	M.fr.VII,53.
D. proteoides, Lindley, Bot. Regist. XXV, App. XXXIII (1839)	W.A.	—	—	—	—	—	—	B.fl.V,582	
D. runcinata, Meissner in Hook. Kew Misc. IV, 210 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.V,582	
D. obtusa, R. Brown in Transact. Linn. Soc. X, 214 (1809) ...	W.A.	—	—	—	—	—	—	B.fl.V,582	M.fr.V,185.
D. bipinnatifida, R. Brown, prot. nov. 39 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,583	
D. pteridifolia, R. Brown in Transact. Linn. Soc. X, 215 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,583	M.fr.VII,54.
D. calophylla, R. Brown, prot. nov. 40 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,583	M.fr.VII,54.

THYMELEAE.

A. L. de Jussieu, gen. plant. 76 (1789).

PIMELEA, Banks & Solander in Gaert. de fruct. I, 186 (1788). (Thecanthes, Gymnococca, Heterolaena, Calyptrostegia, Macrostegia;—Banksia, Forst., 1776.)

P. punicea, R. Brown, prodr. 359 (1810)	—	—	—	—	—	—	N.A.	B.fl.VI,6	M.fr.V,74;VII,3.
P. concreta, F. v. M., fragm. V, 73 (1865)	—	—	—	—	—	—	N.A.	B.fl.VI,6	M.fr.V,73.
P. cornucopiae, Vahl, enum. plant. I, 305 (1804)	—	—	—	—	—	Q.	—	B.fl.VI,6	M.fr.VII,3.
P. sanguinea, F. v. M., fragm. I, 84 (1858)	—	—	—	—	—	Q.	—	B.fl.VI,7	M.fr.VII,3;XI,47.
P. alpina, F. v. M., second general report, 14 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,7	M.fr.VII,4.
P. longifolia, Banks & Solander in Rees, Cycl. XXVI (1814) ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,7	M.fr.VIII,9.
P. cinerea, R. Brown, prodr. 361 (1810)	—	—	T.	—	—	—	—	B.fl.VI,8	
P. Milligani, Meissner in De Candolle, prodr. XIV, 509 (1856) ...	—	—	T.	—	—	—	—	B.fl.VI,8	
P. spectabilis, Lindley, Bot. Regist. t. 33 (1841)	W.A.	—	—	—	—	—	—	B.fl.VI,9	M.fr.VII,4.
P. Lehmanniana, Meissner in Lehm. pl. Preiss. I, 603 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.VI,9	M.fr.VII,4.
P. hispida, R. Brown, prodr. 360 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,10	
P. rosea, R. Brown, prodr. 360 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,10	
P. ferruginea, Labillardière, Nov. Holl. pl. spec. I, 10, t. 5 (1804)	W.A.	—	—	—	—	—	—	B.fl.VI,10	M.fr.VII,4.
P. brachyphylla, Bentham, Fl. Austr. VI, 11 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,11	
P. sylvestris, R. Brown, prodr. 361 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,11	M.fr.VII,2&5.
P. brevifolia, R. Brown, prodr. 359 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,12	M.fr.VII,4.
P. Maxwelli, F. v. M. in Benth. Fl. Austr. VI, 12 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VI,12	
P. angustifolia, R. Brown, prodr. 360 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,13	M.fr.VII,3.
P. nervosa, Meissner in Lehm. pl. Preiss. II, 269 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.VI,13	M.fr.VII,3.
P. sulphurea, Meissner in Mohl & Schlecht. Bot. Zeitung, 396 (1848)	W.A.	—	—	—	—	—	—	B.fl.VI,14	M.fr.VII,5.
P. floribunda, Meissner in De Candolle, prodr. XIV, 505 (1856)	W.A.	—	—	—	—	—	—	B.fl.VI,14	M.fr.VII,5.
P. suaveolens, Meissner in Lehm. pl. Preiss. I, 603 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.VI,14	M.fr.VII,2,4,5.
P. physodes, Hooker, icon. plant. t. 865 (1852)	W.A.	—	—	—	—	—	—	B.fl.VI,15	M.fr.VII,4.
P. glauca, R. Brown, prodr. 360 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,15	M.fr.VII,4;XI,123.
P. colorans, Cunningham in De Cand. prodr. XIV, 499 (1856) ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,16	
P. collina, R. Brown, prodr. 359 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,16	
P. spathulata, Labillardière, Nov. Holl. pl. spec. I, 9, t. 4 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,17	
P. linifolia, Smith, Specim. of Bot. of N. Holl. 31, t. 11 (1793) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,17	M.fr.VII,3.
P. ligustrina, Labillardière, Nov. Holl. pl. spec. I, 9, t. 3 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,18	M.fr.I,87;IV,163;VII,5.
P. humilis, R. Brown, prodr. 361 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,19	M.fr.VII,4;XI,47.
P. sericea, R. Brown, prodr. 361 (1810)	—	—	T.	—	—	—	—	B.fl.VI,19	M.fr.VII,5.
P. nivea, Labillardière, Nov. Holl. pl. spec. I, 10, t. 6 (1804) ...	—	—	T.	—	—	—	—	B.fl.VI,20	M.fr.VII,5.
P. imbricata, R. Brown, prodr. 361 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,20	M.fr.VII,5.
P. villifera, Meissner in Lehm. pl. Preiss. II, 271 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.VI,21	
P. drupacea, Labillardière, Nov. Holl. pl. spec. I, 10, t. 7 (1804)	—	—	T.	V.	—	—	—	B.fl.VI,21	M.fr.VII,8.
P. haematostachya, F. v. M., fragm. I, 84 (1858)	—	—	—	—	—	Q.	—	B.fl.VI,22	M.fr.I,84;VII,8;XI,47.
P. spicata, R. Brown, prodr. 362 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,22	M.fr.VII,7;VIII,9.
P. spiculigera, F. v. M. in Bentham Fl. Austr. VI, 23 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VI,23	M.fr.XI,46.
P. Forrestiana, F. v. M., fragm. XI, 46 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,46.
P. filiformis, J. Hooker in Lond. Journ. VI, 280 (1847)	—	—	T.	—	—	—	—	B.fl.VI,23	M.fr.VII,7.
P. latifolia, R. Brown, prodr. 362 (1810)	—	—	—	—	—	Q.	—	B.fl.VI,23	M.fr.IV,49;VII,7.
P. simplex, F. v. M. in Linnaea XXV, 443 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,23	
P. sericostachya, F. v. M., fragm. IV, 162 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,24	M.fr.XI,47.
P. trichostachya, Lindley in Mitch. Trop. Austr. 355 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,24	M.fr.VII,7.
P. leptostachya, Bentham, Fl. Austr. VI, 24 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,24	
P. argentea, R. Brown, prodr. 362 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,25	M.fr.VII,7.
P. clavata, Labillardière, Nov. Holl. pl. spec. I, 11 (1804) ...	W.A.	—	—	—	—	—	—	B.fl.VI,25	M.fr.VI,160;VII,5.
P. axiflora, F. v. M., First general Report 17 (1853)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,26	M.fr.VII,7.
P. leptospermoides, F. v. M., fragm. VII, 2 (1870)	—	—	—	—	—	Q.	—	B.fl.VI,27	M.fr.VII,2.
P. microcephala, R. Brown, prodr. 361 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,27	M.fr.XI,27,47.
P. pauciflora, R. Brown, prodr. 360 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,27	M.fr.VII,8;XI,47.
P. elachantha, F. v. M., First general Report 17 (1853)	—	—	—	V.	—	—	—	B.fl.VI,28	M.fr.VII,6.
P. pygmaea, F. v. M., in Schlecht. Linnaea XXVI, 346 (1853)	—	—	T.	—	—	—	—	B.fl.VI,28	
P. serpyllifolia, R. Brown, prodr. 360 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,29	M.fr.VII,6.
P. flava, R. Brown, prodr. 361 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,29	M.fr.VII,6.
P. petrophila, F. v. M. in Schlecht. Linnaea XXV, 442 (1852) ...	—	S.A.	—	—	N.S.W.	—	—	B.fl.VI,30	
P. Bowmanni, F. v. M. in Benth. Fl. Austr. VI, 30 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,30	M.fr.VII,6.
P. ammocharis, F. v. M. in Hook. Kew Misc. IX, 24 (1857) ...	—	—	—	—	—	—	N.A.	B.fl.VI,30	M.fr.VII,5;XI,47.
P. curviflora, R. Brown, prodr. 362 (1810))	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,31	M.fr.VII,6;VIII,9.
P. hirsuta, Meissner in De Cand. prodr. XIV, 513 (1856) ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,32	
P. altior, F. v. M., fragm. I, 84 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,32	M.fr.VII,7.
P. octophylla, R. Brown, prodr. 361 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,32	M.fr.VII,6.

P. petraea, Meissner in Schlecht. Linnaea XXVI, 347 (1853) ... — S.A. — — N.S.W. — — B.fl.VI,33
P. phylicoides, Meissner in Lehm. pl. Preiss. II, 271 (1847) ... — S.A. — V. N.S.W. — — B.fl.VI,33 M.fr.VII,6.
P. Eyrei, F. v. M., fragm. V, 109 (1865) ... ... W.A. — — — — — — B.fl.VI,34 M.fr.V,109.
P. longiflora, R. Brown, prodr. 361 (1810)... ... ... W.A. — — — — — — B.fl.VI,34 M.fr.VII,5.
P. stricta, Meissner in Schlecht. Linnaea XXVI, 348 (1853) ... — S.A. T. V. — — — B.fl.VI,34 M.fr.VII,4.
P. Preissii, Meissner in Lehm. pl. Preiss. I, 601 (1845) ... ... W.A. — — — — — — B.fl.VI,35 M.fr.VII,3.
P. Holroydi, F. v. M., fragm. VI, 159, t. 59 (1868) ... ... — — — — — — N.A. B.fl.VI,35 M.fr.250.

DRAPETES, Lamarck, Journ. d'hist. nat. I, 119, t. 10 (1792). (Kelleria, Daphnobryon).
D. Tasmanica, J. Hooker in Kew Misc. V, 299, t. 7 (1853) ... — — T. V. N.S.W. — — B.fl.VI,36 M.fr.VII,1.

WICKSTROEMIA, Endlicher, prodr. fl. insul. Norfolk. 47 (1833). (Daphne partly).
W. Indica, C. A. Meyer in Bull. de l'Acad. de Petersb. I, 357 (1843) — — — — N.S.W. Q. N.A. B.fl.VI,37 M.fr.VII,1.

PHALERIA, Jack in Hook. Comp. to Bot. Mag. I, 156 (1835). (Drymispermum).
P. Blumei, Bentham, Fl. Austr. VI, 38 (1873) ... ... — — — — — Q. — B.fl.VI,38
P. Neumanni, F. v. M., fragm. VIII, 9 (1873) ... ... — — — — — Q. — B.fl.VI,38 M.fr.V,26.
P. Clerodendron, F. v. M., fragm. VIII, 9 (1873) ... ... — — — — — Q. — B.fl.VI,38 M.fr.VII,1.

CORNACEAE.

Humboldt, Bonpland & Kunth, nov. gen. Americ. III, 430 (1818).

STYLIDIUM, Loureiro, Fl. Cochinch. I, 220 (1790). (Marlea, Rhytidandra, Pseudalangium).
S. Vitiense, F.v.M.; Rhytid., Asa Gray, Bot. Wilk. Ex. I, 303, t. 28 (1854) — — — — N.S.W. Q. — B.fl.III,386

RUBIACEAE.

A. L. de Jussieu, gen. plant. 196 (1789), from B. de Jussieu (1759).

SARCOCEPHALUS, Afzelius in Transact. hort. soc. Lond. V, 422, t. 18 (1824).
S. cordatus, Miquel, Fl. Ind. Batav. II, 133 (1856) ... ... — — — — — Q. N.A. B.fl.III,492

OLDENLANDIA, Plumier, nov. plant. Americ. gen. 42, t. 36 (1703). (Hedyotis, Synaptantha).
O. corymbosa, Linné, sp. pl. 119 (1753) ... ... — — — — — Q. — — M.fr.VII,45.
O. paniculata, Linné, spec. plant. edit. sec. 1667 (1763) ... ... — — — — — Q. — — M.fr.VIII,47.
O. auricularia, F. v. M.; Hedyotis, Linné, spec. plant. 101 (1753) — — — — — Q. — B.fl.III,404
O. polyclada; Hedyotis, F. v. M., fragm. VIII, 146 (1874) ... — — — — — Q. — — M.fr.VIII,146.
O. coerulescens; Hedyotis, F. v. M., fragm. IV, 38 (1863) ... — — — — — Q. — B.fl.III,404 M.fr.IV,38.
O. spermacocoides; Hedyotis, F. v. M., fragm. VIII, 146 (1874) — — — — — — N.A. — M.fr.VIII,146.
O. Crouchiana; Hedyotis, F. v. M., fragm. X, 85 (1876) ... — — — — — — N.A. — M.fr.X,85.
O. mitrasacmoides; Hedyotis, F. v. M., fragm. IV, 37 (1863) ... — — — — — — N.A. B.fl.III,404 M.fr.IV,37.
O. galioides; Hedyotis, F. v. M., fragm. IV, 38 (1863) ... ... — — — — — Q. N.A. B.fl.III,404 M.fr.IV,38.
O. scleranthoides; Hedyotis, F. v. M., fragm. IV, 39 (1863) ... — — — — — — N.A. B.fl.III,405 M.fr.IV,39.
O. elatinoides, F. v. M.; Hedyotis, Benth., Fl. Austr. III, 405 (1866) W.A. — — — — — — B.fl.III,405 M.fr.VIII,147.
O. tillaeacea; Hedyotis, F. v. M., fragm. IV, 39 (1863) ... — S.A. — — N.S.W. Q. — B.fl.III,405 M.fr.IV,39.
O. trachymenioides; Hedyotis, F. v. M., fragm. IV, 40 (1863) ... — — — — — Q. — B.fl.III,406 M.fr.IV,40.
O. pterospora; Hedyotis, F. v. M., fragm. IV, 40 (1863)... ... — — — — — — N.A. B.fl.III,407 M.fr.IV,40,177.

DENTELLA, R. et G. Forster, char. gen. 25, t. 13 (1776). (Lippaya).
D. repens, R. & G. Forster, char. gen. 25, t. 13 (1776) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.III,407 M.fr.IX,187.

OPHIORRHIZA, Linné, fl. Zeylan. 190 et 239 (1747).
O. Australiana, Bentham, Fl. Austr. III, 407 (1866) ... ... — — — — — Q. — B.fl.III,408 M.fr.IX,187.

ABBOTTIA, F. v. M., fragm. IX, 181 (1875).
A. singularis, F. v. M., fragm. IX, 181 (1875) ... ... — — — — — Q. — — M.fr.IX,181.

GARDENIA, Ellis in Phil. Transact. LI, 935, t. 23 (1761).
G. edulis, F. v. M., fragm. I, 54 (1858) ... ... — — — — — — N.A. B.fl.III,408 M.fr.I,54.
G. resinosa, F. v. M., fragm. I, 54 (1858) ... ... — — — — — — N.A. B.fl.III,408 M.fr.I,54.
G. pyriformis, Cunningham in Benth. Fl. Austr. III, 409 (1866) — — — — — — N.A. B.fl.III,409
G. megasperma, F. v. M., fragm. I, 54 (1858) ... ... — — — — — — N.A. B.fl.III,409 M.fr.I,54.
G. Macgillivraei, Bentham, Fl. Austr. III, 409 (1866) ... ... — — — — — Q. — B.fl.III,409 M.fr.VII,46.
G. ochreata, F. v. M., fragm. I, 55 (1858) ... ... — — — — — Q. — B.fl.III,409 M.fr.155;VII,46.
G. suffruticosa, R. Brown in Benth. Fl. Austr. III, 410 (1866)... — — — — — — N.A. B.fl.III,410
G. fucata, R. Brown in Benth. Fl. Austr. III, 410 (1866) ... — — — — — — N.A. B.fl.III,410
G. Jardinei, F. v. M. in Benth. Fl. Austr. III, 410 (1866) ... — — — — — Q. — B.fl.III,410 M.fr.VII,46.

RANDIA, Houston in Linné, hort. Cliffort. 485 (1737).
R. hirta, F. v. M., fragm. VII, 46 (1869) ... ... — — — — — Q. — — M.fr.IX,181.
R. chartacea, F. v. M., fragm. IX, 180 (1875) ... ... — — — — N.S.W. Q. — B.fl.III,410 M.fr.VII,47.
R. Benthamiana, F. v. M., fragm. IX, 180 (1875) ... ... — — — — N.S.W. Q. — — M.fr.IX,180.
R. Moorei, F. v. M. in Benth. Fl. Austr. III, 411 (1866) ... — — — — N.S.W. Q. — B.fl.III,411
R. stipularis, F. v. M., Papuan plants I, 69 (1876) ... ... — — — — N.S.W. — — — M.fr.VII,47;IX,70,181.
R. Fitzalani, F. v. M., Rep. Burdek. Exped. 12 (1860) ... ... — — — — — Q. — B.fl.III,411 M.fr.V,19;IX,180.
R. sessilis, F. v. M., fragm. VII, 47 (1869) ... ... — — — — — Q. N.A. — M.fr.VII,47. [180.
R. densiflora, Bentham, Fl. Hongk. 155 (1861) ... ... — — — — N.S.W. Q. N.A. B.fl.III,412 M.fr.II,132; III,166; IX,

DIPLOSPORA, De Candolle, prodr. IV, 477 (1830). (Discospermum.)
D. Australis, Bentham, Fl. Austr. III, 413 (1866) ... ... — — — — — Q. — B.fl.III,413
D. ixoroides, F. v. M., fragm. IX, 182 (1875) ... ... — — — — — Q. — B.fl.III,416 M.fr.IX,182.

IXORA, Linné, syst. nat. 8 (1735); Linné, gen. pl. 27 (1737). (Pavetta, Webera, Stylocorne.)
I. Pavetta, Roxburgh, Fl. Ind. I, 395, ed. Carey (1820)... ... — — — — — Q. N.A. B.fl.III,414 M.fr.IX,182.
I. tomentosa, Roxburgh, Fl. Ind. I, 396, ed. Carey (1820) ... — — — — — — N.A. B.fl.III,414
I. coccinea, Linné, sp. pl. 110 (1753) ... ... — — — — — — N.A. B.fl.III,415
I. Timorensis, Decaisne in Nouv. Ann. du Mus. III, 419 (1834)... — — — — — Q. N.A. B.fl.III,415 M.fr.V,18;IX,183.
I. Becklerii, Bentham, Fl. Austr. III, 415 (1866)... ... ... — — — — N.S.W. Q. — B.fl.III,415 M.fr.IX,183.

I. pentamera, Bentham, Fl. Austr. III, 416 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,416				
I. Dallachyana; F. v. M., Webera, Benth., Fl. Austr. III, 412 (1866)	—	—	—	—	—	Q.	—	B.fl.III,412				
I. expandens; Webera, F. v. M., fragm. VI, 25 (1867)	—	—	—	—	—	—	N.A.	—	M.fr.VI,25.			
COFFEA, Linné, syst. nat. 8 (1735); Linné, gen. pl. 55 (1737), from Ray (1691).												
C. Bengalensis, Roxburgh, hort. Bengal. 15 (1814)	—	—	—	—	—	Q.	—	—	M.fr.VIII,147.			
TIMONIUS, Rumphius, herb. Amboin. III, 216, t. 140 (1743). (Polyphragmon.)												
T. Rumphii, De Candolle, prodr. IV, 461 (1830)	—	—	—	—	—	Q.	N.A.	B.fl.III,417	M.fr.II,134;IX,187.			
SCYPHIPHORA, K. F. Gaertner, de fruct. III, 91, t. 196 (1805). (Epithinia.)												
S. hydrophylacea, K. F. Gaertner, de fruct. III, 91, t 196 (1805)	—	—	—	—	—	Q.	N.A.	B.fl.III,418	M.fr.VII,46;IX,187.			
GUETTARDA, Osbeck in Linné, spec. plant. 991 (1753). (Antirrhoea, Guettardella.)												
G. speciosa, Linné, spec. plant. 991 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.III,419	M.fr.IX,183.			
G. tenuiflora, F. v. M. fragm. VII, 48 (1869)	—	—	—	—	—	Q.	—	B.fl.III,418	M.fr.IX,183.			
G. putaminosa, F. v. M., fragm. IX, 183 (1875)	—	—	—	—	—	Q.	—	B.fl.III,419	M.fr.IV,92.			
G. myrtoides, F. v. M., fragm. IX, 184 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,184.			
HODGKINSONIA, F. v. M., fragm. II, 132 (1861).												
H. ovatiflora, F. v. M., fragm. II, 132 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,420	M.fr.III,166.			
CANTHIUM, Lamarck, Encycl. method. I, 602 (1783). (Psydrax, Plectronia partly.)												
C. cymosum, Persoon, synops. plant. I, 200 (1805)	—	—	—	—	—	Q.	—	—	M.fr.IX,185.			
C. latifolium, F. v. M. in Benth. Fl. Austr. III, 421 (1866) ...	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,421				
C. attenuatum, R. Brown in Benth. Fl. Austr. III, 421 (1866) ...	—	—	—	—	—	—	N.A.	B.fl.III,421				[135.
C. lucidum, Hooker & Arnott, Bot. of Beechey's voy. 65 (1841)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,421	M.fr.II,133; IX,185; XI,			
C. oleifolium, Hooker in Mitch. Trop. Austr. 397 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,422				
C. buxifolium, Bentham, Fl. Austr. III, 422 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,422				
C. vacciniifolium, F. v. M. in Trans. phil. Inst. Vict. III, 47 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III.422	M.fr.II,134;IX,186.			
C. coprosmoides, F. v. M. in Trans. phil. Inst. Vict. III, 47 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,422				
MORINDA, Vaillant in act. Acad. Par. 202 (1722).												
M. citrifolia, Linné, spec. plant. 176 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.III,422				
M. acutifolia, F. v. M., fragm. IX, 179 (1875)	—	—	—	—	—	Q.	—	B.fl.III,421	M.fr.IX,179.			
M. jasminoides, Cunningham in Hook. Bot. Mag. t. 3351 (1834)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,423				
M. umbellata, Linné, spec. plant. 176 (1753)	—	—	—	—	—	Q.	—	B.fl.III,423				
M. reticulata, Bentham, Fl. Austr. III, 424 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,423	M.fr.IX,180.			
COELOSPERMUM, Blume, Bijdr. 994 (1826). (Pogonolobus.)												
C. paniculatum, F. v. M., fragm. V, 19 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.III.425	M.fr.VII,46;IX,185.			
C. reticulatum, Bentham, Fl. Austr. III, 425 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,425	M.fr.I,56;IX,185.			
HYDNOPHYTUM, Jack in Transact. Linn. Soc. XIV, 124 (1823).												
H. formicarum, Jack in Transact. Linn. Soc. XIV, 124 (1823)...	—	—	—	—	—	Q.	N.A.	—	M.fr.VII,45.			
MYRMECODIA, Jack in Transact. Linn. Soc. XIV, 122 (1823).												
M. echinata, Gaudichaud, Bot. voy. Bonite 472, t. 96 (1844) ...	—	—	—	—	—	Q.	N.A.	—	M.fr.VII,45.			
LASIANTHUS, Jack in Transact. Linn. Soc. XIV, 125 (1823).												
L. strigosus, Wight in Calc. Journ. nat. hist. VI, 512 (1846) ...	—	—	—	—	—	Q.	—	B.fl.III,426				
PSYCHOTRIA, Linné, syst. ed. decim. 929 (1759). (Uragoga, 1737, Grumilia.)												
P. nesophila, F. v. M., fragm. II, 135 (1861)	—	—	—	—	—	Q.	N.A.	B.fl.III.427	M.fr.IX,184.			
P. Dallachyana, Bentham, Fl. Austr. III, 427 (1866)	—	—	—	—	—	Q.	—	B.fl.III,427				
P. nematopoda, F. v. M., fragm. VII, 48 (1869)	—	—	—	—	—	Q.	—	—	M.fr.IX,184.			
P. loniceroides, Sieber in De Cand., prodr. IV, 523 (1830) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,427	M.fr.IX,184.			
P. daphnoides, Cunningham in Hook. Bot. Mag. 3228 (1833) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,428	M.fr.IX,184.			
P. poliostemma, Bentham, Fl. Austr. III. 427 (1866)	—	—	—	—	—	Q.	—	B.fl.III,428				
P. Fitzalani, Bentham, Fl. Austr. III, 427 (1866)	—	—	—	—	—	Q.	—	B.fl.III,428				
P. Carronis, Moore & F. v. M., fragm. VII, 49 (1869)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,70,185.			
GEOPHILA, D. Don, prodr. fl. Nepal. 136 (1825). (Cephaelis partly.)												
G. reniformis, D. Don, prodr. fl. Nepal. 136 (1825)	—	—	—	—	—	Q.	—	—	M.fr.VII,45;VIII,147.			
COPROSMA. R. et G. Forster, char. gen. 137, t. 69 (1776). (Nertera, Marquisia.)												
C. Nertera, F. v. M., fragm. IX, 186 (1875)	—	—	T.	V.	N.S.W.	—	—	—	M.fr.IX,186.			
C. reptans, F. v. M., fragm. IX, 186 (1875)	—	—	—	V.	—	—	—	—	M.fr.IX,186.			
C. pumila, J. Hooker in Lond. Journ. VI, 465 (1847)	—	—	T.	V.	—	—	—	B.fl.III,430	M.fr.IX,186.			
C. nitida, J. Hooker in Lond. Journ. VI, 465 (1847)	—	—	T.	V.	—	—	—	B.fl.III,430	M.fr.IX,186.			
C. Billardieri, J. Hooker in Lond. Journ. VI, 465 (1847) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,430				
C. hirtella, Labillardière, Nov. Holl. pl. spec. I, 70, t. 95 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,429	M.fr.IX,186.			
C. Baueri, Endlicher, iconogr. gen. pl. XI, t. 111 (1838) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,69.			
C. putida, C. Moore & F. v. M., fragm. VII, 45 (1869)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,69.			
C. lanceolaris, F. v. M., fragm. IX, 70 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,70.			
OPERCULARIA, J. Gaertner, de fruct. I, 111, t. 24 (1788).												
O. vaginata, Labillardière, Nov. Holl. pl. spec. I, 34, t. 46 (1804)	W.A.	—	—	—	—	—	—	B.fl.III,433				
O. spermacocea, Labillard., Nov. Holl. pl. spec. I, 35, t. 47 (1804)	W.A.	—	—	—	—	—	—	B.fl.III,433				
O. scabrida, Schlechtendal, Linnaea XX, 604 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,433				
O. hirsuta, F. v. M. in Benth. Fl. Austr. III, 434 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,434				
O. aspera, Gaertner, de fruct. I, 112, t. 24 (1788)...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,434				
O. hispida, Sprengel, syst. veg. I, 385 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.III,434				
O. diphylla, Gaertner, de fruct. I, 113 (1788)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,434				
O. ovata, J. Hooker in Lond. Journ. VI, 465 (1847)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,435				
O. varia, J. Hooker in Lond. Journ. VI, 466 (1847)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,435				
O. rubioides, A. L. de Jussieu in Annal. du Mus. IV, 428 (1804)	W.A.	—	—	—	—	—	—	B.fl.III,435				
O. volubilis, R. Brown in Benth. Fl. Austr. III, 435 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,435				
O. hispidula, Endlicher in Hueg. enum. 58 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,436				
O. echinocephala, Bentham, Fl. Austr. III, 436 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,436				
O. apiciflora, Labillardière, Nov. Holl. pl. spec. I, 35, t. 48 (1804)	W.A.	—	—	—	-	—	—	B.fl.III,436				

POMAX, Solander in Gaertn., de fruct. I, 112 (1788).

P. umbellata, Solander in Gaertn., de fruct. I, 112 (1788) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,437	M.fr.IX,187.

ELEUTHRANTHES, F. v. M., fragm. IV, 92 (1864).

E. opercularina, F. v. M., fragm. IV, 92 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,437	M.fr.IV,92.

KNOXIA, Linné, Fl. Zeyl. 189 (1747).

K. corymbosa, Willdenow, spec. pl. I, 582 (1798)	—	—	—	—	—	Q.	N.A.	B.fl.III,438	M.fr.IX,187.

SPERMACOCE, Dillenius, hort. Elth. 369, t. 277 (1732).

S. brachystoma, R. Brown in Benth. Fl. Austr. III, 439 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,439	M.fr.IV,41.
S. pogostoma, Bentham, Fl. Austr. III, 440 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,440	
S. leptoloba, Bentham, Fl. Austr. III, 440 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,440	
S. marginata, Bentham, Fl. Austr. III, 440 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,440	
S. multicaulis, Bentham, Fl. Austr. III, 440 (1866)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,440	
S. exserta, Bentham, Fl. Austr. III, 441 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,441	
S. membranacea, R. Brown in Benth. Fl. Austr. III, 441 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,441	
S. debilis, Bentham, Fl. Austr. III, 441 (1866)	—	—	—	—	—	Q.	—	B.fl.III,441	
S. inaperta, F. v. M., fragm. IV, 43 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,441	M.fr.IV,43.
S. stenophylla, F. v. M., fragm. IV, 43 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,442	M.fr.IV,43.
S. laevigata, F. v. M., fragm. IV, 41 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,442	M.fr.IV,41.
S. breviflora, F. v. M. in Benth. Fl. Austr. III, 442 (1866) ...	—	—	—	—	—	—	N.A.	B.fl.III,442	
S. auriculata, F. v. M., fragm. IV, 42 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,442	M.fr.IV,42.
S. suffruticosa, R. Brown in Benth. Fl. Austr. III, 443 (1866) ...	—	—	—	—	—	—	N.A.	B.fl.III,443	

ASPERULA, Dodoens, pemptad. 355 (1583).

A. geminifolia, F. v. M., fragm. V, 147 (1865)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,443	M.fr.IX,188;XI,27.
A. oligantha, F. v. M. in Neerl. Kruidk. Arch. IV, 111 et 112 (1859)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,444	M.fr.IX,187;XI,27.

GALIUM, Dodoens, pemptad. 335 (1583), from Dioscorides.

G. umbrosum, Solander in G. Forst. prodr. 89 (1786)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,446	M.fr.IX,188.
G. australe, De Candolle, prodr. IV, 608 (1830)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,446	

CAPRIFOLIACEAE.

Adanson, Familles des plantes II, 133 (1763).

SAMBUCUS, Tournefort, inst. 606, t. 396 (1700), from Dodoens (1583). (Tripetelus.)

S. xanthocarpa, F. v. M. in Transact. phil. Inst. Vict. I, 42 (1855)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,398	
S. Gaudichaudiana, De Candolle, prodr. IV, 322 (1830)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,398	

PASSIFLOREAE.

A. L. de Jussieu in Ann. du Mus. VI, 102 (1805).

PASSIFLORA, Plukenet, phytograph. 202 et 282, t. 104, 210, 211, 212 (1692). (Disemma.)

P. Herbertiana, Lindley, Bot. Regist. t. 737 (1823)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,311	M.fr.IX,68.
P. cinnabarina, Lindley in Gard. Chron. 724 (1855)	—	—	—	V.	N.S.W.	—	—	—	M.fr.IX,68.
P. aurantia, G. Forster, fl. ins. Austr. prodr. 62 (1786)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,312	M.fr.IX,68.
P. brachystephanea, F. v. M. in Benth. Fl. Austr. III, 312 (1866)	—	—	—	—	—	Q.	—	B.fl.III,312	M.fr.I.56.

MODECCA, Lamarck, Encycl. méthod. IV, 208 (1797).

M. australis, R. Brown in De Cand. prodr. III, 337 (1828) ...	—	—	—	—	—	Q.	N.A.	B.fl.III,313	M.fr.IX,69.

CUCURBITACEAE.

Haller, enum. stirp. Helv. praef. 34 (1742).

TRICHOSANTHES, Linné, gen. plant. 295 (1737).

T. pentaphylla, F. v. M. in Benth. Fl. Austr. III, 314 (1866) ...	—	—	—	—	—	Q.	—	B.fl.III,314	
T. cucumerina, Linné, spec. plant. 1008 (1753)	—	—	—	—	—	—	N.A.	B.fl.III,314	
T. palmata, Roxburgh, Fl. Ind. III, 704 (1832)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,315	M.fr.VI,187.
T. Hearnii, F. v. M. in Benth. Fl. Austr. III, 315 (1866) ...	—	—	—	—	—	Q.	—	B.fl.III,315	

LAGENARIA, Seringe in Mém. Soc. Genèv. III, 25, t. 2 (1825).

L. vulgaris, Seringe in De Cand. prodr. III, 279 (1828)	—	—	—	—	—	Q.	—	B.fl.III,316	M.fr.VII,62.

LUFFA, Tournefort in act. Acad. Paris. 84, t. 2 (1706), from Vesling (1638).

L. Aegyptiaca, Miller, Gardener's Dictionary (1731)	—	—	—	—	—	Q.	—	B.fl.III,316	M.fr.III,107.
L. graveolens, Roxburgh, Fl. Ind. III, 716 (1832)	—	—	—	—	—	—	N.A.	B.fl.III,316	M.fr.III,106;IV,175.
L. foetida, Cavanilles, icon. I, 7, t. 9—10 (1791)	—	—	—	—	—	—	N.A.	—	M.fr.XI,138.

ZANONIA, Linné, coroll. 19 (1737). (Alsomitra.)

Z. Capricornica, F. v. M., fragm. VII, 61 (1870)	—	—	—	—	—	Q.	—	—	M.fr.VII,61.
Z. Hookeri, F. v. M., fragm. VI, 188 (1868)	—	—	—	—	—	Q.	—	—	M.fr.V,181;VII,62.
Z. Stephensiana, F. v. M., fragm. VIII, 181 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,181.

CUCUMIS, Tournefort, inst. 104, t. 31 (1700), from Plinius.

C. trigonus, Roxburgh, Fl. Ind. III, 722 (1832)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,317	M.fr.VI,187.
C. Chate, Linné, syst. nat. edit. decim. 5 (1759)	—	S.A.	—	—	N.S.W.	Q.	N.A.	—	

BENINCASA, Savi in Bibl. Ital. IX, 158 (1818).

B. vacua; Cucurbita vacua, F. v. M., fragm. VI, 186 (1868) ...	—	—	—	—	—	Q.	—	—	M.fr.VI,186.

MOMORDICA, Tournefort, institut. rei herbar. 103, t. 29 et 30 (1700).

M. Balsamina, Linné, sp. pl. 1009 (1753)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,318	M.fr.VI,187.
M. Charantia, Linné, sp. pl. 1009 (1753)	—	—	—	—	—	Q.	—	—	

BRYONOPSIS, Arnott in Hook. Journ. of Bot. III, 274 (1841). (Bryonia partly.)

B. laciniosa, Naudin in Annal. des sc. nat. sér. cinq. VI, 30 (1869)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,319	
B. Pancheri, Naudin in Annal. des sc. nat. sér. cinq. V, 30 (1869)	—	—	—	—	N.S.W.	—	—	—	

MELOTHRIA, Linné, coroll. 1 (1737). (Zehneria, Mukia.)

M. Cunninghamii, Bentham, Fl. Austr. III, 320 (1866)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,320	M.fr.VI,188;VII,62.
M. Baueriana, F. v. M., fragm. VI, 188 (1868)	—	—	—	—	N.S.W.	—	—	—	M.fr.VI,188.
M. Muelleri, Bentham, Fl. Austr. III, 320 (1866)...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,320	
M. Maderaspatana, Cogniaux in De Cand. mon. phan. III, 623 (1881)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,321	M.fr.VI,187.

SICYOS, Linné, syst. nat. 9 (1735); Linné, gen. plant. 297 (1737).

S. angulata, Linné, spec. plant. 1013 (1753)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,322	M.fr.VI,187;VII,62.

COMPOSITAE.

Vaillant in act. Acad. Paris. 143 (1718).

CENTRATHERUM, Cassini in Bull. de la Soc. philom. (1817).

C. muticum, Lessing in Schlecht. Linnaea II, 320 (1829) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,460	

PLEUROCARPAEA, Bentham, Fl. Austr. III, 460 (1866).

P. denticulata, Bentham, Fl. Austr. III, 460 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,460	

VERNONIA, Schreber, gen. plant. II, 541 (1791).

V. cinerea, Lessing in Schlecht. Linnaea II, 291 (1829)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,459	
V. vagans, De Candolle, prodr. V, 32 (1843)	—	—	—	—	—	Q.	—	—	M.fr.VI,234.

ELEPHANTOPUS, Vaillant in act. Acad. Par. 309 (1719).

E. scaber, Linné, spec. plant. 814 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.III,461	

ADENOSTEMMA, R. et G. Forster, char. gen. 89, t. 45 (1776).

A. viscosum, R. & G. Forster, char. gen. 89, t. 45 (1776) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,462	

AGERATUM, Linné, gen. plant. 247 (1737).

A. conyzoides, Linné, spec. plant. 839 (1753)	—	—	—	—	—	Q.	—	B.fl.III,462	

EUPATORIUM, Tournefort, inst. 455, t. 259 (1700), from C. Bauhin (1623).

E. cannabinum, Linné, spec. pl. 838 (1753)...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,462	M.fr.V,62.

LAGENOPHORA, Cassini in Bull. de la Soc. philom. 34 (1818). (Ixauchenus, Solenogyne, Emphysopus.)

L. Billardieri, Cassini in Diction. XXV, 111 (1822)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,507	
L. Huegelii, Bentham in Hueg. enum. 59 (1837)	W.A.	S.A.	T.	V.	—	—	—	B.fl.III,507	
L. Solenogyne, F. v. M., fragm. V, 62 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,508	M.fr.V,62.
L. Emphysopus, J. Hooker, Fl. Tasman. I, 189 (1860)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,508	

BRACHYCOME, Cassini in Bull. de la Soc. philom. 199 (1816). (Bellidis subgenus, Pacquerina, Brachystephium, Steiroglossa, Silphiospermum.)

B. diversifolia, Fischer & Meyer, ind. sem. hort. Petrop. 31 (1835)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,511	
B. segmentosa, Moore & F. v. M., fragm. VIII, 144 (1874) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,144.
B. melanocarpa, Sonder & F. v. M. in Schl. Linn. XXV, 476 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,511	
B. radicans, Steetz in Lehm. pl. Preiss. I, 429 (1845)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,512	
B. goniocarpa, Sonder & F. v. M. in Schl. Linn. XXV, 474 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,512	
B. pachyptera, Turczaninow in Bull. de Mosc. XXIV, 175 (1851)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,512	
B. iberidifolia, Bentham in Hueg. enum. 59 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,512	M.fr.XI,123.
B. pusilla, Steetz in Lehm. pl. Preiss. I, 427 (1845)	W.A.	—	—	—	—	—	—	B.fl.III,513	
B. microcarpa, F. v. M., fragm. I, 50 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,513	M.fr.I,50.
B. Stuartii, Bentham, Fl. Austr. III, 513 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,513	
B. scapigera, De Candolle, prodr. VII, 277 (1838)...	—	—	—	V.	N.S.W.	—	—	B.fl.III,513	
B. Muelleri, Sonder in Schlecht. Linnaea XXV, 475 (1852) ...	—	S.A.	—	—	—	—	—	B.fl.III,514	
B. graminea, F. v. M., fragm. I, 49 (1858)...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,514	M.fr.VIII,142.
B. angustifolia, Cunningham in De Cand., prodr. V, 306 (1836)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,514	M.fr.I,49.
B. linearifolia, De Candolle, prodr. V, 306 (1836)...	—	—	—	—	N.S.W.	—	—	B.fl.III,515	
B. basaltica, F. v. M., fragm. I, 50 (1858)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,515	M.fr.I,50.
B. trachycarpa, F. v. M. in Schlecht. Linnaea XXV, 339 (1852)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,515	
B. exilis, Sonder in Linnaea XXV, 473 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,516	
B. ptychocarpa, F. v. M. in Transact. phil. Soc. Vict. I, 43 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.III,516	
B. debilis, Sonder in Linnaea XXV, 477 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,516	
B. decipiens, J. Hooker in Lond. Journ. VI, 114 (1847)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,516	
B. cardiocarpa, F. v. M. in Benth. Fl. Austr. III, 516 (1866) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,517	
B. nivalis, F. v. M. in Transact. phil. Soc. Vict. I, 43 (1854) ...	—	—	—	V.	N.S.W.	—	—	B.fl.III,517	
B. scapiformis, De Candolle, prodr. V, 306 (1836)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,517	
B. stricta, De Candolle, prodr. V, 305 (1836)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,518	
B. heterodonta, De Candolle, prodr. V, 305 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,518	
B. Billardieri, Bentham, Fl. Austr. III, 518 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,518	
B. cheilocarpa, F. v. M. in Wing's South. Sc. Record, Aug. (1882)	W.A.	—	—	—	—	—	—	—	
B. ciliaris, Lessing, syn. comp. 172 (1832)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,518	
B. calocarpa, F. v. M. in Schlecht. Linnaea XXV, 399 (1852) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,519	M.fr.XI,27.
B. chrysoglossa, F. v. M. in Transact. phil. Soc. Vict. I, 44 (1854)	—	S.A.	—	V.	N.S.W.	—	—	—	
B. marginata, Bentham in Hueg. enum. 60 (1837)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,519	
B. Sieberi, De Candolle, prodr. V, 306 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,520	
B. discolor, C. Stuart in Benth. Fl. Austr. III, 520 (1866) ...	—	—	—	—	N.S.W.	—	—	B.fl.III,520	
B. multifida, De Candolle, prodr. V, 306 (1836)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,520	M.fr.VIII,145.
B. latisquamea, F. v. M., fragm. XI, 16 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,16.
B. glandulosa, Bentham, Fl. Austr. III, 520 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,520	
B. collina, Bentham, Fl. Austr. III, 521 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,521	

ERODIOPHYLLUM, F. v. M., fragm. IX, 119 (1875).

E. Elderi, F. v. M., fragm. IX, 120 (1875)...	—	S.A.	—	—	—	—	—	—	M.fr.IX,119.

MINURIA, De Candolle, prodr. V, 298 (1836). (Therogeron, Elachothamnus, Kippistia.)

M. leptophylla, De Candolle, prodr. V, 298 (1836)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,498	M.fr.IX,119;X,56.
M. Cunninghamii, Bentham, Fl. Austr. III, 498 (1866)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,498	

M. Candollei, F. v. M., fragm. IX, 119 (1875)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,499	M.fr.X,56.
M. suaedifolia, F. v. M. in Bentham, Fl. Austr. III, 499 (1866)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,499	M.fr.X,56.
CALOTIS, R. Brown in Bot. Regist. t. 504 (1820). (Huenefeldia, Goniopogon, Cheiroloma.)									
C. dentex, R. Brown in Bot. Regist. 504 (1820)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,501	
C. cuneifolia, R. Brown in Bot. Regist. t. 504 (1820)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,501	
C. glandulosa, F. v. M. in Transact. Vict. Inst. 129 (1855) ...	—	—	—	V.	N.S.W.	—	—	B.fl.III,502	
C. cymbacantha, F. v. M. in Schlecht. Linnaea XXV, 400 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,502	
C. erinacea, Steetz in Lehm. pl. Preiss. I, 424 (1845)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,502	
C. scabiosifolia, Sonder & F. v. M. in Linnaea XXV, 471 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,503	M.fr.XI,27.
C. scapigera, Hooker in Mitch. Trop. Austr. 75 (1848)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,503	M.fr.XI,27.
C. anthemoides, F. v. M. in Transact. phil. Soc. Vict. I, 44 (1854)	—	—	—	V.	—	—	—	B.fl.III,504	
C. lappulacea, Bentham in Hueg. enum. 60 (1837)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,504	
C. microcephala, Bentham, Fl. Austr. III, 504 (1866)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,504	
C. breviseta, Bentham in Hueg. enum. 60 (1837)	—	—	—	—	—	—	N.A.	B.fl.III,505	
C. plumulifera, F. v. M. in Transact. phil. Inst. Vict. III, 57 (1859)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,505	
C. porphyroglossa, F. v. M. in Benth. Fl. Austr. III, 505 (1866)	—	S.A.	—	—	N.S.W.	—	N.A.	B.fl.III,505	
C. pterosperma, R. Brown in Benth. Fl. Austr. III, 505 (1866)...	—	—	—	—	—	—	N.A.	B.fl.III,505	
C. hispidula, F. v. M. in Transact. Vict. Inst. 130 (1855) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,506	
C. Kempei, F. v. M. in Transact. Roy. Soc. S.Austr. IV, 112 (1881)—		S.A.	—	—	—	—	—	—	M.fr.XI,139.
ASTER, Tournefort, inst. 481, t. 274 (1700), from Dioscorides. (Olearia, Celmisia, Eurybia, Steetzia.)									
A. megalophyllus, F. v. M., fragm. V, 70 (1865)	—	—	—	V.	N.S.W.	—	—	B.fl.III,468	M.fr.V,70.
A. chrysophyllus, Cunningham in De Cand. prodr. V, 266 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,468	
A. alpicola, F. v. M., fragm. V, 70 (1865)	—	—	—	V.	N.S.W.	—	—	B.fl.III,468	M.fr.V,70.
A. rosmarinifolius, Cunningham in De Cand. prodr. V, 268 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,468	
A. viscosus, Labillardière, Nov. Holl. pl. spec. II, 53, t. 203 (1806)	—	—	T.	V.	—	—	—	B.fl.III,468	M.fr.V,71.
A. Sonderi, F. v. M., fragm. V, 83 (1865)	—	S.A.	—	—	—	—	—	B.fl.III,469	M.fr.V,83.
A. pannosus, F. v. M., fragm. V, 83 (1865)	—	S.A.	—	V.	—	—	—	B.fl.III,469	M.fr.V,83.
A. oliganthemus; Olearia, F. v. M. in Benth. Fl. Austr. III, 469 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,469	
A. argophyllus, Labillard., Nov. Holl. pl. spec. II, 52, t. 201 (1806)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,470	M.fr.V,68.
A. cydonifolius, Cunningham in De Cand. prodr. V, 267 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,470	
A. myrsinoides, Labillard., Nov. Holl. pl. spec. II, 53, t. 202 (1806)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,471	M.fr.V,69.
A. persoonioides, Cunningham in De Cand. prodr. V, 267 (1836)	—	—	T.	—	—	—	—	B.fl.III,471	M.fr.V,69.
A. obcordatus, F. v. M., fragm. V, 69 (1865)	—	—	T.	—	—	—	—	B.fl.III,471	M.fr.V,69.
A. pinifolius, F. v. M., fragm. V, 71 (1865)	—	—	T.	—	—	—	—	B.fl.III,472	M.fr.V,71.
A. ledifolius, Cunningham in De Cand. prodr. V, 269 (1836) ...	—	—	T.	—	—	—	—	B.fl.III,472	
A. dentatus, Andrews, Bot. Reposit. t. 61 (1799)...	—	—	—	—	N.S.W.	—	—	B.fl.III,472	M.fr.V,82.
A. stellulatus, Labillard., Nov. Holl. pl. spec. II, 50, t. 196 (1806)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,473	M.fr.V,81.
A. asterotrichus, F. v. M., fragm. V, 79 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,473	M.fr.I,111.
A. gravis, F. v. M., fragm. V, 82 (1865)	—	—	—	—	N.S.W.	—	—	B.fl.III,474	M.fr.V,82.
A. Nernstii, F. v. M., fragm. V, 81 (1865)...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,474	M.fr.V,81.
A. hygrophilus, Cunningham in De Cand. prodr. V, 269 (1836) ...	—	—	—	—	—	Q.	—	B.fl.III,474	
A. Siemssenii, F. v. M., fragm. V, 71 (1865)	—	—	—	—	N.S.W.	—	—	B.fl.III,475	M.fr.I,50.
A. tubuliflorus, F. v. M., fragm. V, 65 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,475	M.fr.V,65.
A. axillaris, F. v. M., fragm. V, 64 (1865)...	W.A.	S.A.	T.	V.	N.S.W.	—	N.A.	B.fl.III,475	M.fr.V,64.
A. revolutus; Olearia, F. v. M. in Benth. Fl. Austr. III, 476 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,476	
A. exilifolius, F. v. M., fragm. V, 69 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,476	M.fr.V,69.
A. ramulosus, Labillard., Nov. Holl. plant. spec. II, 51, t. 198 (1806)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,476	
A. florulentus, F. v. M., fragm. V, 82 (1865)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,477	M.fr.V,82.
A. microphyllus, Persoon, synops. pl. II, 442 (1807)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,478	M.fr.V,70.
A. Mitchelli, F. v. M., fragm. V, 78 (1865)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,478	M.fr.V,78.
A. exiguifolius, F. v. M., fragm. V, 67 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,478	M.fr.V,67.
A. Cassiniae, F. v. M., fragm. V, 68 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,479	M.fr.V,68.
A. cyanodiscus, F. v. M., fragm. V, 82 (1865)	—	—	—	—	N.S.W.	—	—	B.fl.III,479	M.fr.V,82.
A. pimeleoides, Cunningham in De Cand. prodr. V, 268 (1836)...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,479	
A. iodochrous, F. v. M., fragm. V, 81 (1865)	—	—	—	V.	N.S.W.	—	—	B.fl.III,480	M.fr.II,110.
A. adenolasius, F. v. M., fragm. V, 67 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,480	M.fr.V,67.
A. conocephalus, F. v. M., fragm. V, 79 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,480	M.fr.V,79.
A. calcareus; Olearia, F. v. M. in Benth. Fl. Austr. III, 481 (1866)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,481	
A. magniflorus, F. v. M., fragm. V, 80 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,481	M.fr.V,80.
A. Muelleri; Eurybia, Sonder in Schlecht. Linn. XXV, 459 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,481	
A. Stuartii, F. v. M., fragm. V, 76 (1865)	—	S.A.	—	—	—	—	—	B.fl.III,481	M.fr.I,202.
A. decurrens, Cunningham in De Cand. prodr. V, 269 (1836) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,481	
A. glutescens, F. v. M., fragm. V, 77 (1865)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,482	M.fr.V,77.
A. orarius, F. v. M., fragm. V, 78 (1865)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,482	M.fr.V,78.
A. vernicosus, F. v. M., fragm. V, 67 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,482	M.fr.V,67.
A. teretifolius, F. v. M., fragm. V, 77 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,483	M.fr.V,77.
A. Turczaninowii, F. v. M., fragm. V, 67 (1865)...	W.A.	—	—	—	—	—	—	B.fl.III,483	M.fr.V,67.
A. ellipticus, Cunningham in De Cand. prodr. V, 271 (1836) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,483	M.fr.I,16.
A. glandulosus, Labillard., Nov. Holl. pl. spec. II, 50, t. 197 (1806)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,484	
A. heleophilus, F. v. M., fragm. V, 66 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,484	M.fr.V,66.
A. muricatus, F. v. M., fragm. V, 66 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,484	M.fr.V,66.
A. Benthami, F. v. M.; Olcariastricta, Benth. Fl. Austr. III, 485 (1866)	—	—	—	V.	—	—	—	B.fl.III,485	
A. Steetzii, F. v. M., fragm. V, 66 (1865)	W.A.	—	—	—	—		—	B.fl.III,485	M.fr.V,66.
A. Cunninghami, F. v. M.; Eurybia, De Cand. prodr. V, 269 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,486	
A. paucidentatus, F. v. M., fragm. V, 66 (1865)	W.A.	—	—	—	—		—	B.fl.III,485	M.fr.V,66.
A. megalodontus, F. v. M., fragm. VIII, 244 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,244.
A. adenophorus, F. v. M., fragm. V, 78 (1865)	—	—	—	V.	N.S.W.	—	—	B.fl.III,486	M.fr.I,111.
A. homolepis, F. v. M., fragm. V, 65 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,486	M.fr.V,65.
A. xerophilus, F. v. M., fragm. V, 76 (1865)	—	—	—	—	—	Q.	—	B.fl.III,486	M.fr.I,51,76&86.
A. Ferresii, F. v. M., fragm. V, 75 (1865)	—	S.A.	—	—	—	—	N.A.	B.fl.III,487	M.fr.III,18.

A. exul, Lindley, Bot. Regist. XXV, App. XXIV (1839) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,487	M.fr.V,75.
A. argutus, R. Brown in Benth. Fl. Austr. III, 488 (1866) ...	—	—	—	—	—	—	N.A.	B.fl.III,488	
A. Huegelii, F. v. M., fragm. V, 79 (1865)...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,488	M.fr.V,79.
A. Balli, F. v. M., fragm. VIII, 143 (1874)...	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,143.
A. Mooneyi, F. v. M., fragm. VIII, 144 (1874)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,144.
A. Celmisia, F. v. M., fragm. V, 84 (1865)...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,489	M.fr.V,84.
VITTADINIA, Ach. Richard, Voy. d'Astrol. bot. 250 (1832). (Eurybiopsis, Microgyne.)									
V. brachycomoides, F. v. M., fragm. V, 86 (1865)	—	—	—	—	—	Q.	N.A.	B.fl.III,490	M.fr.V,86.
V. australis, A. Richard, Voy. d'Astrol. 251 (1832)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,490	M.fr.XI,20.
V. scabra, De Candolle, prodr. V, 281 (1836)	—	—	—	—	N.S.W	Q.	—	B.fl.III,491	
V. macrorrhiza, A. Gray in Proc. Am. Acad. V, 118 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.III,492	
PODOCOMA, Cassini in Bull. de la Soc. philom. 137 (1817). (Ixiochlamys.)									
P. cuneifolia, R. Brown, App. to Sturt, Centr. Austr. 17 (1849)	—	S.A.	—	—	N.S.W.	—	N.A	B.fl.III,493	
ERIGERON, Linné, hort. Cliffort. 407 (1737). (Haplopappus partly.)									
E. pappochromus, Labillard., Nov. Holl. pl. spec. II, 47, t. 193 (1806)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,494	
E. ambiguus, F. v. M. in Transact. phil. Inst. Vict. III, 58 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.III,494	
E. sessilifolius, F. v. M., fragm. X, 100 (1881)	—	—	—	—	—	—	N.A.	—	M.fr.X,137.
E. minurioides, Bentham, Fl. Austr. III, 495 (1866)	—	—	—	V.	—	—	—	B.fl.III,495	
E. conyzoides, F. v. M. in Transact. phil. Soc. Vict. I, 105 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.III,495	
CONYZA, Lamarck. Encycl. méthod. II, 87, 89 (1786).									
C. viscidula, Wallich, numer. list 3006 (1828)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,496	
C. Aegyptiaca, Aiton, hort. Kew. III, 183 (1789)...	—	—	—	—	—	Q.	N.A.	B.fl.III,497	
BLUMEA, De Candolle in Guillemin, Arch. de Bot. II, 514 (1833).									
B. glandulosa, De Candolle in Wight, contrib. 14 (1834)... ...	—	—	—	—	—	Q.	N.A	B.fl.III,525	
B. amplectens, De Candolle in Wight, contrib. 13 (1834)... ...	—	—	—	—	—	Q.	—	B.fl.III,525	
B. integrifolia, De Candolle, prodr. V, 433 (1836)...	—	—	—	—	—	—	N.A.	B.fl.III,525	
B. diffusa, R. Brown in Benth. Fl. Austr. III, 525 (1866) ...	—	—	—	—	—	—	N.A.	B.fl.III,525	
B. hieracifolia, De Candolle in Wight, contrib. 15 (1834)... ...	—	—	—	—	—	Q.	N.A.	B.fl.III,526	
B. Cunninghamii, De Candolle, prodr. V, 435 (1836)	—	—	—	—	—	—	N.A.	B.fl.III,526	
B. acutata, De Cand. in Ann. du Mus. d'hist. nat. Par. III, 409 (1834)	—	—	—	—	—	Q.	—	—	M.fr.IX,160.
B. lacera, De Candolle in Wight, contrib. 14 (1834)	—	—	—	—	—	Q.	N.A	B.fl.III,526	
PLUCHEA, Cassini in Bull. de la Soc. philom. 31 (1817). (Spiropodium, Eyrea.)									
P. Indica, Lessing in Schlecht. Linnaea VI, 150 (1831)	—	—	—	—	—	Q.	N.A.	B.fl III,527	M.fr.X,101.
P. tetranthera, F. v. M., Rep. Babb. Exped. 12 (1860)	—	—	—	—	—	Q.	N.A	B.fl.III,528	M.fr.I,34.
P. baccharoides, F. v. M. in Benth. Fl. Austr. III, 528 (1866) ...	—	—	—	—	—	Q.	—	B.fl.III,528	
P. Eyrea, F. v. M., Rep. Babb. Exped. 11, 12 (1860)	W.A.	S.A.	—	—	N.S.W.	Q.	NA.	B.fl.III,528	
P. squarrosa, Bentham, Fl. Austr. III, 529 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,529	
P. dentex, R. Brown in Benth. Fl. Austr. III, 529 (1866) ...	—	—	—	—	—	Q.	—	B.fl.III,529	
PTERIGERON, De Candolle, prodr. V, 293 (1836). (Streptoglossa, Oliganthemum.)									
P. decurrens, De Candolle, prodr. V, 293 (1836)	—	—	—	—	—	—	N.A.	B.fl.III,531	
P. liatroides, Bentham, Fl. Austr. III, 532 (1866)...	W.A.	S.A.	—	—	—	—	—	B.fl.III,532	
P. odorus, Bentham, Fl. Austr. III, 532 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,532	
P. macrocephalus, Bentham, Fl. Austr. III, 532 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,532	
P. microglossus, Bentham, Fl. Austr. III, 532 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,532	
P. dentatifolius, F. v. M., fragm. IX, 119 (1875)..	—	S.A.	—	—	—	—	—	—	M.fr.X,119.
P. adscendens, Bentham, Fl. Austr. III, 533 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,533	
P. filifolius, Bentham, Fl. Austr. III, 533 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,533	
THESPIDIUM, F. v. M. in Journ. of Landsborough's Exped. app. (1862).									
T. basiflorum, F. v. M. in Journ. of Landsb. Exped. App. (1862)	—	—	—	—	—	—	N.A.	B.fl.III,534	M.fr.XI,101.
COLEOCOMA, F. v. M. in Hook. Kew Misc. IX, 19 (1857).									
C. Centaurea, F. v. M. in Hook. Kew Misc. IX, 19 (1857) ...	—	—	—	—	—	—	N.A.	B.fl.III,533	M.fr.X,101.
EPALTES, Cassini in Bull. de la Soc. philom. 139 (1818). (Sphaeromorphaea partly, Ethuliopsis.)									
E. Cunninghamii, Bentham, Fl. Austr. III, 530 (1866)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,530	
E. australis, Lessing in Linnaea V, 148 (1831)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,530	M.fr.XI,101.
E. pleiochaeta, F. v. M., fragm. X, 100 (1877)	—	—	—	—	N.S.W.	—	—	—	M.fr.X,100.
E. Harrisii, F. v. M., fragm. XI, 101 (1881)	—	—	—	—	—	Q.	—	—	M.fr.XI,101.
SPHAERANTHOS, Vaillant in act. Acad. Par. 289 (1719). (Sphaeranthus.)									
S. hirtus, Willdenow, spec. plant. III, 2395 (1800)	—	—	—	—	—	Q.	N.A.	B.fl.III,521	M.fr.III,138.
S. microcephalus, Willdenow, spec. plant. III, 2395 (1800) ...	—	—	—	—	—	—	N.A.	B.fl.III,522	M.fr.III,138.
PTEROCAULON, Elliot, Sketch of the Bot. of S. Carolina and Georgia II, 323 (1824.) (Monenteles.)									
P. verbascifolius, Bentham & J. Hooker, gen. pl. II, 294 (1873)	—	—	—	—	—	Q.	N.A.	B.fl.III,523	
P. Billardieri, F. v. M., Papuan plants, 43 (1876)...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,523	M.fr.XI,101.
P. sphacelatus, Bentham & J. Hooker, gen. plant. II, 94 (1873)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,523	
P. glandulosus, Bentham & J. Hooker, gen. plant. II, 94 (1873)	—	—	—	—	—	Q.	N.A.	B.fl.III,523	
P. sphaeranthoides, Bentham & J. Hooker, gen. plant. II, 94 (1873)	—	—	—	—	—	—	N.A.	B.fl.III,524	
STUARTINA, Sonder in Schlechtendal's Linnaea XXV, 521 (1852).									
S. Muelleri, Sonder in Schlecht. Linnaea XXV, 522 (1852) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,657	M.fr.VIII,145;XI,27.
GNAPHALIUM, Linné, gen. plant. 250 (1737), from J. & C. Bauhin (1619). (Euchiton.)									
G. luteo-album, Linné, spec. plant. 851 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,653	M.fr.V,150.
G. Japonicum, Thunberg, Fl. Japon. 311 (1784)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,653	M.fr.V,150;XI,27.
G. alpigenum, F. v. M., second gen. Report 12 (1854)	—	—	T.	V.	—	—	—	B.fl.III,654	M.fr.V,150.
G. purpureum, Linné, spec. plant. 854 (1753)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,655	
G. Indicum, Linné, spec. plant. 852 (1753)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,655	M.fr.V,149.
G. indutum, J. Hooker in Lond. Journ. of Bot. VI, 121 (1847)...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,655	M.fr.V,150.
G. Traversii, J. Hooker, Handb. N. Zeal. Fl. 154	—	—	—	V.	N.S.W.	—	—	B.fl.III,656	

ANTENNARIA, Gaertner, de fruct. II, 410, t. 167 (1791). (Raoulia partly.)

A. Planchoni, F. v. M.; Gnaphalium, Hook. Fl. Tasm. I, 217, t. 62 (1860)	—	—	—	—	—	—	—	—		
A. uniceps, F. v. M. in Transact. phil. Soc. Vict. t. 105 (1854)..	—	—	—	V.	N.S.W.	—	—	B.fl.III,652	M.fr.VIII,145.	

LEONTOPODIUM, R. Brown in Transact. Linn. Soc. XII, 124 (1817). (Raoulia partly).

L. Catipes, F. v. M. in Papers of the Roy. Soc. of Tasm. 44 (1882)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,651		
L. Meredithae; Antennaria, F. v. M. in Pap. Roy. Soc. of Tasm. 15 (1870)	—	—	T.	—	—	—	—	—		

PTERYGOPAPPUS, J. Hooker in Lond. Journ. of Bot. VII, 120 (1847).

P. Lawrencii, J. Hooker in Lond. Journ. VII, 120 (1847) ...	—	—	T.	—	—	—	—	B.fl.III,656		

PODOTHECA, Cassini in Diction. XXIII, 561 (1822). (Podospermum, Phaenopoda, Lophoclinium.)

P. gnaphalioides, Graham in Bot. Mag. t. 3920 (1842)	W.A.	—	—	—	—	—	—	B.fl.III,601		
P. angustifolia, Lessing, synops. gen. compos. 273 (1832) ..	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,601	M.fr.XI,20.	
P. pygmaea, A. Gray in Hook. Kew Misc. IV, 227 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.III,602		
P. chrysantha, Bentham, Fl. Austr. III, 602 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,602		
P. fuscescens, Bentham, Fl. Austr. III, 602 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,602		

IXIOLAENA, Bentham in Hueg. enum. 66 (1837).

I. brevicompta, F. v. M., fragm. I, 53 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,597	M.fr.I,53.	
I. leptolepis, Bentham, Fl. Austr. III, 597 (1866)...	—	S.A.	—	V.	N.S.W.	—	N.A	B.fl.III,597		
I. supina, F. v. M. in Transact. Vict. Inst. 37 (1855)	—	S.A.	T.	—	—	—	—	B.fl.III,598		
I. tomentosa, Sonder & F. v. M. in Linnaea XXV, 504 (1852) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,598	M.fr.I,53;XI,27.	
I. viscosa, Bentham in Hueg. enum. 66 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,598		

PODOLEPIS, Labillardière, Nov. Holl. pl. specim. II, 57, t. 28 (1806). (Scalia, Panaetia, Scaliopsis, Siemssenia, Stylolepis, Rutidochlamys.)

P. rhytidochlamys, F. v. M., fragm. IV, 79 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,603	M.fr.IV,79.	
P. longipedata, Cunningham in De Cand. prodr. VI, 163 (1837)...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,604	M.fr.I,112.	
P. acuminata, R. Brown in Ait. hort. Kew. ed. sec. V, 82 (1813)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,604	M.fr.XI,27.	
P. canescens, Cunningham in De Cand. prodr. VI, 163 (1837) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,605		
P. aristata, Bentham in Hueg. enum. 64 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,605		
P. pallida, Turczaninow in Bull. de Mosc. XXIV, 78 (1851) ...	W.A.	—	—	—	-	—	N.A.	B.fl.III,605		
P. nutans, Steetz in Lehm. pl. Preiss. I, 464 (1845)	W.A.	—	—	—	—	—	—	B.fl.III,605		
P. gracilis, Graham in Edinb. New phil. journ. V, 379 (1828) ...	W.A.	—	—	—	—	—	—	B.fl.III,606		
P. rugata, Labillardière, Nov. Holl. pl. spec. II, 57, t. 208 (1806)	W.A.	S.A.	—	V.	—	—	—	B.fl.III,606		
P. Lessoni, Bentham, Fl. Austr. III, 606 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,606	M.fr.XI,27.	
P. Siemssenia, F. v. M. in Benth. Fl. Austr. III, 607 (1866) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,607		
P. microcephala, Bentham, Fl. Austr. 607 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,607		

ATHRIXIA, Ker in Bot. Regist. VIII, 681 (1823). (Asteridea, Chrysodiscus, Trichostegia.)

A. australis, Steetz in Lehm. pl. Preiss. I, 482 (1845)	W.A.	—	—	—	—	—	—	B.fl.III,599		
A. gracilis, Bentham, Fl. Austr. III, 599 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,599		
A. multiceps, Bentham, Fl. Austr. III, 599 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,599		
A. stricta, Bentham, Fl. Austr. III, 600 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,600		
A. tenella, Bentham, Fl. Austr. III, 600 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,600		
A. chaetopoda, F. v. M., fragm. X, 56 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,56.	

LEPTORRHYNCHOS, Lessing, syn. composit. 273 (1832). (Rhytidanthe.)

L. squamatus, Lessing, syn. comp. 273 (1832)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,608	M.fr.I,52.	
L. panaetioides, Bentham, Fl. Austr. III, 609 (1866)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,609		
L. tenuifolius, F. v. M., fragm. I, 52 (1858)	—	S.A.	—	V.	—	—	—	B.fl.III,609	M.fr.I,52.	
L. pulchellus, F. v. M. in Linnaea XXV, 500 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,610	M.fr.I,53;XI,27.	
L. Baileyi, F. v. M., fragm. X, 101 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,101.	
L. elongatus, De Candolle, prodr. VI, 160 (1837)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,610	M.fr.XI,27,28.	
L. medius, Cunningham in De Cand. prodr. VI, 160 (1837) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	—	M.fr.XI,85.	
L. Waitzia, Sonder in Schlecht. Linnaea XXV, 501 (1852) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,610		
L. nitidulus, De Candolle, prodr. VI, 160 (1837)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,611		

WAITZIA, Wendland, Collect. pl. II, 13, t. 42 (1808). (Viraya, Morna, Pterochaete.)

W. corymbosa, Wendland, Coll. pl. II, 13, t. 42 (1808)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,635		
W. aurea, Steetz in Lehm. pl. Preiss. I, 452 (1845)	W.A.	—	—	—	—	—	—	B.fl.III,636		
W. nivea, Bentham, Fl. Austr. III, 636 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,636		
W. Steetziana, Lehmann, pl. Preiss. I, 454 (1845)	W.A.	—	—	—	—	—	—	B.fl.III,636		
W. podolepis, Steetz in Lehm. pl. Preiss. I, 450 (1845)	W.A.	—	—	—	—	—	—	B.fl.III,637		
W. paniculata, F. v. M. in Benth. Fl. Austr. III, 637 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,637		

HELIPTERUM, De Candolle, prodr. VI, 211 (1837). (Argyrocome, 1822, Pteropogon, Rodanthe, Xyridanthe, Anisolepis, Hyalosperma, Triptilodiscus, Acroclinium, Monencyanthes, Dimorpholepis, Duttonia, Cassiniola.)

H. Manglesii, F. v. M., fragm. V, 200 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,640	M.fr.X,108.	
H. Margaritae, F. v. M., fragm. XI, 48 (1878)	—	—	—	—	—	—	N.A.	—	M.fr.XI,48.	
H. roseum, Bentham, Fl. Austr. III, 640 (1866)	W.A.	S.A.	—	—	—	—	—	B.fl.III,640	M.fr.X,108.	
H. anthemoides, De Candolle, prodr. VI, 216 (1837)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,641	M.fr.X,108.	
H. polygalifolium, De Candolle, prodr. VI, 216 (1837)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,641	M.fr.X,107;108.	
H. rubellum, Bentham, Fl. Austr. III, 641 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,641	M.fr.X,108.	
H. chlorocephalum, Bentham, Fl. Austr. III, 641 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,641		
H. floribundum, De Candolle, prodr. VI, 217 (1837)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,642	M.fr.X,108.	
H. Pyrethrum, Bentham, Fl. Austr. III, 642 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,642	M.fr.X,110.	
H. heteranthum, Turczan. in Bull. de Mosc. XXIV, 198 (1851)	W.A.	S.A.	—	—	—	—	—	B.fl.III,642	M.fr.III,137;X,108.	
H. Kendallii, F. v. M., fragm. VIII, 168 (1874)	W.A.	—	—	—	—	—	—	—	M.fr.VIII,168;X,107.	
H. stipitatum, F. v. M. in Bentham, Fl. Austr. III, 643 (1866)	—	S.A.	—	—	—	—	—	B.fl.III,643	M.fr.III,133;X,109.	
H. incanum, De Candolle, prodr. VI, 215 (1837)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,643		[110.
H. Cotula, De Candolle, prodr. VI, 215 (1837)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,644	M.fr.II,157;III,134;X,	
H. hyalospermum, F. v. M. in Benth., Fl. Austr. III, 644 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,644	M.fr.X,109.	
H. condensatum, F. v. M., fragm. III, 136 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.III,645	M.fr.III,136.	
H. polyphyllum, F. v. M., fragm. I, 35 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,645	M.fr.I,35.	

H. Humboldtianum, De Candolle, prodr. VI, 216 (1866) ...	W.A.	—	—	—	—	—	N.A.	B.fl.III,645	M.fr.III,135;X,109.
H. Haigii, F. v. M., fragm. X, 107 (1877)	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,107. [20.
H. tenellum, Turczaninow in Bull. de Mosc. XXIV, 198 (1851)	W.A.	S.A.	—	—	—	—	—	B.fl.III,646	M.fr.III,135;X,109;XI,
H. strictum, Bentham, Fl. Austr. III, 646 (1866)	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.III,646	M.fr.X,109.
H. corymbiflorum, Schlechtendal, Linnaea XXI, 448 (1848) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,647	M.fr.X,109.
H. pygmaeum, Bentham, Fl. Austr. III, 647 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,647	M.fr.X,109.
H. spicatum, F. v. M. in Bentham, Fl. Austr. III, 647 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,647	M.fr.X,109.
H. Charsleyae, F. v. M., fragm. VIII, 168 (1874)...	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,168;XI,134.
H. moschatum, Bentham, Fl. Austr. III, 648 (1866)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,648	M.fr.X,107.
H. Tietkensii, F. v. M., fragm. VIII, 227 (1874)...	—	S.A.	—	—	—	—	—	—	M.fr.VIII,227.
H. pterochaetum, Bentham, Fl. Austr. III, 648 (1866)	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,648	M.fr.X,109.
H. polycephalum, Bentham, Fl. Austr. III, 649 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,649	M.fr.III,139;X,109.
H. corymbosum, Bentham, Fl. Austr. III, 649 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,649	M.fr.V,200.
H. laeve, Bentham, Fl. Austr. III, 649 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,649	M.fr.V,200.
H. exiguum, F. v. M. in Transact. Vict. Inst. 39 (1855)... ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,649	M.fr.X,109.
H. Dimorpholepis, Bentham, Fl. Austr. III, 650 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,650	M.fr.X,109;XI,27.

HELICHRYSUM, Vaillant in Act. Acad. Par. 290 (1719), from Theophrastos and Dioscorides. (Elichrysum, Schoenia, Petalolepis, Faustula, Ozothamnus, Swammerdamia, Lawrencella, Argyrophanes, Chrysocephalum, Conanthodium, Xanthochrysum, Argyroglottis, Acanthocladium, Raoulia partly.)

H. Cassinianum, Gaudichaud in Freyc. Voy. Bot. 466, t. 87 (1826)	W.A.	S.A.	—	—	—	—	—	B.fl.III,616	
H. Lawrencella, F. v. M. in Benth. Fl. Austr. III, 616 (1866) ...	W.A.	S.A.	—	—	—	—	—	B.fl.III,616	
H. subulifolium, F. v. M., fragm. III, 134 (1863)..	W.A.	—	—	—	—	—	—	B.fl.III,616	M.fr.III,134.
H. filifolium, F. v. M., fragm. III, 134 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,617	M.fr.III,134.
H. semifertile, F. v. M., Rep. Babb. Exped. 14 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,617	
H. Baxteri, Cunningham in De Cand. prodr. VI, 193 (1837) ...	—	S.A.	V.	—	N.S.W.	—	—	B.fl.III,617	
H. Ayersii, F. v. M., fragm. VIII, 167 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.IX,195.
H. rutidolepis, De Candolle, prodr. VI, 194 (1837)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,618	
H. scorpioides, Labillardière, Nov. Holl. pl. spec. II, 45, t. 191 (1806)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,618	M.fr.VIII,167.
H. oligochaetum, F. v. M., fragm. VI, 235 (1868)...	W.A.	—	—	—	—	—	N.A.	—	M.fr.VI,235.
H. Tepperi, F. v. M. in Wing's South. Science Record, II, 1 (1882)	—	S.A.	—	—	—	—	—	—	
H. obtusifolium, Sonder & F. v. M. in Linnaea XXV, 515 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,619	
H. Calvertianum, F. v. M., fragm. X, 108 (1877)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,136.
H. Spiceri, F. v. M., fragm. XI, 47 (1878)	—	—	T.	—	—	—	—	—	M.fr.XI,47.
H. dealbatum, Labillardière, Nov. Holl. pl. spec. II, 45, t. 190 (1806)	—	—	T.	—	—	—	—	B.fl.III,619	
H. pumilum, J. Hooker, Fl. Tasman. I, 213, t. 60 (1860) ...	—	—	T.	—	—	—	—	B.fl.III,619	
H. lucidum, Henckel, adumbr. pl. hort. Hal. 5 (1806)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,620	M.fr.XI,48.
H. elatum, Cunningham in De Cand. prodr. VI, 193 (1837) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,621	
H. adenophorum, F. v. M. in Transact. Vict. Inst. 38 (1855) ...	—	S.A.	—	—	—	—	—	B.fl.III,622	
H. leucopsidium, De Candolle, prodr. VI, 193 (1837)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,622	M.fr.VI,255.
H. Blandowskianum, Steetz in Schlecht. Linnaea XXV, 512 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,622	
H. oxylepis, F. v. M., fragm. I, 35 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,623	M.fr.I,35.
H. collinum, De Candolle, prodr. VI, 190 (1837)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,623	
H. rupicola, De Candolle, prodr. VI, 190 (1837)	—	—	—	—	—	Q.	—	B.fl.III,623	
H. podolepideum, F. v. M., Rep. Babb. Exped. 13 (1858) ...	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,624	
H. Gilesii, F. v. M., fragm. X, 85 (1876)	W.A.	—	—	—	—	—	N.A.	—	M.fr.X,86.
H. ambiguum, Turczaninow in Bull. de Mosc. 195 (1851) ...	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.III,609	
H. apiculatum, De Candolle, prodr. VI, 195 (1837)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,624	
H. semipapposum, De Candolle, prodr. VI, 195 (1837)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,625	
H. Dockerii, F. v. M. in Benth. Fl. Austr. III, 626 (1866) ...	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,626	M.fr.II,150.
H. Agyroglottis, Bentham, Fl. Austr. III, 626 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,626	
H. ramosum, De Candolle, prodr. VI, 181 (1837)...	W.A.	—	—	—	—	—	—	B.fl.III,626	
H. cordatum, De Candolle, prodr. VI, 180 (1837)...	W.A.	—	—	—	—	—	—	B.fl.III,627	
H. obovatum, De Candolle, prodr. VI, 180 (1837)...	—	—	—	—	N.S.W.	—	—	B.fl.III,627	M.fr.II,89.
H. Bidwillii, Bentham, Fl. Austr. III, 627 (1866)...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,627	
H. Becklerii, F. v. M. in Benth. Fl. Austr. III, 627 (1866) ...	—	—	—	—	N.S.W.	—	—	B.fl.III,627	M.fr.I,183.
H. diotophyllum, F. v. M., fragm. V, 150 (1866)...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,628	M.fr.VIII,46.
H. Thomsoni, F. v. M., fragm. VIII, 45 (1873)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,45.
H. diosmifolium, Don in Steud. Nom. Bot. ed. sec. 738 (1841) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,628	
H. retusum, F. v. M., fragm. VIII, 46 (1873)	—	S.A.	—	V.	N.S.W.	—	—	—	M.fr.VIII,46.
H. decurrens, F. v. M., fragm. VIII, 46 (1873)	—	S.A.	—	V.	N.S.W.	—	—	—	M.fr.VIII,46.
H. Cunninghamii, Bentham, Fl. Austr. III, 629 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,629	
H. reticulatum, Lessing in Steud. Nom. Bot. ed. sec. 740 (1841)	—	—	T.	—	—	—	—	B.fl.III,629	
H. cinereum, F. v. M. in Benth. Fl. Austr. III, 629 (1866) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,629	M.fr.V,199.
H. bracteolatum, Bentham, Fl. Austr. III, 630 (1866)	—	—	T.	—	—	—	—	B.fl.III,630	
H. Kempei, F. v. M. in Melb. Chemist, Jan. (1882)	—	S.A.	—	—	—	—	—	—	
H. cassinioides, Bentham, Fl. Austr. III, 630 (1866)	—	—	—	—	—	Q.	—	B.fl.III,630	
H. Gunnii, F. v. M. in Benth. Fl. Austr. III, 630 (1866) ...	—	—	T.	—	—	—	—	B.fl.III,630	M.fr.V,199.
H. rosmarinifolium, Lessing, synops. gen. comp. 274 (1832) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,631	
H. ledifolium, F. v. M. in Benth. Fl. Austr. III, 630 (1866) ...	—	—	T.	—	—	—	—	B.fl.III,630	M.fr.V,199.
H. ferrugineum, Lessing, synops. gen. comp. 274 (1832)... ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,631	
H. antennarium, F. v. M. in Benth. Fl. Austr. III, 632 (1866)...	—	—	T.	—	—	—	—	B.fl.III,632	M.fr.V,199.
H. obcordatum, F. v. M. in Benth. Fl. Austr. III, 632 (1866) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,632	M.fr.V,199.
H. Backhousii, F. v. M. in Benth. Fl. Austr. III, 632 (1866) ...	—	—	T.	—	—	—	—	B.fl.III,632	M.fr.V,199.
H. cuneifolium, F. v. M. in Benth. Fl. Austr. III, 633 (1866) ...	—	—	—	V.	—	—	—	B.fl.III,633	
H. baccharoides, F. v. M. in Benth. Fl. Austr. III, 633 (1866)...	—	—	T.	V.	—	—	—	B.fl.III,633	M.fr.V,199.
H. lepidophyllum, F. v. M. in Benth. Fl. Austr. III, 633 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,633	M.fr.V,199.
H. scutellifolium, Bentham, Fl. Austr. III, 633 (1866)	—	—	T.	—	—	—	—	B.fl.III,633	M.fr.V,199.
H. pholidotum, F. v. M. in Benth. Fl. Austr. III, 634 (1866) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,634	M.fr.II,131;V,199.
H. lycopodioides, Bentham, Fl. Austr. III, 634 (1866)	—	—	T.	—	—	—	—	B.fl.III,634	M.fr.V,199.
H. selaginoides, F. v. M. in Benth. Fl. Austr. III, 634 (1866) ...	—	—	T.	—	—	—	—	B.fl.III,634	M.fr.V,199.

CASSINIA, R. Brown in Transact. Linn. Soc. XII, 126 (1817). (Non zoologorum, Apolochlamys, Achromolaena, Chromochiton.)

C. leptocephala, F. v. M., fragm. III, 138 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.III,585	M.fr.III,138.
C. compacta, F. v. M., fragm. I, 18 (1858)...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,585	M.fr.I,18.
C. denticulata, R. Brown in Transact. Linn. Soc. XII, 127 (1817)	—	—	—	—	N.S.W.	—	—	B.fl.III,586	
C. longifolia, R. Brown in Transact. Linn. Soc. XII, 127 (1817)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,586	
C. aurea, R. Brown in Transact. Linn. Soc. XII, 127 (1817) ...	—	—	—	—	N.S.W.	—	—	B.fl.III,586	
C. aculeata, R. Brown in Transact. Linn. Soc. XII, 127 (1817)...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,586	
C. laevis, R. Brown in Transact. Linn. Soc. XII, 128 (1817) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,587	
C. tenuifolia, Bentham, Fl. Austr. III, 585 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,585	
C. quinquefaria, R. Brown in Transact. Linn. Soc. XII, 128 (1817)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,587	
C. arcuata, R. Brown in Transact. Linn. Soc. XII, 128 (1817) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,587	
C. subtropica, F. v. M., fragm. I, 17 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,588	M.fr.I,17.
C. Theodori, F. v. M., fragm. V, 148 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,588	M.fr.V,148.
C. spectabilis, R. Brown in Transact. Linn. Soc. XII, 128 (1817)	—	S.A.	T.	V.	—	—	—	B.fl.III,588	

HUMEA, Smith, Exot. bot. I, t. 1 (1804). (Haeckeria, Calomeria.)

H. elegans, Smith, Exot. bot. I, t. 1 (1804)	—	—	—	V.	N.S.W.	—	—	B.fl.III,589	M.fr.I,17.
H. cassiniacea, F. v. M., fragm. I, 17 (1858)	—	S.A.	—	—	—	—	—	B.fl.III,589	M.fr.I,17.
H. punctulata, F. v. M., fragm. III, 137 (1863)	—	S.A.	—	—	—	—	—	B.fl.III,589	M.fr.III,137.
H. ozothamnoides, F. v. M., fragm. I, 17 (1858)	—	—	—	V.	N.S.W.	—	—	B.fl.III,590	M.fr.VIII,142.
H. squamata, F. v. M., fragm. XI, 86 (1880)	—	S.A.	—	V.	—	—	—	—	M.fr.XI,86.

ACOMIS, F. v. M., fragm. II, 89 (1860).

A. Rutidosis, F. v. M., fragm. II, 89 (1860)	—	—	—	—	N.S.W.	—	—	B.fl.III,591	M.fr.II,89.
A. macra, F. v. M., fragm. IV, 145 (1864)...	—	—	—	—	—	Q.	—	B.fl.III,591	M.fr.IV,145.

RUTIDOSIS, De Candolle, prodr. VI, 158 (1837). (Pumilo, Actinopappus, Lepidocoma.)

R. leiolepis, F. v. M. in Transact. Vict. Inst. 131 (1855) ...	—	—	—	V.	—	—	—	B.fl.III,593	
R. leptorrhynchoides, F. v. M., fragm. V, 148 (1866)	—	—	—	V.	N.S.W.	—	—	B.fl.III,593	M.fr.V,148.
R. Brownii, Bentham, Fl. Austr. III, 594 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,594	
R. helichrysoides, De Candolle, prodr. VI, 159 (1837)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,594	
R. leucantha, F. v. M., fragm. I, 35 (1858)	—	—	—	—	—	Q.	—	B.fl.III,594	M.fr.I,35.
R. Murchisonii, F. v. M., fragm. I, 34 (1858)	—	—	—	—	—	Q.	—	B.fl.III,594	M.fr.I,34.
R. Pumilo, Bentham, Fl. Austr. III, 595 (1866)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,595	

PITHOCARPA, Lindley, Bot. Regist. XXV, App. XXIII (1839).

P. corymbosa, Lindley, Bot. Regist. XXV, App. XXIII (1839)	W.A.	—	—	—	—	—	—	B.fl.III,590	

AMMOBIUM, R. Brown in Bot. Magaz. t. 2459 (1824).

A. alatum, R. Brown in Bot. Magaz. t. 2459 (1824)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,583	
A. craspedioides, Bentham, Fl. Austr. III, 584 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,584	

IXODIA, R. Brown in Ait. hort. Kew. sec. ed. IV, 17 (1812).

I. achilleoides, R. Brown in Ait. hort. Kew. sec. ed. IV, 17 (1812)	—	S.A.	—	V.	—	—	—	B.fl.III,583	

MILLOTIA, Cassini in Ann. sc. nat. XVII, 416 (1829).

M. tenuifolia, Cassini in Ann. sc. nat. XVII, 416 (1829)... ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,596	
M. Greevesii, F. v. M., fragm. III, 18, t. 19 (1863)	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,596	M.fr.III,18.
M. Kempei, F. v. M. in Wing's South. Sc. Record II, 1 (1882)...	—	S.A.	—	—	—	—	—	—	

TOXANTHUS, Turczaninow in Bull. de Mosc. XXIV, 177 (1851). (Anthocerastes, Scyphocoronis.)

T. perpusillus, Turczaninow in Bull. Mosc. XXIV, 177 (1851) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,592	M.fr.XI,49.
T. Muelleri, Bentham, Fl. Austr. III, 592 (1866)	W.A.	S.A.	—	—	—	—	—	B.fl.III,592	M.fr.XI,49.
T. major, Turczaninow in Bull. de Mosc. XXIV, 64 (1851) ...	W.A.	—	—	—	—	—	—	B.fl.III,592	M.fr.XI,49.

QUINETIA, Cassini in Diction. sc. nat. LX, 579 (1830).

Q. Urvillei, Cassini in Diction. LX, 579 (1830)	W.A.	S.A.	—	—	—	—	—	B.fl.III,595	

PHACELOTHRIX, F. v. M., fragm. XI, 49 (1878).

P. cladochaeta, F. v. M., fragm. XI, 49 (1878)	—	—	—	—	—	Q.	—	—	M.fr.V,199;XI,49.

ERIOCHLAMYS, Sonder and F. v. M. in Schlecht. Linnaea XXV, 488 (1852).

E. Behrii, Sonder & F. v. M. in Linnaea XXV, 488 (1852) ...	--	S.A.	—	V.	N.S.W.	—	—	B.fl.III,591	

MYRIOCEPHALUS, Bentham in Hueg. enum. 61 (1837). (Hyalolepis, Antheidosorus, Gilberta, Lamprochlaena, Elachopappus, Polycalymma; some readily to be restituted.)

M. rhizocephalus, Bentham, Fl. Austr. III, 557 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,557	M.fr.III,155.
M. nudus, A. Gray in Hook. Kew Misc. III, 174 (1851)	W.A.	—	—	—	—	—	—	B.fl.III,558	M.fr.III,157.
M. appendiculatus, Bentham in Hueg. enum. 61 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,558	[145.
M. Rudallii, F. v. M., fragm. III, 157 (1863)	W.A.	S.A.	—	—	—	—	—	B.fl.III,559	M.fr.III,157;VI,235;VII,
M. gracilis, Bentham, Fl. Austr. III, 559 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,559	M.fr.VIII,169.
M. helichrysoides, A. Gray in Hook. Kew Misc. III, 175 (1851)	W.A.	—	—	—	—	—	—	B.fl.III,559	
M. suffruticosus, Bentham, Fl. Austr. III, 559 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,559	
M. Stuartii, Bentham, Fl. Austr. III, 560 (1866)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,560	
M. Guerinae, F. v. M., fragm. VIII, 169 (1874)	W.A.	—	—	—	—	—	—	—	M.fr.VIII,169.

ANGIANTHUS, Wendland, Collect. pl. II, 31 (1809). (Siloxerus, Styloncerus, Ogcerostylus, Cylindrosorus, Phyllocalymma, Skirrophorus, Chrysocoryne, Eriocladium, Pogonolepis, Piptostemma, Epitriche, Gamozygis, Cephalosorus partly, Hyalochlamys, Dithyrostegia, Pleuropappus; some readily to be restituted.)

A. humifusus, Bentham, Fl. Austr. III, 563 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,563	
A. tomentosus, Wendland, Collect. pl. II, 31, t. 48 (1809) ...	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.III,562	
A. pleuropappus, Bentham, Fl. Austr. III, 563 (1866)	—	S.A.	—	—	—	—	—	B.fl.III,563	
A. brachypappus, F. v. M. in Trans. phil. Soc. Vict. I, 44 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,563	
A. myosuroides, Bentham, Fl. Austr. III, 563 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,563	
A. tenellus, Bentham, Fl. Austr. III, 564 (1866)	W.A.	S.A.	—	—	—	—	—	B.fl.III,564	
A. pusillus, Bentham, Fl. Austr. III, 564 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,564	
A. Milnei, Bentham, Fl. Austr. III, 564 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,564	
A. Cunninghamii, Bentham, Fl. Austr. III, 565 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,565	
A. phyllocephalus, Bentham, Fl. Austr. III, 565 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,565	M.fr.III,159.

A. micropoides, Bentham, Fl. Austr. III, 565 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,565							
A. microcephalus, Bentham, Fl. Austr. III, 566 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,566	M.fr.III,158.						
A. platycephalus, Bentham, Fl. Austr. III, 566 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,566							
A. Drummondii, Bentham, Fl. Austr. III, 566 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,566							
A. Preissianus, Bentham, Fl. Austr. III, 566 (1866)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,566							
A. pygmaeus, Bentham, Fl. Austr. III, 567 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,567							
A. globifer, Bentham, Fl. Austr. III, 567 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,567							
A. demissus, Bentham, Fl. Austr. III, 567 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,567							
A. strictus, Bentham, Fl. Austr. III, 568 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,568							
A. plumiger, Bentham, Fl. Austr. III, 568 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,568							
A. amplexicaulis, Bentham, Fl. Austr. III, 568 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,568							
GNEPHOSIS, Cassini in Bull. de la Soc. philom. 45 (1820). (Cephalosorus partly, Nematopus, Crossolepis, Leptotriche, Trichanthodium, Cyathopappus; some readily to be restituted.)															
G. Burkittii, Bentham, Fl. Austr. III, 570 (1866)	—	S.A.	—	—	—	—	—	B.fl.III,570							
G. eriocarpa, Bentham, Fl. Austr. III, 570 (1866)	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,570							
G. macrocephala, Turcz. in Bull. Soc. Mosc. XXIV, 190 (1851)...	W.A.	—	—	—	—	—	—	B.fl.III,570							
G. skirrophora, Bentham, Fl. Austr. III, 570 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,570							
G. cyathopappa, Bentham, Fl. Austr. III, 571 (1866)	—	S.A.	—	—	N.S.W.	—	N.A.	B.fl.III,571							
G. codonopappa, F. v. M. in Giles, geogr. trav. 217 (1875) ...	—	S.A.	—	—	—	—	—	—	M.fr.IX,2.						
G. leptoclada, Bentham, Fl. Austr. III, 571 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,571	M.fr.III,158.						
G. arachnoidea, Turcz. in Bull. Soc. Mosc. XXIV, 189 (1851) ...	W.A.	S.A.	—	—	—	—	—	B.fl.III,571							
G. tenuissima, Cassini in Bull. de la Soc. philom. 43 (1820) ..	W.A.	—	—	—	—	—	—	B.fl.III,572							
G. acicularis, Bentham, Fl. Austr. III, 572 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,572							
G. pygmaea, Bentham, Fl. Austr. III, 572 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,572							
G. brevifolia, Bentham, Fl. Austr. III, 572 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,572							
G. eriocephala, Bentham, Fl. Austr. III, 573 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,573							
DECAZESIA, F. v. M., fragm. XI, 71 (1879).															
D. hecatocephala, F. v. M., fragm. XI, 72 (1879)...	—	—	—	—	—	—	N.A.	—	M.fr.XI,72.						
CALOCEPHALUS, R. Brown in Transact. Linn. Soc. XII, 106 (1817). (Leucophyta, Pachysura, Blennospora, Achrysum.)															
C. Drummondii, Bentham, Fl. Austr. III, 574 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,574							
C. Brownii, F. v. M., Rep. Rabb. Exped. 13 (1858)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,574							
C. Sonderi, F. v. M., Rep. Babb. Exped. 13 (1858)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,775							
C. lacteus, Lessing, syn. gen. comp. 271 (1832)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,575							
C. citreus, Lessing, syn. gen. comp. 271 (1832)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,575							
C. angianthoides, Bentham, Fl. Austr. III, 575 (1866)	W.A.	—	T.	—	—	—	—	B.fl.III,575							
C. Francisii, Bentham, Fl. Austr. III, 576 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,576							
C. platycephalus, Bentham, Fl. Austr. 115, 576 (1866)	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,576							
C. multiflorus, Bentham, Fl. Austr. III, 576 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,576							
C. aeruoides, Bentham, Fl. Austr. III, 576 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,576							
CEPHALIPTERUM, A. Gray in Hook. Kew Misc. IV, 271 (1852).															
C. Drummondii, A. Gray in Hook. Kew Misc. IV, 272 (1852) ...	W.A.	S.A.	—	—	—	—	—	B.fl.III,577							
GNAPHALODES, A. Gray in Hook. Kew Misc. IV, 228 (1852).															
G. uliginosum, A. Gray in Hook. Kew Misc. IV, 228 (1852) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,578	M.fr.XI,27.						
G. condensatum, A. Gray in Hook. Kew Misc. IV, 228 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.III,578							
G. filifolium, Bentham, Fl. Austr. III, 578 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,578							
CRASPEDIA, G. Forster, florul. insul. Austr. prodr. 306 (1786). (Richea, Pycnosorus.)															
C. Richea, Cassini in Diction. XI, 353 (1818)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,579							
C. pleiocephala, F. v. M. in Schlecht. Linnaea XXV, 404 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,580	M.fr.XI,27.						
C. chrysantha, Bentham, Fl. Austr. III, 580 (1866)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,580							
C. globosa, Bentham, Fl. Austr. III, 580 (1866)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,580							
CHTHONOCEPHALUS, Steetz in Lehm. pl. Preiss. I, 444 (1845). (Chamaesphaerion, Gyrostephium, Lachnothalamus.)															
C. tomentellus, Bentham, Fl. Austr. III, 581 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,581							
C. Pseudevax, Steetz in Lehm. pl. Preiss. I, 445 (1845)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,582							
C. pygmaeus, Bentham, Fl. Austr. III, 582 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,582							
NABLONIUM, Cassini in Diction. XXXIV, 101 (1825).															
N. calyceroides, Cassini, Dict. XXXIV, 101 (1825)	—	—	T.	—	—	—	—	B.fl.III,545							
CHRYSOGONUM, Linné, hort. Cliffort. 424 (1737). (Moonia, Pentalepis.)															
C. trichodesmoides, Bentham & J. Hooker, gen. pl. II, 350 (1873)	—	—	—	—	—	—	N.A.	B.fl.III,540	M.fr.VIII,145.						
C. ecliptoides, Bentham & J. Hooker, gen. pl. II, 350 (1873) ...	—	—	—	—	—	—	N.A.	B.fl.III,540	M.fr.VIII,145.						
C. procumbens, Bentham & J. Hooker, gen. pl. II, 350 (1873) ...	—	—	—	—	—	—	N.A.	B.fl.III,540	M.fr.VIII,145.						
SIEGESBECKIA, Linné, hort. Cliffort. 412 (1737).															
S. orientalis, Linné, spec. plant. 900 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,535							
ENHYDRA, Loureiro, Flor. Cochinch. II, 510 (1790). (Enydra.)															
E. paludosa, De Candolle, prodr. V, 637 (1836)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,546	M.fr.III,139.						
ECLIPTA, Linné, mantissa altera, 157 (1771).															
E. alba, Hasskarl, plant. Javan. rar. 528 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,536							
E. latifolia, Linné, fil. suppl. 378 (1781)	—	—	—	—	—	Q.	N.A.	—	M.fr.VIII,145.						
E. platyglossa, F. v. M., fragm. II, 135 (1861)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,536	M.fr.XI,27.						
WEDELIA, Jacquin, stirp. Amer. hist. 217, t. 130 (1763). (Wollastonia.)															
W. calendulacea, Lessing, synops. gen. comp. 222 (1832) ...	—	—	—	—	—	Q.	—	B.fl.III,537							
W. urticifolia, De Candolle in Wight, Contrib. Bot. Ind. 18 (1834)	—	—	—	—	—	—	N.A.	B.fl.III,538							
W. spilanthoides, F. v. M., fragm. V, 64 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,538	M.fr.V,64.						
W. verbesinoides, F. v. M. in Benth. Fl. Austr. III, 538 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,538							
W. biflora, De Candolle in Wight, Contrib. Bot. Ind. 18 (1834)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,539							
W. asperrima, Bentham, Fl. Austr. III, 539 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,539							

SPILANTHES, Jacquin, stirp. Amer. hist. 214, t. 126 (1763).

S. grandiflora, Turczaninow in Bull. de Mosc. XXIV, 185 (1851)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,541	M.fr.V,63.
S. anactina, F. v. M., fragm. V, 63 (1865)	—	—	—	—	—	—	N.A.	B.fl.III,541	M.fr.V,63.

BIDENS, Dillenius, hort. Eltham. 51 and 52, t. 43 et 44 (1732), from Cesalpini (1583). (Possibly immigrated.)

B. tripartitus, Linné, spec. plant. 831 (1753)	—	—	—	V.	N.S.W.	—	—	B.fl.III,543	
B. pilosus, Linné, spec. plant. 832 (1753)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,543	
B. bipinnatus, Linné, spec. plant. 832 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,543	

GLOSSOGYNE, Cassini in Diction. LI, 475 (1827). (Diodontium.)

G. tenuifolia, Cassini in Dict. LI, 475 (1827)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,544	M.fr.I,51.
G. retroflexa, F. v. M., fragm. I, 51 (1858)	—	—	—	—	—	Q.	—	B.fl.III,544	
G. filifolia, F. v. M. in Benth. Fl. Austr. III, 544 (1866) ...	—	—	—	—	—	—	N.A.	B.fl.III,544	

FLAVERIA, A. L. de Jussieu, gen. plant. 186 (1789).

F. Australasica, Hooker in Mitch. Trop. Austr. 118 (1848) ...	—	—	—	—	—	Q.	N.A.	B.fl.III,546	M.fr.I,183.

SOLIVA, Ruiz et Pavon, prodr. 113, t. 24 (1794). (Gymnostyles; possibly immigrated.)

S. anthemifolia, R. Brown in Transact. Linn. Soc. XII, 102 (1817)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,552	

COTULA, Linné, syst. nat. 9 (1735); Linné, gen. pl. 256 (1737). (Gymnogyne, Strongylospermum, Pleiogyne, Symphyomera, Stenosperma, Leptinella.)

C. filifolia, Thunberg, prodr. pl. Capens. 161 (1800)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,548	M.fr.VI,114.
C. coronopifolia, Linné, sp. pl. 892 (1753)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,549	M.fr.VI,114.
C. Gymogyne, F. v. M. in Benth. Fl. Austr. III, 549 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,549	
C. australis, J. Hooker, Fl. N. Zeal. I, 128 (1853)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,550	
C. alpina, J. Hooker, Fl. Tasman. I, 192, t. 51 (1860)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,550	
C. Drummondii, Bentham, Fl. Austr. III, 550 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,550	
C. reptans, Bentham, Fl. Austr. III, 551 (1866)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,551	
C. Filicula, J. Hooker in Benth. Fl. Austr. III, 551 (1866) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,551	

CENTIPEDA, Loureiro, Fl. Cochinch. II, 492 (1790). (Myriogyne, Sphaeromorphaea partly.)

C. racemosa, F. v. M.; Myriogyne, Hook. in Mitch. Tr. Austr. 353 (1848)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,553	
C. orbicularis, Loureiro, Fl. Cochinch. II, 492 (1790)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,553	M.fr.VIII,142.
C. Cunninghami, F. v. M., fragm. VIII, 143 (1874)	W.A.	S.A.	—	V.	N.S.W.	—	—	—	
C. thespidioides, F. v. M., fragm. VIII, 143 (1874)	—	S.A.	—	V.	N.S.W.	—	—	—	

ABROTANELLA, Cassini in Dict. XXXVI, 27 (1825). (Scleroleima, Trineuron.)

A. fosterioides, J. Hooker, Hand. N. Zeal. Fl. 139 (1864) ...	—	—	T.	—	—	—	—	B.fl.III,554	
A. nivigena, F. v. M. in Benth. Fl. Austr. III, 554 (1866) ...	—	—	—	V.	—	—	—	B.fl.III,554	
A. scapigera, F. v. M. in Benth. Fl. Austr. III, 554 (1866) ...	—	—	T.	—	—	—	—	B.fl.III,554	

ELACHANTHUS, F. v. M. in Schlecht. Linnaea XXV, 410 (1852).

E. pusillus, F. v. M. in Schlecht. Linnaea XXV, 411 (1852) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,555	

CERATOGYNE, Turczaninow in Bull. de Mosc. XXIV, 68 (1851). (Diotosperma.)

C. obionoides, Turczaninow in Bull. de Mosc. XXIV, 68 (1851)	W.A.	—	—	—	—	—	—	B.fl.III,555	

ISOETOPSIS, Turczaninow in Bull. de Mosc. XXIV, 175 (1851).

I. graminifolia, Turcz. in Bull. de Mosc. XXIV, 175, t. 3 (1851)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,556	

EMILIA, Cassini in Bulletin de la Soc. philomat. 68 (1817).

E. purpurea, Cassini in Diction. des sc. nat. XXXIV, 393 (1825)	—	—	—	—	—	Q.	—	—	M.fr.XII.

GYNURA, Cassini in Diction. des sc. nat. XXXIV, 391 (1825).

G. Pseudo-China, De Candolle, prodr. VI, 299 (1837)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,661	

SENECIO, Tournefort, inst. 456, t. 260 (1700), from Plinius. (Bedfordia, Centropappus.)

S. Gregorii, F. v. M., pl. of Greg. Search-Exped. for Leichh. 7 (1859)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,661	
S. platylepis, De Candolle, prodr. VI, 371 (1837)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,662	M.fr.XI,27.
S. papillosus, F. v. M. in Transact. phil. Inst. Vict. II, 69 (1857)	—	—	T.	—	—	—	—	B.fl.III,662	
S. primulifolius, F. v. M. in Transact. phil. Inst. Vict. II, 69 (1857)	—	—	T.	—	—	—	—	B.fl.III,662	
S. pectinatus, De Candolle, prodr. VI, 372 (1837)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,662	
S. spathulatus, A. Richard, sert. Astrol. 125 (1833)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,663	
S. megaglossus, F. v. M. in Linnaea XXV, 419 (1852)	—	S.A.	—	—	—	—	—	B.fl.III,663	M.fr.VI,255.
S. magnificus, F. v. M. in Linnaea XXV, 418 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,663	
S. insularis, Bentham, Fl. Austr. III, 666 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,664	
S. Centropappus, F. v. M., annual report 26 (1858)	—	—	T.	—	—	—	—	B.fl.III,664	
S. macranthus, A. Richard, sert. Astrol. 126 (1833)	—	—	—	—	N.S.W.	—	—	B.fl.III,664	
S. Daltoni, F. v. M., fragm. VI, 27 (1867)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VI,27.
S. lautus, Solander in G. Forst. prodr. 91 (1786)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,665	M.fr.XI,20.
S. capillifolius, J. Hooker in Lond. Journ VI, 123 (1847) ...	—	—	T.	—	—	—	—	B.fl.III,667	
S. vagus, F. v. M. in Transact. phil. Soc. Vict. I, 46 (1854) ...	—	—	—	V.	N.S.W.	—	—	B.fl.III,667	
S. amygdalifolius, F. v. M., fragm. I, 232 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,668	M.fr.I,232.
S. velleioides, Cunningham in De Cand. prodr. VI, 374 (1837) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,668	
S. australis, A. Richard, sert. Astrol. 131, t. 39 (1833)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,668	
S. Behrianus, Sonder and F. v. M. in Linnaea XXV, 527 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,669	
S. leucoglossus, F. v. M., fragm. II, 15 (1861)	W.A.	—	—	—	—	—	—	B.fl.III,669	M.fr.II,15.
S. brachyglossus, F. v. M. in Schlecht. Linnaea XXV, 525 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,669	M.fr.XI,27.
S. Georgianus, De Candolle, prodr. VI, 371 (1837)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,670	
S. Gilberti, Turczaninow in Bull. Soc. Mosc. XXIV, 208 (1851)	W.A.	—	—	—	—	—	—	B.fl.III,670	
S. ramosissimus, De Candolle, prodr. VI, 371 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,671	
S. odoratus, Horneman, hort. reg. bot. Hafn. II, 809 (1815) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,671	
S. Cunninghamii, De Candolle, prodr. VI, 371 (1837)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,671	
S. anethifolius, Cunningham in De Cand. prodr. VI, 371 (1837) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,672	
S. Bedfordii, F. v. M. annual report 26 (1858)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,673	
S. Billardieri, F. v. M. annual report 26 (1858)	—	—	T.	—	—	—	—	B.fl.III,673	

ERECHTITES, Rafinesque, Fl. Ludov. 65 (1817).

E. prenanthoides, De Candolle, prodr. VI, 296 (1837)	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,658	
E. Atkinsoniae, F. v. M., fragm. V, 88 (1866)	...	...	. .	—	—	—	—	N.S.W.	Q.	—	B.fl.III,658	M.fr.V,88.
E. arguta, De Candolle, prodr. VI, 296 (1837)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,659	
E. mixta, De Candolle, prodr. VI, 297 (1837)	..	...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,659	
E. quadridentata, De Candolle, prodr. VI, 295 (1837)	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,660	
E. hispidula, De Candolle, prodr. VI, 296 (1837)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,660	

CYMBONOTUS, Cassini in Diction. XXXV, 397 (1825).

C. Lawsonianus, Gaudichaud in Freyc. Voy. Bot. 462, t. 86 (1826)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,674	M.fr.XI,27.

SAUSSUREA, De Candolle in Ann. du Mus. XVI, 156 (1810). (Aplotaxis, Haplotaxis.)

S. carthamoides, Bentham, Fl. Hongk. 168 (1861)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,456	

CENTAUREA, Linné, hort. Cliffort. 420 (1737). (Leuzea partly.)

C. Australis, Bentham & J. Hooker, gen. plant. II, 479 (1873)	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,457	

TRICHOCLINE, Cassini in Bull. de la Soc. philom. 13 (1817). (Amblysperma.)

T. scapigera, Bentham & J. Hooker, gen. plant. II, 497 (1873)	...	W.A.	—	—	—	—	—	—	—	B.fl.III,676	

MICROSERIS, D. Don in Edinb. phil. Mag. XI, 388 (1832). (Phyllopappus, Scozonera partly.)

M. Forsteri, J. Hooker, Fl. N. Zeal. I, 151 (1853)	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,676	M.fr.XI,27.

CREPIS, Linné, gen. pl. 240 (1737). (Youngia.)

C. Japonica, Bentham, Fl. Hongk. 194 (1861)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,679	

CAMPANULACEAE.

A. L. de Jussieu, gen. plant. 163 (1789), from B. de Jussieu (1759).

LOBELIA, Linné, Fl. Lappon. 227 (1737). (Rapuntium, Pratia, Grammatotheca, Holostigma.)

L. heterophylla, Labillard., Nov. Holl. pl. spec. I, 52, t. 74 (1804)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,124	M.fr.IV,184;X,41.
L. Browniana, Roemer & Schultes, syst. veg. V, 71 (1819)	...	—	S.A.	T.	V.	N.S.W.	Q.	—	—	
L. simplicicaulis, R. Brown, prodr. 564 (1810)	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,124	M.fr.IV,184;X,41.
L. microsperma, F. v. M., fragm. X, 41 (1876)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	—	M.fr.X,41.
L. dentata, Cavanilles, icon. pl. VI, 14, t. 522 (1801)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,125	M.fr.IV,183;X,42.
L. gracilis, Andrews, Bot. Reposit. t. 340 (1805)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,125	M.fr.IV,183;X,42.
L. rhytidosperma, Bentham, Fl. Austr. IV, 126 (1869)	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,126	M.fr.X,40.
L. tenuior, R. Brown, prodr. 564 (1810)	..	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,126	M.fr.IV,183;X,42.
L. rhombifolia, De Vriese in Lehm. pl. Preiss. I, 397 (1845)	...	W.A.	S.A.	—	V.	—	—	—	B.fl.IV,126	M.fr.X,42.
L. parvifolia, R. Brown, prodr. 564 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,127	M.fr.IV,183.
L. trigonocaulis, F. v. M., fragm. I, 18 (1858)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,127	M.fr.I,18,242.
L. Bergiana, Chamisso in Schlecht. Linnaea VIII, 217 (1835)	...	W.A.	—	—	—	—	—	—	B.fl.IV,127	M.fr.II,20;IV,171;X,43.
L. anceps, Thunberg, prodr. plant. cap. 40 (1794)	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,128	M.fr.IV,184;VI,114; X,43
L. surrepens, J. Hooker, Fl. Tasman. I, 237, t. 69 (1860)	...	—	—	T.	—	—	—	—	B.fl.IV,129	M.fr.IV,183.
L. membranacea, R. Brown, prodr. 563 (1810)	...	...	...	—	—	—	—	—	Q.	—	B.fl.IV,129	M.fr.IV,171;X,42.
L. stenophylla, Bentham, Fl. Austr. III, 130 (1869)	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,130	M.fr.X,42.
L. humistrata, F. v. M. in Benth. Fl. Austr. III, 130 (1869)	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,130	M.fr.IV,171;X,42.
L. dioica, R. Brown, prodr. 565 (1810)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.IV,130	M.fr.IV,183;X,43.
L. purpurascens, R. Brown, prodr. 563 (1810)	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,131	M.fr.IV,183;X,42.
L. pratioides, Bentham, Fl. Austr. IV, 131 (1869)	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,131	M.fr.X,43.
L. irrigua, R. Brown, prodr. 563 (1810)	...	...	...	...	—	—	T.	—	—	—	B.fl.IV,132	
L. gelida, F. v. M., fragm. IV, 183 (1864)	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,132	M.fr.IV,183.
L. platycalyx, F. v. M., fragm. IV, 183 (1864)	...	...	...	—	—	T.	V.	—	—	—	B.fl.IV,133	M.fr.X,43.
L. concolor, R. Brown, prodr. 563 (1810)	..	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,133	M.fr.X,43.
L. Benthami, F. v. M.; Pratia puberula, Benth. Fl. Austr. IV, 133 (1869)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,133	
L. pedunculata, R. Brown, prodr. 563 (1810)	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,133	M.fr.X,42.

ISOTOMA, R. Brown, prodr. 565 (1810). (Lobelia partly, Enchysia partly, Laurencia partly.)

I. Brownii, G. Don, gen. syst. III, 716 (1834)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,134	M.fr.X,40.
I. axillaris, Lindley, Bot. Regist. t. 964 (1826)	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,135	M.fr.X,39.
I. petraea, F. v. M. in Linnaea XXV, 420 (1852)	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.IV,135	M.fr.X,39.
I. pusilla, Bentham in Hueg. enum. 75 (1837)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,135	
I. Gulliverii, F. v. M., fragm. X, 39 (1876)	...	...	...	—	—	—	—	—	—	N.A.	—	M.fr.X,39.
I. scapigera, G. Don, gen. syst. III, 716 (1834)	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.IV,136	M.fr.IV,183;X,39.
I. fluviatilis, F. v. M. in Bentham, Fl. Austr. IV, 136 (1869)	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,136	M.fr.X,39.

WAHLENBERGIA, Schrader, cat. hort. bot. Goetting. (1814).

W. gracilis, A. De Candolle, Monogr. comp. 142 (1830)	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,137	

CANDOLLEACEAE.

F. v. M., fragm. VIII, 41 (1873). (Stylideae, R. Brown, prodr. 565, 1810.)

CANDOLLEA, Labillardière in Ann. du Mus. VI, 451 (1805). (Stylidium, Ventenata, Forsteropsis.)

C. (St.) carnosa, Bentham in Hueg. enum. 71 (1837)	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,6	
C. (St.) pilosa, Labillardière, Nov. Holl. pl. spec. II, 63, t. 213 (1806)	W.A.	—	—	—	—	—	—	B.fl.IV,7	
C. (St.) reduplicata, R. Brown, prodr. 568 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,7	
C. (St.) scabrida, Lindley, Bot. Regist. XXV, App. XXVIII (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,8	
C. (St.) hirsuta, R. Brown, prodr. 568 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,8	
C. (St.) crossocephala, F. v. M., fragm. VI, 5 (1866)	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,8	M.fr.VI,5,254;X,58.
C. (St.) juncea, R. Brown, prodr. 569 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,9	M.fr.X,58.
C. (St.) guttata, R. Brown, prodr. 571 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,9	
C. (St.) repens, R. Brown, prodr. 571 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,9	M.fr.X,58.
C. (St.) graminifolia, Swartz, Mag. der nat. Ges. Berl. 49 (1807)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,10	M.fr.VI,6;X,58.
C. (St.) linearis, Swartz, Mag. der nat. Ges. Berl. 50 (1807)	...	—	—	—	—	N.S.W.	—	—	B.fl.IV,11	
C. (St.) elongata, Bentham, Fl. Austr. IV, 11 (1869)	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,11	M.fr.X,86.
C. (St.) spinulosa, R. Brown, prodr. 569 (1815)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.IV,11	

C. (St.) limbata, F. v. M., fragm. X, 57 (1876)	W.A.	—	—	—	—	—	—	—	B.fl.IV,11	M.fr.X,57.
C. (St.) caespitosa, R. Brown, prodr. 569 (1810)	W.A.	—	—	—	—	—	—	—	—	M.fr.X,57.
C. (St.) squamellosa, De Candolle, prodr. VII, 782 (1838) ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,12	
C. (St.) violacea, R. Rrown, prodr. 569 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,12	
C. (St.) lutea, R. Brown, prodr. 570 (1810)...	W.A.	—	—	—	—	—	—	—	B.fl.IV,12	M.fr.X,58,86.
C. (St.) filifera, R. Brown, prodr. 569 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,13	
C. (St.) ciliata, Lindley, Bot. Regist. XXV, App. XXVIII (1839)	W.A.	—	—	—	—	—	—	—	B.fl.IV,13	
C. (St.) sobolifera, F. v. M. in transact. Vict. Inst. 131 (1855) ...	—	—	—	V.	—	—	—	—	B.fl.IV,13	
C. (St.) Floodii, F. v. M., fragm. I, 149 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.IV,13	M.fr.VI,6;X,58.	
C. (St.) disperma, F. v. M., fragm. IV, 93 (1864)...	W.A.	—	—	—	—	—	—	—	B.fl.IV,14	M.fr.IV,93.
C. (St.) calcarata, R. Brown, prodr. 570 (1810)	W.A.	S.A.	—	V.	—	—	—	—	B.fl.IV,14	M.fr.VI,6.
C. (St.) perpusilla, J. Hooker in Lond. Journ. of Bot. VI, 266 (1847)	W.A.	S.A.	T.	V.	—	—	—	—	B.fl.IV,15	M.fr.VI,78;X,58.
C. (St.) eriorrhiza, R. Brown, prodr. 569 (1810)	—	—	—	—	—	Q.	—	—	B.fl.IV,15	M.fr.I,147;VI,6;X,58.
C. (St.) debilis, F. v. M., fragm. I, 149 (1859)	—	—	—	—	N.S.W.	Q.	—	—	B.fl.IV,15	M.fr.VI,6;X,58.
C. (St.) floribunda, R. Brown, prodr. 569 (1810)	—	S.A.	—	—	—	—	N.A.	—	B.fl.IV,15	M.fr.I,148;VI,6;X,58.
C. (St.) leptorrhiza, F. v. M., fragm. I, 148 (1859)	—	—	—	—	—	—	N.A.	—	B.fl.IV,16	M.fr.I,148.
C. (St.) assimilis, R. Brown, prodr. 569 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,16	M.fr.X,57.
C. (St.) rupestris, Sonder in Lehm. pl. Preiss. I, 375 (1845) ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,16	
C. (St.) spathulata, R. Brown, prodr. 569 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,17	
C. (St.) Barleei, F. v. M., fragm. VI, 5, t. 69 (1866)	W.A.	—	—	—	—	—	—	—	B.fl.IV,17	M.fr.VI,5,249.
C. (St.) lineata, Sonder in Lehm. pl. Preiss. I, 376 (1845) ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,17	
C. (St.) glauca, Labillard., Nov. Holl. pl. spec. II, 64, t. 214 (1806)	W.A.	—	—	—	—	—	—	—	B.fl.IV,17	
C. (St.) amoena, R. Brown, prodr. 570 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,18	
C. (St.) striata, Lindley, Bot. Regist. XXV, App. XXVIII (1839)	W.A.	—	—	—	—	—	—	—	B.fl.IV,18	
C. (St.) diversifolia, R. Brown, prodr. 570 (1810)...	W.A.	—	—	—	—	—	—	—	B.fl.IV,18	M.fr.X,58.
C. (St.) articulata, R. Brown, prodr. 570 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,19	
C. (St.) Brunoniana, Bentham in Hueg. enum. 72 (1837)... ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,19	
C. (St.) diuroides, Lindley, Bot. Regist. XXV, App. XXIX (1839)	W.A.	—	—	—	—	—	—	—	B.fl.IV,20	
C. (St.) scandens, R. Brown, prodr. 570 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,20	M.fr.VI,78.
C. (St.) verticillata, F. v. M., fragm. IV, 94 (1864)	W.A.	—	—	—	—	—	—	—	B.fl.IV,20	M.fr.IV,94.
C. (St.) glandulosa, Salisbury, Parad. Lond. t. 77 (1807) ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,20	
C. (St.) laricifolia, Richard in Pers. syn. II, 210 (1807)	—	—	—	—	N.S.W.	Q.	—	—	B.fl.IV,21	
C. (St.) Preissii, F. v. M., fragm. III, 122 (1863)...	W.A.	—	—	—	—	—	—	—	B.fl.IV,21	M.fr.III,122.
C. (St.) imbricata, Bentham in Hueg. enum. 73 (1837)	W.A.	—	—	—	—	—	—	—	B.fl.IV,21	
C. (St.) adpressa, Bentham, Fl. Austr. IV, 22 (1869)	W.A.	—	—	—	—	—	—	—	B.fl.IV,22	
C. (St.) despecta, R. Brown, prodr. 571 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	—	B.fl.IV,22	M.fr.VI,58.
C. (St.) utricularioides, Bentham in Hueg. enum. 73 (1837) ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,23	
C. (St.) pygmaea, R. Brown, prodr. 571 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,23	
C. (St.) longitubea, Bentham in Hueg. enum. 73 (1837)	W.A.	—	—	—	—	—	—	—	B.fl.IV,23	
C. (St.) diffusa, R. Brown, prodr. 571 (1810)	—	—	—	—	—	Q.	—	—	B.fl.IV,23	
C. (St.) fissiloba, F. v. M., fragm. I, 154 (1859)	—	—	—	—	—	—	N.A.	—	B.fl.IV,24	M.fr.I,154.
C. (St.) trichopoda, F. v. M., fragm. X, 86 (1877)	—	—	—	—	—	Q.	—	—	—	M.fr.X,86.
C. (St.) alsinoides, R. Brown, prodr. 572 (1810)	—	—	—	—	—	Q.	N.A.	—	B.fl.IV,24	M.fr.I,151;VI,6;X,58.
C. (St.) tenerrima, F. v. M., fragm. I, 150 (1859)...	—	—	—	—	—	—	N.A.	—	B.fl.IV,24	M.fr.I,150.
C. (St.) brachyphylla, Sonder in Lehm. pl. Preiss. I, 386 (1845)	W.A.	—	—	—	—	—	—	—	B.fl.IV,24	
C. (St.) capillaris, R. Brown, prodr. 570 (1810)	—	—	—	—	—	Q.	—	—	B.fl.IV,25	
C. (St.) rotundifolia, R. Brown, prodr. 571 (1810)...	—	—	—	—	—	Q.	N.A.	—	B.fl.IV,25	M.fr.I,151.
C. (St.) schizantha, F. v. M., fragm. I, 152 (1859)	—	—	—	—	—	—	N.A.	—	B.fl.IV,25	M.fr.I,152.
C. (St.) lobuliflora, F. v. M., fragm. I, 153 (1859)...	—	—	—	—	—	—	N.A.	—	B.fl.IV,25	M.fr.I,153.
C. (St.) uliginosa, Swartz in Mag. nat. Fr. Berl. 52, t. 2 (1807)	—	—	—	—	—	Q.	—	—	B.fl.IV,26	
C. (St.) pulchella, Sonder in Lehm. pl. Preiss. I, 381 (1845) ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,26	
C. (St.) petiolaris, Sonder in Lehm. pl. Preiss. I, 382 (1845) ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,26	
C. (St.) emarginata, Sonder in Lehm. pl. Preiss. I, 383 (1845) ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,27	
C. (St.) corymbosa, R. Brown, prodr. 571 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,27	
C. (St.) lepida, F. v. M. in Benth. Fl. Austr. IV, 27 (1869) ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,27	
C. (St.) streptocarpa, Sonder in Lehm. pl. Preiss. I, 385 (1845)	W.A.	—	—	—	—	—	—	—	B.fl.IV,27	
C. (St.) uniflora, Sonder in Lehm. pl. Preiss. I, 381 (1845) ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,28	M.fr.VI,6.
C. (St.) pedunculata, R. Brown, prodr. 571 (1810)	—	—	—	—	—	Q.	N.A.	—	B.fl.IV,28	M.fr.VI,91.
C. (St.) pachyrrhiza, F. v. M., fragm. I, 152 (1859)	—	—	—	—	—	—	N.A.	—	B.fl.IV,28	M.fr.I,152.
C. (St.) muscicola, F. v. M., fragm. I, 153 (1859)...	—	—	—	—	—	—	N.A.	—	B.fl.IV,29	M.fr.I,153.
C. (St.) crassifolia, R. Brown, prodr. 571 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,29	
C. (St.) pycnostachya, Lindley, Bot. Regist. XXV, App. XXIX (1839)	W.A.	—	—	—	—	—	—	—	B.fl.IV,29	
C. (St.) pubigera, Sonder in Lehm. pl. Preiss. 1, 383 (1845) ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,30	
C. (St.) canaliculata, Lindley, Bot. Regist. XXV, App. XXIX (1839)	W.A.	—	—	—	—	—	—	—	B.fl.IV,30	
C. (St.) leptophylla, De Candolle, prodr. VII, 783 (1838) ..	W.A.	—	—	—	—	—	—	—	B.fl.IV,30	
C. (St.) dichotoma, De Candolle, prodr. VII, 783 (1838)... ...	W.A.	—	—	—	—	—	—	—	B.fl.IV,30	
C. (St.) bulbifera, Bentham in Hueg. enum. 73 (1837)	W.A.	—	—	—	—	—	—	—	B.fl.IV,31	
C. (St.) breviscapea, R. Brown, prodr. 572 (1810)...	W.A.	—	—	—	—	—	—	—	B.fl.IV,31	
C. (St.) eglandulosa, F. v. M., fragm. I, 150 (1859)	—	—	—	—	N.S.W.	Q.	—	—	B.fl.IV,31	M.fr.VI,6;X,58.
C. (St.) fasciculata, R. Brown, prodr. 572 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,32	
C. (St.) falcata, R. Brown, prodr. 572 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,32	
C. (St.) rhynchocarpa, Sonder in Lehm. pl. Preiss. I, 389 (1845)	W.A.	—	—	—	—	—	—	—	B.fl.IV,33	
C. (St.) adnata, R. Brown, prodr. 572 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,33	M.fr.X,58.
LEEWENHOEKIA, R. Brown, prodr. 572 (1810). (Levenhookia, Coleostylis.)										
L. pusilla, R. Brown, prodr. 573 (1810)	W.A.	—	—	—	—	—	—	—	B.fl.IV,34	M.fr.X.58.
L. dubia, Sonder in Lehm. pl. Preiss. I, 392 (1845)	W.A.	S.A.	T.	V.	N.S.W.	—	—	—	B.fl.IV,34	M.fr.III,121;X,58.
L. Sonderi, F. v. M., fragm. I, 18 (1858)	—	—	—	V.	—	—	—	—	B.fl.IV,35	
L. pauciflora, Bentham in Hueg. enum. 74 (1837)...	W.A.	—	—	—	—	—	—	—	B.fl.IV,35	M.fr.VI,77.
L. leptantha, Bentham, Fl. Austr. IV, 35 (1869)	W.A.	—	—	—	—	—	—	—	B.fl.IV,35	
L. stipitata, F. v. M., fragm. IV, 94 (1864)	W.A.	—	—	—	—	—	—	—	B.fl.IV,36	M.fr.IV,94.
L. Preissii, F. v. M., fragm. IV, 94 (1864)...	W.A.	—	—	—	—	—	—	—	B.fl.IV,36	M.fr.IV,94.

PHYLLACHNE, R. & G. Forster, char. gen. 115, t. 58 (1776). (Forstera, Helophyllum, Oreostylidium.)

P. bellidifolia, F. v. M., fragm. VIII, 39 (1873)	—	—	T.	—	—	—	—	—			M.fr.X,58.

DONATIA, R. et G. Forster, char. gen. 9, t. 5 (1776).

D. Novae Zealandiae, J. Hooker, Fl. N. Zeal. I, 81, t. 20 (1853)	—	—	T.	—	—	—	—	—		B.fl.II,450	M.fr.VIII,46;X,58.

GOODENIACEAE.

R. Brown, prodr. 573 (1810).

BRUNONIA, Smith in Transact. Linn. Soc. X, 366 (1809).

B. australis, Smith in Transact. Linn. Soc. X, 367, t. 28 (1809)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—			B.fl.IV,121	M.fr.X,111.

DAMPIERA, R. Brown, prodr. 587 (1810). (Linschotenia.)

D. luteifolia, F. v. M., fragm. X, 11 (1876)	W.A.	—	—	—	—	—	—			—	M.fr.X,11.
D. Linschotenii, F. v. M., fragm. VI, 28 (1866)	—	—	—	—	—	Q.	—			B.fl.IV,108	M.fr.VI,28.
D. spicigera, Bentham, Fl. Austr. IV, 109 (1869)...	W.A.	—	—	—	—	—	—			B.fl.IV,109	
D. candicans, F. v. M., fragm. X, 86 (1876)	W.A.	S.A.	—	—	—	—	—			—	M.fr.X,86.
D. teres, Lindley, Bot. Regist. XXV, App. XXVII (1839) ...	W.A.	—	—	—	—	—	—			B.fl.IV,109	
D. trigona, De Vriese in Lehm. pl. Preiss. I, 401 (1845)... ...	W.A.	—	—	—	—	—	—			B.fl.IV,109	M.fr.II,17;VIII,245;X,11
D. prostrata, De Vriese, Goodenov. 83 (1854)	W.A.	—	—	—	—	—	—			B.fl IV,110	
D. alata, Lindley, Bot. Regist. XXV, App. XXVII (1839) ...	W.A.	—	—	—	—	—	—			B.fl.IV,110	M.fr.X,11.
D. coronata, Lindley, Bot. Regist. XXV, App. XXVII (1839)...	W.A.	—	—	—	—	—	—			B.fl.IV,110	
D. carinata, Bentham, Fl. Austr. IV, 111 (1869)	W.A.	—	—	—	—	—	—			B.fl.IV,111	
D. sacculata, F. v. M. in Benth. Fl. Austr. IV, 111 (1869) ...	W.A.	—	—	—	—	—	—			B.fl.IV,111	
D. incana, R. Brown, prodr. 588 (1810)	W.A.	—	—	—	—	—	—			B.fl.IV,111	M.fr.II,17.
D. hederacea, R. Brown, prodr. 588 (1810)...	W.A.	—	—	—	—	—	—			B.fl.IV,112	M.fr.X,12.
D. ferruginea, R. Brown, prodr. 588 (1810)	—	—	—	—	—	Q.	—			B.fl.IV,112	M.fr.VI,29.
D. Brownii, F. v. M., fragm. VI, 29 (1866)	—	—	—	V.	N.S.W.	—	—			B.fl.IV,112	M.fr.VI,29.
D. lanceolata, Cunningham in De Cand., prodr. VII, 503 (1839)	—	S.A.	—	V.	N.S.W.	—	—			B.fl.IV,113	M.fr.VI,29.
D. altissima, F. v. M. in Benth. Fl. Austr. IV, 113 (1869) ...	W.A.	—	—	—	—	—	—			B.fl.IV,113	
D. marifolia, Bentham, Fl. Austr. IV, 114 (1869)...	—	S.A.	—	V.	N.S.W.	—	—			B.fl.IV,114	
D. rosmarinifolia, Schlechtendal, Linnaea XX, 603 (1847) ...	—	S.A.	—	V.	N.S.W.	—	—			B.fl.IV,114	M.fr.VI,29;X,12.
D. lavandulacea, Lindley, Bot. Regist. XXV, App. XXVII (1839)	W.A.	—	—	—	—	—	—			B.fl.IV,114	
D. juncea, Bentham, Fl. Austr. IV, 115 (1869)	W.A.	—	—	—	—	—	—			B.fl.IV,115	
D. oligophylla, Bentham, Fl. Austr. IV, 115 (1869)	W.A.	—	—	—	—	—	—			B.fl.IV,115	
D. loranthifolia, F. v. M. in Benth. Fl. Austr. IV, 115 (1869) ...	W.A.	—	—	—	—	—	—			B.fl.IV,115	
D. stricta, R. Brown, prodr. 589 (1810)	—	S.A.	T.	V.	N.S.W.	—	—			B.fl.IV,116	M.fr.X,12;XI,121.
D. Scottiana, F. v. M., fragm. XI, 120 (1881)	—	—	—	—	N.S.W.	—	—			—	M.fr.XI,120.
D. leptoclada, Bentham, Fl. Austr. IV, 116 (1869)	W.A.	—	—	—	—	—	—			B.fl.IV,116	
D. fasciculata, R. Brown, prodr. 588 (1810)	W.A.	—	—	—	—	—	—			B.fl.IV,116	M.fr.XI,121.
D. subspicata, Bentham, Fl. Austr. IV, 117 (1869)	W.A.	—	—	—	—	—	—			B.fl.IV,117	
D. triloba, Lindley, Bot. Regist. XXV, App. XXVII (1839) ...	W.A.	—	—	—	—	—	—			B.fl.IV,117	
D. linearis, R. Brown, prodr. 588 (1810)	W.A.	—	—	—	—	—	—			B.fl.IV,117	
D. cuneata, R. Brown, prodr. 588 (1810)	W.A.	—	—	—	—	—	—			B.fl.IV,118	
D. sericantha, F. v. M. in Benth. Fl. Austr. IV, 118 (1869) ...	W.A.	—	—	—	—	—	—			B.fl.IV,118	
D. parvifolia, R. Brown, prodr. 589 (1810)...	W.A.	—	—	—	—	—	—			B.fl.IV,118	
D. glabrescens, Bentham, Fl. Austr. IV, 119 (1869)	W.A.	—	—	—	—	—	—			B.fl.IV,119	
D. adpressa, Cunningham in De Cand., prodr. VII, 503 (1839)...	—	—	—	—	N.S.W.	Q.	—			B.fl.IV,119	
D. diversifolia, De Vriese in Lehm. pl. Preiss. I, 403 (1845) ...	W.A.	—	—	—	—	—	—			B.fl.IV,119	
D. eriocephala, De Vriese, Goodenov. 118, t. 21 (1854)	W.A.	—	—	—	—	—	—			B.fl.IV,120	M.fr.VI,29;X,12.
D. Wellsiana, F. v. M., fragm. X, 12 (1876)	W.A.	—	—	—	—	—	—			—	M.fr.X,12.

DIASPASIS, R. Brown, prodr. 586 (1810).

D. filifolia, R. Brown, prodr. 587 (1810)	W.A.	—	—	—	—	—	—			B.fl.IV,104	M.fr.I,206;II,181;VIII,58

LESCHENAULTIA, R. Brown, prodr. 581 (1810). (Latourea.)

L. formosa, R. Brown, prodr. 581 (1810)	W.A.	—	—	—	—	—	—			B.fl.IV,40	M.fr.VI,226.
L. oblata, Sweet, flor. Australas. 26, 46, t. 46 (1828)	W.A.	—	—	—	—	--	--			—	M.fr.VI,26.
L. chlorantha, F. v. M., fragm. II, 20 (1860)	W.A.	—	—	—	—	—	—			B.fl.IV,40	M.fr.II,20.
L. linarioides, De Candolle, prodr. VII, 519 (1818)	W.A.	—	—	—	—	—	—			B.fl.IV,40	
L. tubiflora, R. Brown, prodr. 581 (1810)	W.A.	—	—	—	—	—	—			B.fl.IV,41	
L. superba, F. v. M., fragm. VI, 10 (1866)	W.A.	—	—	—	—	—	—			B.fl.IV,41	M.fr.VI,10.
L. striata, F. v. M., fragm. VIII, 245 (1874)	—	S.A.	—	—	—	—	—			—	M.fr.VIII,245.
L. acutiloba, Bentham, Fl. Austr. IV, 41 (1869)	W.A.	—	—	—	—	—	—			B.fl.IV,41	
L. laricina, Lindley, Bot. Regist. XXV, App. XXVII (1839) ...	W.A.	—	—	—	—	—	—			B.fl.IV,41	
L. hirsuta, F. v. M., fragm. VI, 9 (1866)	W.A.	—	—	—	—	—	—			B.fl.IV,42	M.fr.VI,9.
L. longiloba, F. v. M., fragm. VI, 10 (1866)	W.A.	—	—	—	—	—	—			B.fl.IV,42	M.fr.VI,10.
L. biloba, Lindley, Bot. Regist. XXV, App. XXVII (1839) ...	W.A.	—	—	—	—	—	—			B.fl.IV,42	
L. expansa, R. Brown, prodr. 581 (1810)	W.A.	—	—	—	—	—	—			B.fl.IV,42	
L. floribunda, Bentham in Hueg. enum. 70 (1837)...	W.A.	—	—	—	—	—	—			B.fl.IV,43	
L. heteromera, Bentham, Fl. Austr. IV, 43 (1869)	W.A.	—	—	—	—	—	—			B.fl.IV,43	
L. divaricata, F. v. M., fragm. III, 33, 167 (1863)	—	S.A.	—	—	—	—	—			B.fl.IV,43	M.fr.VIII,246.
L. filiformis, R. Brown, prodr. 581 (1810)	—	—	—	—	—	Q.	N.A.			B.fl.IV,44	M.fr.VI,9,249.
L. agrostophylla, F. v. M., fragm. VI, 8, pl. XLVII (1866) ...	—	—	—	—	—	—	N.A.			B.fl.IV,44	M.fr.VI,8;249.

ANTHOTIUM, R. Brown, prodr. 582 (1810).

A. humile, R. Brown, prodr. 582 (1810)	W.A.	—	—	—	—	—	—			B.fl.IV,44	M.fr.VIII,58.
A. rubriflorum, F. v. M. in Benth. Fl. Austr. IV, 45 (1869) ...	W.A.	—	—	—	—	—	—			B.fl.IV,45	

CATOSPERMA, Bentham in Hooker, icon. pl. t. 1028 (1868).

C. Muelleri, Bentham in Hook. icon. pl. t. 1028 (1868)	—	—	—	—	—	—	N.A.			B.fl.IV,83	M.fr.I,121;VIII,58.

SCAEVOLA, Linné, mantiss. II, 145 (1771). (Pogonetes, Crossotoma, Temminckia, Kamphusia, Molkenboeria, Merkusia, Verreauxia.)

S. Koenigii, Vahl, symb. bot. III, 36 (1794)	—	—	—	—	—	Q.	N.A.			B fl.IV,86	M.fr.XI,76.
S. enantophylla, F. v. M., fragm. VIII, 58 (1873)...	—	—	—	—	—	Q.	—			—	M.fr.VI,225.

S. spinescens, R. Brown, prodr. 586 (1810)...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,87	M.fr.XI,20,27,76.
S. Groeneri, F. v. M., fragm. VI, 15 (1866)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,88	M.fr.VIII,58.
S. tomentosa, Gaudichaud in Freyc. Voy. Bot. 460, t. 81 (1826)	W.A.	—	—	—	—	—	—	B.fl.IV,88	
S. atriplicina, F. v. M., fragm. II, 18 (1860)	W.A.	—	—	—	—	—	—	B.fl.IV,88	M.fr.III,181.
S. striata, R. Brown, prodr. 586 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,89	
S. phlebopetala, F. v. M., fragm. II, 18 (1860)	W.A.	—	—	—	—	—	—	B.fl.IV,89	M.fr.II,VI,16;XI,76.
S. pilosa, Bentham in Hueg. enum. 69 (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,89	
S. hispida, Cavanilles, icon. et descr. pl. VI, 7, t. 510 (1801) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,90	M.fr.XI,76.
S. apterantha, F. v. M., fragm. I, 121 (1859)	—	—	—	V.	—	—	—	B.fl.IV,90	M.fr.I,121.
S. Hookeri, F. v. M. in Benth. Fl. Austr. IV, 90 (1869)... ...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,90	M.fr.XI,76.
S. parvifolia, F. v. M. in Benth. Fl. Austr. IV, 91 (1869) ...	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.IV,91	M.fr.VIII,58;X,59;XI,76.
S. oxyclona, F. v. M., fragm. X, 58 (1876)...	W.A.	—	—	—	—	—	—	—	M.fr.X,58.
S. restiacea, Bentham, Fl. Austr. IV, 91 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,91	[76.
S. depauperata, R. Brown, App. Sturt's Exped. 20 (1849) ...	—	S.A.	—	—	—	—	—	B.fl.IV,91	M.fr.III,33;VIII,58;XI,
S. tortuosa, Bentham, Fl. Austr. IV, 91 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,91	
S. Cunninghamii, De Candolle, prodr. VII, 508 (1838)	—	—	—	—	—	—	N.A.	B.fl.IV,92	
S. collaris, F. v. M., Rep. Babb. Exped. 15 (1858)	—	S.A.	—	—	—	—	—	B.fl.IV,92	M.fr.VIII,58;XI,76.
S. angulata, R. Brown, prodr. 586 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,92	
S. nitida, R. Brown, prodr. 584 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,93	
S. globulifiora, Labillardière, Nov. Holl. pl. spec. I, 55, t. 78 (1804)	W.A.	—	—	—	—	—	—	B.fl.IV,93	
S. porocarya, F. v. M., fragm. II, 19 (1860)	W.A.	—	—	—	—	—	—	B.fl.IV,94	M.fr.II,19;XI,76.
S. attenuata, R. Brown, prodr. 583 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,94	
S. glandulifera, De Candolle, prodr. VII, 510 (1838)	W.A.	—	—	—	—	—	—	B.fl.IV,94	
S. anchusifolia, Bentham in Hueg. enum. 68 (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,94	
S. holosericea, De Vriese in Lehm. pl. Preiss. I, 408 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,95	
S. suaveolens, R. Brown, prodr. 585 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,95	
S. revoluta, R. Brown, prodr. 586 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,96	
S. ovalifolia, R. Brown, prodr. 584 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,96	M.fr.VIII,58;XI,76.
S. crassifolia, Labillardière, Nov. Holl. pl. spec. I, 56, t. 79 (1804)	W.A.	S.A.	—	V.	—	—	—	B.fl.IV,96	M.fr.VIII,245;XI,20,76.
S. macrostachya, Bentham, Fl. Austr. IV, 97 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,97	M.fr.VIII,58;XI,76.
S. longifolia, De Vriese in Lehm. pl. Preiss. I, 410 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,97	
S. lanceolata, Bentham in Hueg. enum. 69 (1837)...	W.A.	—	—	—	—	—	—	B.fl.IV,97	M.fr.I,207;XI,76.
S. thesioides, Bentham in Hueg. enum. 68 (1837)...	W.A.	—	—	—	—	—	—	B.fl.IV,98	M.fr.XI,77.
S. macrophylla, Bentham, Fl. Austr. IV, 98 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,98	
S. platyphylla, Lindley, Bot. Regist. XXV, App. XXVI (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,98	M.fr.XI,77.
S. auriculata, Bentham, Fl. Austr. IV, 99 (1869)...	W.A.	—	—	—	—	—	—	B.fl.IV,99	M.fr.XI,77.
S. aemula, R. Brown, prodr. 584 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,99	M.fr.XI,77.
S. humilis, R. Brown, prodr. 585 (1810)	—	S.A.	—	—	—	—	—	B.fl.IV,100	
S. amblyanthera, F. v. M., fragm. I, 121 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,100	M.fr.I,121.
S. microphylla, Bentham, Fl. Austr. IV, 100 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,100	
S. cuneiformis, Labillard., Nov. Holl. pl. spec. I, 56, t. 80 (1804)	W.A.	—	—	—	—	—	—	B.fl.IV,101	M.fr.XI,77.
S. microcarpa, Cavanilles, icon. et. descr. pl. VI, 6, t. 509 (1801)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,101	M.fr.VI,16;XI,77.
S. linearis, R. Brown, prodr. 586 (1810)	—	S A.	—	—	—	—	—	B.fl.IV,102	
S. Oldfieldii, F. v. M., fragm. II, 19 (1860)	W.A.	—	—	—	—	—	—	B.fl.IV,102	M.fr.II,19;XI,77.
S. paludosa, R. Brown, prodr. 586 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,102	M.fr.XI,77.
S. sericophylla, F. v. M. in Benth. Fl. Austr. IV, 102 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,102	M.fr.XI,77.
S. canescens, Bentham in Hueg. enum. 69 (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,103	
S. humifusa, De Vriese in Lehm. pl. Preiss. I, 410 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,103	M.fr.XI,77.
S. fasciculata, Bentham in Hueg. enum. 68 (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,104	M.fr.XI,77.
S. stenophylla, Bentham, Fl. Austr. IV, 104 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,104	M.fr.I,113;XI,77.
S. Reinwardtii, De Vriese in Lehm. pl. Preiss. I, 409 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,106	
S. Verreauxii, F. v. M., Bot. Teachings, 65 (1877)	W.A.	—	—	—	—	—	—	B.fl.IV,105	
SELLIERA, Cavanilles in Anal. Cienz. Nat. I, 41, t. 5 (1799).									
S. radicans, Cavanilles, icon. et descr. pl. V, 49, t. 474 (1799) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,82	M.fr.VIII,58.
S. exigua, F. v. M., fragm. III, 142 (1862)...	W.A.	—	—	—	—	—	—	B.fl.IV,82	M.fr.III,142.
CALOGYNE, R. Brown, prodr. 579 (1810). (Distylis.)									
C. pilosa, R. Brown, prodr. 579 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,81	M.fr.VI,7,28.
C. Berardiana, F. v. M., fragm. VI, 7 (1867)	W.A.	—	—	—	—	—	N.A.	B.fl.IV,81	M.fr.VI,7;VIII,58.
C. heteroptera, F. v. Mueller, fragm. X, 43 (1876)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,43.
C. purpurea, F. v. M., fragm. VIII, 57 (1873)	—	—	—	—	—	—	N.A.	—	M.fr.VIII,57.
GOODENIA, Smith in Transact. Linn. Soc. II, 347 (1794). (Goodenoughia, Tetraphylax, Picrophyta, Stekhovia, Aillya.)									
G. phylicoides, F. v. M., fragm. I, 206 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,55	M.fr.I,206.
G. viscida, R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,55	M.fr.I,120;III,35.
G. xanthotricha, De Vriese, Gooden. 155 (1854)	W.A.	—	—	—	—	—	—	B.fl.IV,56	M.fr.VI,13.
G. scapigera, R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,56	M.fr.I,114;III,163;XI,50.
G. quadrilocularis, R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,57	M.fr.III,141;IV,13.
G. Ramelii, F. v. M., fragm. III, 20, t. 17 (1862)...	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,57	M.fr.VIII,245;X,13.
G. pinifolia, De Vriese, Gooden. 157, t. 30 (1854)...	W.A.	—	—	—	—	—	—	B.fl.IV,58	
G. decurrens, R. Brown, prodr. 575 (1810)...	—	—	—	—	N.S.W.	—	—	B.fl.IV,58	
G. racemosa, F. v. M., fragm. I, 114 (1859)	—	—	—	—	—	Q.	—	B.fl.IV,58	M.fr.I,114.
G. bellidifolia, Smith in Transact. Linn. Soc. II, 349 (1794) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,58	M.fr.XI,50.
G. stelligera, R. Brown, prodr. 575 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,59	M.fr.VI,14.
G. ovata, Smith in Transact. Linn. Soc. II, 347 (1794)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,59	M.fr.X,110.
G. amplexans, F. v. M. in Transact. phil. Inst. Vict. II, 70 (1859)	—	S.A.	—	V.	—	—	—	B.fl.IV,60	M.fr.VI,15.
G. strophiolata, F. v. M., fragm. I, 119 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,60	M.fr.I,119.
G. varia, R. Brown, prodr. 576 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,61	M.fr.I,205.
G. laevis, Bentham, Fl. Austr. IV, 61 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,61	
G. barbata, R. Brown, prodr. 576 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,61	M.fr.II,110,176;VI,15.
G. arthrotricha, F. v. M. in Benth. Fl. Austr. IV, 62 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,62	
G. disperma, F. v. M., fragm. I, 113 (1859)	—	—	—	—	—	Q.	—	B.fl.IV,62	M.fr.IV,145;VI,15.

G. geniculata, R. Brown, prodr. 577 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,62	M.fr.VI,14.
G. hederacea, Smith in Transact. Linn. Soc. II, 349 (1794) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,63	M.fr.VI,15,110.
G. xanthosperma, F. v. M., fragm. X, 12 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,12.
G. hirsuta, F. v. M., fragm. III, 35 (1862)...	—	S.A.	—	—	—	—	N.A.	B.fl.IV,64	M.fr.III,35.
G. heterophylla, Smith in Transact. Linn. Soc. II, 349 (1794) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,64	
G. glabra, R. Brown, prodr. 577 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,64	M.fr.VI,14.
G. strongylophylla, F. v. M., fragm. VI, 12 (1867)	—	—	—	—	—	Q.	—	B.fl.IV,65	M.fr.VI,12.
G. rotundifolia, R. Brown, prodr. 576 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,65	M.fr.VI,15.
G. azurea, F. v. M., fragm. I, 117 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,66	M.fr.I,117.
G. scaevolina, F. v. M., fragm. I, 118 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,66	M.fr.I,118.
G. Stobbsiana, F. v. M., fragm. XI, 49 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,49,137.
G. incana, R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,66	M.fr.I,155;III,141. [50.
G. leptoclada, Bentham, Fl. Austr. IV, 67 (1869)...	W.A.	—	—	—	—	—	—	B.fl.IV,67	M.fr.VI,227,VIII,186;XI,
G. Eatoniana, F. v. M., fragm. VIII, 186 (1874)	W.A.	—	—	—	—	—	—	—	M.fr.XI,51.
G. coerulea, R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,67	M.fr.III,140;XI,20.
G. trichophylla, De Vriese in Benth. Fl. Austr. IV, 67 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,67	
G. Hassallii, F. v. M., fragm. VI, 10, t. 51 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,68	M.fr.XI,50.
G. pterygosperma, R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,68	M.fr.I,155;III,163.
G. Vilmoriniae, F. v. M., fragm. III, 19, t. 16 (1862)	—	S.A.	—	—	—	—	N.A.	B.fl.IV,68	M.fr.III,19;VIII,245.
G. Bonneyana, F. v. M. in Benth. Fl. Austr. IV, 69 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,69	M.fr.VI,226,250.
G. calcarata, F. v. M., fragm. VI, 14 (1867)	—	S.A.	—	—	N.S.W.	—	—	B.fl.IV,69	M.fr.VIII,245.
G. albiflora, Schlechtendal, Linnaea XX, 599 (1847)	—	S.A.	—	—	—	—	—	B.fl.IV,70	
G. Nicholsoni, F. v. M., fragm. I, 203, t. 4 (1859)	—	S.A.	—	—	—	—	—	B.fl.IV,69	M.fr.I,203.
G. Macmillani, F. v. M., fragm. I, 119, t. 5 (1859)	—	—	—	V.	—	—	—	B.fl.IV,69	M.fr.I,119.
G. grandiflora, Sims, Bot. Mag. t. 890 (1806)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,69	M.fr.I,204;VI,14.
G. Chambersii, F. v. M., fragm. I, 204 (1859)	—	S.A.	—	—	—	—	—	B.fl.IV,70	M.fr.I,204.
G. Strangfordii, F. v. M., fragm. VI, 11, t. 52 (1867)	—	—	—	—	—	Q.	N.A.	B.fl.IV,70	
G. Mitchellii, Bentham, Fl. Austr. IV, 71 (1869)...	—	S.A.	—	—	—	Q.	—	B.fl.IV,71	M.fr.XI,75.
G. Mueckeana, F. v. M., fragm. VIII, 56 (1873)	—	S.A.	—	—	—	—	—	—	M.fr.IX,194;XI,133.
G. melanoptera, F. v. M., fragm. I, 115 (1859)	—	—	—	—	—	—	N.A.	—	M.fr.I,115.
G. heterochila, F. v. M., fragm. III, 142 (1863)	—	S.A.	—	—	—	—	N.A.	B.fl.IV,71	M.fr.VIII,245.
G. sepalosa, F. v. M. in Benth. Fl. Austr. IV, 72 (1869)... ...	—	—	—	—	—	—	N.A.	B.fl.IV,72	
G. hispida, R. Brown, prodr. 577 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,72	M.fr.VI,12.
G. auriculata, Bentham, Fl. Austr. IV, 72 (1869)...	—	—	—	—	—	—	N.A.	B.fl.IV,72	
G. Armstrongiana, De Vriese, Goodenov. 138, t. 24 (1854) ...	—	—	—	—	—	—	N.A.	B.fl.IV,73	M.fr.I,205;VII,40.
G. corynocarpa, F. v. M., fragm. II, 16 (1860)	W.A.	—	—	—	—	—	—	B.fl.IV,73	M.fr.II,16;VI,15.
G. mollissima, F. v. M. in Benth. Fl. Austr. IV, 73 (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,73	
G. cycloptera, R. Brown in App. Sturt's Exped. 20 (1849) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,74	M.fr.VI,14;XI,51.
G. tenella, R. Brown, prodr. 577 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,74	M.fr.XI,51. [X,110.
G. elongata, Labillardière, Nov. Holl. pl. spec. I, 52, t. 75 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,74	M.fr.VI,14;VIII,142,245;
G. pinnatifida, Schlechtendal, Linnaea XXI, 440 (1848)... ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,75	M.fr.VI,14;XI,51.
G. coronopifolia, R. Brown, prodr. 577 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,75	
G. heteromera, F. v. M., fragm. I, 115 (1859)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,76	M.fr.VI,14;XI,27.
G. concinna, Bentham, Fl. Austr. IV, 76 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,76	
G. glauca, F. v. M. in Transact. Vict. Inst. 40 (1855)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,76	M.fr.VI,14;XI,27.
G. filiformis, R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,77	M.fr.VI,14;VIII,245.
G. Armitiana, F. v. M., fragm. X, 110 (1877)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,110.
G. microptera, F. v. M., fragm. III, 34 (1862)	—	S.A.	—	—	—	—	N.A.	B.fl.IV,77	M.fr.VI,227;XI,75.
G. bicolor, F. v. M. in Benth. Fl. Austr. IV, 80 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,80	[245;X,110.
G. humilis, R. Brown, prodr. 575 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,79	M.fr.VII,227;VIII,142,
G. paniculata, Smith in Transact. Linn. Soc. II, 348 (1794) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,78	M.fr.I,116.
G. purpurascens, R. Brown, prodr. 578 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,78	M.fr.I,117.
G. gracilis, R. Brown, prodr. 575 (1810)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,79	M.fr.XI,51.
G. lamprosperma, F. v. M., fragm. I, 116 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,79	M.fr.I,116.
G. minutiflora, F. v. M., fragm. VIII, 244 (1874)	—	—	—	—	—	—	N.A.	—	M.fr.VIII,244.
G. claytoniacea, F. v. M. in Benth. Fl. Austr. IV, 79 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,79	M.fr.II,111.
G. Pumilio, R. Brown, prodr. 579 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,80	M.fr.X,111.
VELLEYA, Smith in Transact. Linn. Soc. IV, 217 (1798).	(Velleia, Euthales.)								
V. Daviesii, F. v. M., fragm. X, 10 (1876)...	W.A.	—	—	—	—	—	—	—	M.fr.X,10.
V. panduriformis, Cunningham in Benth. Fl. Austr. IV, 46 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,46	M.fr.XI,73.
V. connata, F. v. M. in Transact. phil. Soc. Vict. I, 18 (1855)...	—	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.IV,46	M.fr.VI,227;VIII,246.
V. perfoliata, R. Brown, prodr. 581 (1810)...	—	—	—	—	N.S.W.	—	—	B.fl.IV,46	
V. discophora, F. v. M., fragm. X, 10 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,10.
V. trinervis, Labillard., Nov. Holl. pl. spec. I, 54, t. 77 (1804)	W.A.	—	—	—	—	—	—	B.fl.IV,47	M.fr.XI,75.
V. macrophylla, Bentham, Fl. Austr. IV, 47 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,47	M.fr.XI,75.
V. paradoxa, R. Brown, prodr. 580 (1810)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,48	M.fr.VI,11;XI,75.
V. cycnopotamica, F. v. M., fragm. VI, 7 (1867)...	W.A.	—	—	—	—	—	—	B.fl.IV,48	M.fr.VI,249;X,10.
V. lyrata, R. Brown, prodr. 580 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,49	
V. macrocalyx, De Vriese, Goodenov. 176, t. 34 (1854)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,49	M.fr.X,75.
V. spathulata, R. Brown, prodr. 580 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,50	M.fr.XI,75.
V. pubescens, R. Brown, prodr. 581 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,49	M.fr.VI,11;XI,75.
V. montana, J. Hooker in Lond. Journ. of Bot. VI, 265 (1847)...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,50	

SYNPETALEAE HYPOGYNAE.

F. v. M. in Woolls, pl. of the neighb. of Sydney 34 (1880).

GENTIANEAE.

Necker in Act. Acad. Theod. Pal. II, 477 (1770), from B. de Jussieu (1759).

LIMNANTHEMUM, Gmelin, Nov. Comm. Acad. Petrop. XIV, 257 (1770). (Villarsia, Liparophyllum.)

L. Indicum, Thwaites, enum. pl. Ceyl. 205 (1862)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,378	M.fr.VI,136.
L. Moonii, Thwaites, enum. pl. Ceyl. 205 (1862)	—	—	—	—	—	Q.	—	—	M.fr.IX,163.
L. minimum, F. v. M., fragm. I, 40 (1858)	—	—	—	—	—	—	N.A.	B.fl.IV,379	M.fr.VI,137. [165.
L. crenatum, F. v. M. in Transact. phil. Soc. Vict. I, 17 (1854)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,379	M.fr.IV,127;VI,137;IX,
L. geminatum, Grisebach, gen. et spec. Gent. 346 (1839)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,380	M.fr.VI,137,254.
L. hydrocharoides, F. v. M. in Benth. Fl. Austr. IV, 380 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,380	M.fr.VI,139;IX,165.
L. exiliflorum, F. v. M., fragm. V, 46 (1865)	—	—	—	—	—	Q.	—	B.fl.IV,381	M.fr.VI,137.
L. exiguum, F. v. M., fragm. I, 40 (1858)	—	—	T.	—	—	—	—	B.fl.IV,381	M.fr.VI,137.
L. Gunnii, J. Hooker, Fl. Tasman. II, 368 (1860)	—	—	T.	—	—	—	—	B.fl.IV,382	M.fr.VI,137;IX,164.
L. capitatum, F. v. M., fragm. IX, 164 (1875)	W.A.	—	—	—	—	—	—	B.fl.IV,375	M.fr.VI,142.
L. calthifolium, F. v. M., fragm. IX, 164 (1875)	W.A.	—	—	—	—	—	—	B.fl.IV,374	M.fr.VI,140.
L. congestiflorum; Villarsia, F. v. M., fragm. VI, 141 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,375	M.fr.VI,141.
L. latifolium, F. v. M., fragm. IX, 164 (1875)	W.A.	—	—	—	—	—	—	B.fl.IV,375	M.fr.VI,141.
L. exaltatum, F. v. M., fragm. IX, 165 (1875)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,376	M.fr.II,21;VI,139.
L. parnassifolium, F. v. M., fragm. IX, 165 (1875)	W.A.	—	—	—	—	—	—	B.fl.IV,376	M.fr.VI,140.
L. violifolium; Villarsia, F. v. M., fragm. VI, 138 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,377	M.fr.VI,138.
L. lasiospermum, F. v. M., fragm. IX, 165 (1875)	W.A.	—	—	—	—	—	—	B.fl.IV,377	M.fr.VI,137.
L. albiflorum; Villarsia, F. v. M., fragm. II, 21 (1860)	W.A.	—	—	—	—	—	—	B.fl.IV,377	M.fr.VI,138.

SEBAEA, Solander in R. Brown, prodr. 451 (1810).

S. ovata, R. Brown, prodr. 452 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,371	M.fr.VII,136;IX,165.
S. albidiflora, F. v. M. in Transact. phil. Soc. Vict. I, 46 (1854)	—	S.A.	T.	V.	—	—	—	B.fl.IV,371	M.fr.VI,136.

ERYTHRAEA, Reneaulme, specim. hist. plant. 77, t. 76 (1611). [20.

E. australis, R. Brown, prodr. 451 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,371	M.fr.VI,136;IX,165;XI,

CANSCORA, Lamarck, Encycl. method. I, 602 (1783). (Orthostemon.)

C. diffusa, R. Brown, prodr. 451 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,372	M.fr.VI,136;IX,165.

GENTIANA, Tournefort, inst. 80, t. 40 (1700), from Dioscorides.

G. saxosa, Forster in Svensk. Kongl. Vet. Ac. Handl. 183, t. 5 (1777)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,373	M.fr.VI,136,255;IX,165.

LOGANIACEAE.

R. Brown in App. Flind. Voy. 564 (1814).

STRYCHNOS, Linné, syst. nat. 8 (1735); Linné, spec. plant. 189 (1753).

S. lucida, R. Brown, prodr. 469 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,369	
S. psilosperma, F. v. M., fragm. IV, 44 (1863)	—	—	—	—	—	Q.	—	B.fl.IV,369	M.fr.VI,131.

FAGRAEA, Thunberg in Kongl. Svensk. Vetensk. Acad. Handl. 125 (1782).

F. racemosa, Jack in Roxb. Fl. Ind. ed. Carey. II, 35 (1824)	—	—	—	—	—	—	N.A.	B.fl.IV,367	M.fr.II,137.
F. Muelleri, Bentham, Fl. Austr. IV, 368 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,368	

MITREOLA, Linné, hort. Cliffort. 492 (1737).

M. oldenlandioides, Wallich, numeric. list 4350 (1828)	—	—	—	—	—	—	N.A.	B.fl.IV,349	M.fr.VI,131.

MITRASACME, Labillardière, Nov. Holl. pl. spec. I, 36, t. 49 (1804). (Mitragyne.)

M. Archeri, J. Hooker, Fl. Tasman. II, 368 (1860)	—	—	T.	—	—	—	—	B.fl.IV,351	
M. montana, J. Hooker, Fl. Tasman. I, 274, t. 88 (1859)	—	—	T.	V.	—	—	—	B.fl.IV,351	M.fr.VI,131.
M. serpillifolia, R. Brown, prodr. 454 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,352	
M. paludosa, R. Brown, prodr. 453 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,352	M.fr.I,134.
M. pilosa, Labillardière, Nov. Holl. pl. spec. I, 36, t. 49 (1804)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,352	M.fr.VI,131.
M. alsinoides, R. Brown, prodr. 453 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,353	
M. polymorpha, R. Brown, prodr. 452 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,353	M.fr.VI,131.
M. longiflora, F. v. M. in Benth. Fl. Austr. IV, 353 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,353	
M. elata, R. Brown, prodr. 453 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,354	M.fr.I,132.
M. exserta, F. v. M., fragm. I, 131 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,354	M.fr.I,131.
M. tenuiflora, Benth. Fl. Austr. IV, 354 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,354	
M. ambigua, R. Brown, prodr. 454 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,354	
M. nudicaulis, Reinwardt in Blume, Bijdr. 849 (1826)	—	—	—	—	—	—	N.A.	B.fl.IV,355	
M. connata, R. Brown, prodr. 454 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,355	M.fr.I,131.
M. laevis, Bentham in Journ. Linn. Soc. I, 93 (1857)	—	—	—	—	—	—	N.A.	B.fl.IV,355	M.fr.I,133.
M. Indica, Wight, icon. pl. Ind. or. IV, t. 1601 (1850)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,356	
M. stellata, R. Brown, prodr. 454 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,356	
M. Cunninghamii, Bentham, Fl. Austr. IV, 357 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,357	
M. pygmaea, R. Brown, prodr. 453 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,357	
M. lutea, F. v. M., fragm. I, 133 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,357	M.fr.IV,170.
M. multicaulis, R. Brown, prodr. 453 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,357	
M. laricifolia, R. Brown, prodr. 453 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,358	M.fr.VI,131.
M. prolifera, R. Brown, prodr. 453 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,358	
M. gentianea, F. v. M., fragm. I, 130 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,358	M.fr.I,130.
M. phascoides, R. Brown, prodr. 454 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,359	
M. paradoxa, R. Brown, prodr. 454 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,359	M.fr.VI,131.
M. distylis, F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854)	—	—	T.	V.	—	—	—	B.fl.IV,359	M.fr.VI,131.

GENIOSTOMA, R. & G. Forster, char. gen. 23, t. 12 (1776).

G. Australianum, F. v. M., fragm. V, 19 (1865)	—	—	—	—	—	Q.	—	B.fl.IV,367	M.fr.V,19.
G. petiolosum, Moore & F. v. M., fragm. VII, 28 (1869)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,193.

LOGANIA, R. Brown, prodr. 454 (1810). (Euosma 1808, restorable.)

L. longifolia, R. Brown, prodr. 456 (1810)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,361	
L. latifolia, R. Brown, prodr. 455 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,361	M.fr.VI,132.
L. crassifolia, R. Brown, prodr. 455 (1810)...	W.A.	S.A.	—	—	—	—	—	B.fl.IV,362	
L. ovata, R. Brown, prodr. 455 (1810)	W.A.	S.A.	—	V.	—	—	—	B.fl.IV,362	
L. buxifolia, F. v. M., fragm. VI, 132 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,362	M.fr.IV,132.
L. stenophylla, F. v. M., fragm. I, 128 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,362	M.fr.I,128.
L. micrantha, Bentham in Journ. Linn. Soc. I, 94 (1857) ...	W.A.	—	—	—	—	—	—	B.fl.IV,363	M.fr.VI,132.
L. linifolia, Schlechtendal, Linnaea XX, 605 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,363	M.fr.VI,131.
L. fasciculata, R. Brown, prodr. 456 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,363	M.fr.VI,131.
L. floribunda, R. Brown, prodr. 456 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,364	M.fr.VI,131.
L. cordifolia, Hooker in Mitch. Trop. Austr. 341 (1848)... ...	—	—	—	—	—	Q.	—	B.fl.IV,364	
L. nuda, F. v. M., fragm. I, 129 (1859)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,365	M.fr.III,163; VI,131.
L. spermacocea, F. v. M., fragm. VI, 134 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,365	M.fr.VI,134.
L. callosa, F. v. M., fragm. VI, 134 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,365	M.fr.VI,134.
L. campanulata, R. Brown, prodr. 456 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,365	M.fr.VI,133.
L. serpillifolia, R. Brown, prodr. 456 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,366	M.fr.VI,133.
L. pusilla, R. Brown, prodr. 456 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,366	M.fr.VI,131.

PLANTAGINEAE.

A. L. de Jussieu, gen. plant. 89 (1789).

PLANTAGO, Tournefort, inst. 126, t. 48 (1700), from l'Ecluse (1576).

P. varia, R. Brown, prodr. 424 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,139	M.fr.XI,20.
P. stellaris, F. v. M., fragm. II, 23 (1860)	—	—	—	V.	N.S.W.	—	—	—	M.fr.VIII,148.
P. Brownii, Rapin in Mém. de la Soc. Linn. Par. VI, 484 (1827)	—	—	T.	—	—	—	—	B.fl.V,141	M.fr.II,23.
P. Gunnii, J. Hooker in Lond. Journ. V, 446, t. 13 (1846) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.V,142	M.fr.VIII,148;XI,134.

PRIMULACEAE.

Ventenat, Tabl. du régn. végét. II, 285 (1799).

ANAGALLIS, Tournefort, inst. 142, t. 59 (1700), from Hippocrates and Dioscorides. (Centunculus, Micropyxis.)

A. pumila, Swartz, Fl. Ind. occ. I, 345 (1797)	—	—	—	—	—	Q.	N.A.	B.fl.IV,270	

LYSIMACHIA, Tournefort, inst. 141, t. 59 (1700), from Dioscorides.

L. salicifolia, F. v. M. in Benth. Fl. Austr. IV, 269 (1869) ...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,269	
L. Japonica, Thunberg, F. Japon. 83 (1784)	—	—	—	—	N.S.W.	—	—	B.fl.IV,269	

SAMOLUS, Tournefort, inst. 143, t. 60 (1700). (Sheffieldia.)

S. Valerandi, Linné, spec. plant. 171 (1753)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,271	
S. repens, Persoon, synops. plant. I, 171 (1805)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,271	M.fr.XI,20.
S. platyphyllus, F. v. M. inedit.	—	S.A.	—	—	—	—	N.A.	—	

MYRSINACEAE.

R. Brown, prodr. 532 (1810). (Primulacearum subordo.)

MAESA, Forskael, Fl. Aegypt. Arab. 66 (1775). (Baeobotrys.)

M. dependens, F. v. M., fragm. V, 107 (1865)	—	—	—	—	—	Q.	—	B.fl.IV,273	M.fr.V,107.
M. haplobotrys, F. v. M., fragm. V, 161 (1866)	—	—	—	—	—	Q.	—	B.fl.IV,273	M.fr.V,161.

SAMARA, Linné, mantiss. II, 144 (1771). (Choripetalum.)

S. Australiana, F. v. M., fragm. VI, 164 (1868)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,274	M.fr.IV,36.

MYRSINE, Linné, syst. nat. 8 (1735); Linné, gen. plant. 54 (1737).

M. urceolata, R. Brown, prodr. 534 (1810)...	—	—	—	—	—	Q.	—	B.fl.IV,274	
M. campanulata, F. v. M., fragm. VI, 235 (1868)... . ..	—	—	—	—	—	Q.	—	—	M.fr.VI,235. [48.
M. crassifolia, R. Brown, prodr. 534 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,275	M.fr.IV,81;VI,164;VIII,
M. platystigma, F. v. M., fragm. VIII, 48 (1873)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,48.
M. variabilis, R. Brown, prodr. 534 (1810)...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,275	
M. achradifolia, F. v. M., fragm. VI, 164 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,275	M.fr.VI,164.

ARDISIA, Swartz, nov. gen. et spec. 3 et 48 (1788).

A. Pseudo-Jambosa, F. v. M., fragm. IV, 81 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,276	M.fr.IV,81.
A. brevipedata, F. v. M., fragm. VI, 163 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,276	M.fr.VI,163.

AEGICERAS, Gaertner, de fruct. I, 216, t. 46 (1788).

A. majus, Gaertner, de fruct. I, 216, t. 46 (1788)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,277	

SAPOTACEAE.

A. L. de Jussieu, gen. 151 (1789), from B. de Jussieu (1759).

NIEMEYERA, F. v. M., fragm. VII, 114 (1870). (Chrysophyllum partly.)

N. prunifera, F. v. M., fragm. VII, 114 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,278	M.fr.VI,26.

AMORPHOSPERMUM, F. v. M., fragm. VII, 112 (1870).

A. antilogum, F. v. M., fragm. VII, 113 (1870)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.X,119.

LUCUMA, Molina, Saggio, 186 (1782). (Sersalisia.)

L. sericea, Bentham & J. Hooker, gen. pl. II, 654 (1876) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,279	M.fr.VII,112.
L. Galactoxylon, Bentham & J. Hooker, gen. pl. II, 654 (1876)	—	—	—	—	—	Q.	—	B.fl.IV,279	

SIDEROXYLON, Dillenius, hort. Eltham. 357, t. 265 (1732). (Achras partly, Sapota partly, Sersalisia partly).

S. Arnhemicum, Bentham & J. Hooker, gen. pl. II, 655 (1876)...	—	—	—	—	—	—	N.A.	B.fl.IV,280	
S. Pohlmanianum, Bentham & J. Hooker, gen. pl. II, 655 (1876)	—	—	—	—	—	Q.	—	B.fl.IV,281	M.fr.V,184.
S. xerocarpum, Bentham & J. Hooker, gen. pl. II, 655 (1876) ...	—	—	—	—	—	Q.	—	B.fl.IV,281	
S. chartaceum, Bentham & J. Hooker, gen. pl. II, 655 (1876) ...	—	—	—	—	—	Q.	—	B.fl.IV,281	

S. Richardi, F. v. M.; Sersalisia laurifolia, A. Rich. sert. Astrol. 84 t. 13 (1834) ... ... — — — — N.S.W. Q. — B.fl.IV,282
S. australe, Bentham & J. Hooker, gen. pl. II, 655 (1876) ... — — — — N.S.W. Q. — B.fl.IV,282
S. Brownii, F.v. M.; Sersalisia obovata, R. Brown, prodr. 530 (1810) — — — — — Q. — B.fl.IV,283
S. myrsinoides, Bentham & J. Hooker, gen. pl. II, 655 (1876) ... — — — — N.S.W. Q. N.A. B.fl.IV,283
S euphlebium; Achras, F. v. M., fragm. VII (1870) ... ... — — — — — Q. — — M.fr.VII,110.
S. Brownlessianum, F. v. M., fragm. VIII, 111 (1870) ... ... — — — — — Q. — — M.fr.VII,111.
S. Howeanum; Achras, F. v. M., fragm. IX, 72 (1875) ... ... — — — — N.S.W. — — — M.fr.IX,72.
S. costatum; Achras, Endlicher, prodr. fl. ins. Norf. 49 (1833) ... — — — — N.S.W. — — — M.fr.IX,72.

HORMOGYNE, A. De Candolle, prodr. VIII, 176 (1844).
H. cotinifolia, A. De Candolle, prodr. VIII, 176 (1844) ... ... — — — — N.S.W. Q. — B.fl.IV,284 M.fr.VII,111.

MIMUSOPS, Linné, Fl. Zeil. 57 (1747).
M. parvifolia, R. Brown, prodr. 531 (1810)... ... ... — — — — -- Q. N.A. B.fl.IV,284 M.fr.VI,178;VII,114.
M. Browniana, Bentham, Fl. Austr. IV, 285 (1869) .. ... — — — — — Q. — B.fl.IV,285

EBENACEAE.

Ventenat, Tabl. II, 443 (1799).

DIOSPYROS, Linné, gen. plant. 143 (1737). (Cargillia.)
D. cordifolia, Roxburgh, pl. Corom. I, 38, t. 50 (1795) ... ... — — — — — — N.A. B.fl.IV,286
D. hebecarpa, Cunningham in Benth. pl. Austr. IV, 286 (1869) — — — — — Q. N.A. B.fl.IV,286
D. maritima, Blume, Bijdrag. 669 (1825) ... ... — — — — — Q. N.A. B.fl.IV,287 M.fr.V,163.
D. mabacea, F. v. M., Docum. intercol. Exhib. 35 (1866) ... — — — — N.S.W. Q. — B.fl.IV,288 M.fr.V,162.
D. Cargillia, F. v. M., Docum. intercol. Exhib. 35 (1866) ... — — — — N.S.W. Q. — B.fl.IV,288 M.fr.V,162.
D. pentamera, F. v. M. & Woolls, Doc. intercol. Exhib. 35 (1866) — — — — N.S.W. Q. — B.fl.IV,288 M.fr.V,163.

MABA, R. & G. Forster, char. gen. 121, t. 61 (1776).
M. laurina, R. Brown, prodr. 527 (1810) ... ... — — — — — Q. — B.fl.IV,289
M. rufa, Labillardière, sert. Austr. Caled. 33, t. 36 (1824) ... — — — — — Q. — B.fl.IV,289 M.fr.V,164;VI,253.
M. hemicycloides, F. v. M. in Benth. Fl. Austr. IV, 290 (1869) — — — — — Q. — B.fl.IV,290
M. fasciculosa, F. v. M., fragm. V, 163 (1866) ... ... — — — — N.S.W. Q. — B.fl.IV,290 M.fr.V,163;X,74.
M. compacta, R. Brown, prodr. 528 (1810)... ... ... — — — — — Q. N.A. B.fl.IV,290
M. reticulata, R. Brown, prodr. 528 (1810)... ... ... — — — — — Q. N.A. B.fl.IV,291
M. geminata, R. Brown, prodr. 527 (1810)... ... ... — — — — — Q. N.A. B.fl.IV,291
M. humilis, R. Brown, prodr. 527 (1810) ... . — — — — — Q. N.A. B.fl.IV,291 M.fr.X,74.
M. buxifolia, Persoon, synops. pl. II, 602 (1807) ... ... — — — — — — N.A. — M.fr.X,74.

STYRACEAE.

L. C. Richard, Analyse du fruit. 48 (1808).

SYMPLOCOS, N. J. Jacquin, enum. plant. Carib. 24 (1760).
S. Stawellii, F. v. M., fragm. V, 60 (1865)... ... ... — — — — N.S.W. Q. — B.fl.IV,292 M.fr.V,60.
S. Thwaitesii, F. v. M., fragm. III, 22 (1862) ... ... — — — — N.S.W. Q. — B.fl.IV,293 M.fr.V,211.

JASMINEAE.

A. L. de Jussieu, gen. 104 (1789), from Necker (1770).

JASMINUM, Tournefort, inst. 597, t. 368 (1700), from l'Ecluse (1611). [43.
J. didymum, G. Forster, prodr. 3 (1786) ... ... — — — — N.S.W. Q. N.A. B.fl.IV,294 M.fr.IV,150;VI,87;VIII,
J. racemosum, F. v. M., fragm. I, 19 (1858) ... ... — — — — — Q. — B.fl.IV,295 M.fr.I,19;VI,87;VIII,43.
J. lineare, R. Brown, prodr. 521 (1810) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.IV,295 M.fr.VI,87;VIII,43.
J. simplicifolium, G. Forster, prodr. 3 (1786) ... ... — — — — N.S.W. Q. N.A. B.fl.IV,296 M.fr.VI,87;VIII,43.
J. aemulum, R. Brown, prodr. 521 (1810) ... ... — — — — — Q. N.A. B.fl.IV,296 M.fr.VI,86. [43.
J. calcareum, F. v. M., fragm. I, 212 (1859) ... ... W.A. S.A. — — — — N.A. B.fl.IV,297 M.fr.I,212;VI,87,VIII,
J. suavissimum, Lindley in Mitch. Trop. Austr. 355 (1848) ... — — — — N.S.W. Q. — B.fl.IV,297 M.fr.I,183;VI,87.

OLEA, Tournefort, inst. 598, t. 370 (1700), from Plinius.
O. paniculata, R. Brown, prodr. 523 (1810) ... ... — — — — N.S.W. Q. — B.fl.IV,297 M.fr.VIII,43.
O. apetala, Vahl, symbol. III, 3 (1794) ... ... — — — — N.S.W. — — — M.fr.VIII,43;IX,169.

LIGUSTRUM, Tournefort, inst. 596, t. 367 (1700), from C. Bauhin (1623).
L. Australianum, F. v. M., fragm. V, 20 (1865) ... ... — — — — — Q. — B.fl.IV,298 M.fr.V,20.

NOTELAEA, Ventenat, Choix des pl. dans le jard. de Cels, t. 25 (1803).
N. ovata, R. Brown, prodr. 524 (1810) ... ... — — — — N.S.W. Q. — B.fl.IV,299 M.fr.VIII,43.
N. longifolia, Ventenat, Choix t. 25 (1803)... ... ... — — — — V. N.S.W. Q. — B.fl.IV,299 M.fr.VIII,43.
N. punctata, R. Brown, prodr. 524 (1810) ... ... — — — — — — Q. — B.fl.IV,300
N. microcarpa, R. Brown, prodr. 524 (1810) ... ... — — — — N.S.W. Q. — B.fl.IV,300 M.fr.VIII,43.
N. ligustrina, Ventenat, Choix t. 25 (1803) ... ... — — — T. V. N.S.W. — — B.fl.IV,300 M.fr.VIII,43,142.
N. linearis, Bentham, Fl. Austr. IV, 300 (1869) ... ... — — — — N.S.W. — — B.fl.IV,300

MAYEPEA, Aublet, Hist. des pl. de la Guian. I, 81, t. 31 (1775). (Chionanthus partly, Linociera partly, Ceranthes.)
M. ramiflora, F.v.M.; Chion. Roxb. Fl. Ind. I, 106 ed. Carey (1820) — — — — — Q. — B.fl.IV,301 M.fr.IV,83.
M. picrophloia; Chionanthus, F. v. M., fragm. III, 139, t. 24 (1863) — — — — — Q. — B.fl.IV,301 M.fr.IV,176.
M. axillaris, F. v. M.; Chionanthus, R. Brown, prodr. 523 (1810) — — — — — Q. N.A. B.fl.IV,301
M. quadristaminea, F. v. M., fragm. X, 89 (1876)... ... ... — — — — N.S.W. — — — M.fr.VIII,41.

APOCYNEAE.

A. L. de Jussieu, gen. plant. 143 (1789).

CHILOCARPUS, Blume, catal. hort. Buitenz. (1823).
C. australis, F. v. M., fragm. II, 90 (1861)... ... ... — — — — N.S.W. Q. — B.fl.IV,303 M.fr.II,90;VI,118.

MELODINUS, R. & G. Forster, char. gen. 37, t. 19 (1776).
M. acutiflorus, F. v. M. in Transact. phil. Soc. Vict. II, 71 (1857) — — — — — Q. — B.fl.IV,304
M. Guilfoylei, F. v. M., fragm. VI, 118 (1868) ... ... — — — — — Q. — B.fl.IV,304 M.fr.VI,118.
M. Baueri, Endlicher, prodr. pl. Norfolk. 59 (1833) ... ... — — — — N.S.W. — — — M.fr.IX,169.

CARISSA, Linné, mantiss, pl. 7 (1767).

C. laxiflora, Bentham, Fl. Austr. IV, 305 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,305	
C. Brownii, F. v. M., fragm. IV, 45 (1863)...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,305	M.fr.IV,45.

CERBERA, Linné, gen. plant. 62 (1737).

C. Odollam, Gaertner, de fruct. 193, t. 124 (1791)...	—	—	—	—	—	Q.	N.A.	B.fl.IV,306	M.fr.VI,117.

ALYXIA, Banks in R. Brown, prodr. 469 (1810). (Gynopogon.)

A. buxifolia, R. Brown, prodr. 470 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,307	M.fr.VI,117.
A. ruscifolia, R. Brown, prodr. 470 (1810)...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,308	M.fr.VI,117.
A. ilicifolia, F. v. M., fragm. IV, 149 (1864)	—	—	—	—	—	Q.	—	B.fl.IV,308	M.fr.IV,149.
A. squamulosa, C. Moore & F. v. M., fragm. VIII, 47 (1873) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.VI,117;VIII,47.
A. obtusifolia, R. Brown, prodr. 470 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,308	
A. Gynopogon, Roemer & Schultes, syst. veg. IV, 440 (1819) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,47.
A. spicata, R. Brown, prodr. 470 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,308	M.fr.VI,117.
A. Lindii, F. v. M., fragm. VIII, 46 (1873)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,46.
A. thyrsiflora, Bentham, Fl. Austr. IV, 309 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,309	
A. Thozetii, F. v. M., fragm. X, 103 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,103.

OCHROSIA, A. L. de Jussieu, gen. 143 (1789). (Lactaria.)

O. elliptica, Labillardière, sert. Austr. Caled. 27, t. 30 (1824) ...	—	—	—	—	—	Q.	—	B.fl.IV,310	M.fr.VI,118.
O. Moorei, F. v. M. in Benth. Fl. Austr. IV, 310 (1869)... ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,310	M.fr.VI,118.
O. Kilneri, F. v. M., fragm. VII, 129 (1871)	—	—	—	—	—	Q.	—	—	M.fr.VIII,129.

NOTONERIUM, Bentham in B. & J. H. gen. II, 698 (1876).

N. Gossei, Bentham in B. & J. H. gen. pl. 698 (1876)	—	S.A.	—	—	—	—	—	—	M.fr.X,103.

TABERNAEMONTANA, Plumier, nov. pl. Amer. gen. 18, t. 30 (1703).

T. orientalis, R. Brown, prodr. 468 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,311	M.fr.VI,117.
T. pubescens, R. Brown, prodr. 468 (1810)...	—	—	—	—	—	Q.	N.A.	B.fl.IV,311	M.fr.VI,118.

VINCA, Rivinus in Rupp. Fl. Jenens. 27 (1818), from Plinius. (Perhaps immigrated.)

V. rosea, Linné, spec. plant. ed. sec. 305 (1762)	—	—	—	—	—	Q.	—	—	M.fr.XI,123.

ALSTONIA, R. Brown in Mem. Wern. Soc. I, 75 (1809).

A. scholaris, R. Brown in Mem. Wern. Soc. I, 75 (1809) ...	—	—	—	—	—	Q.	—	B.fl.IV,312	M.fr.VI,117.
A. verticillosa, F. v. M., fragm. VI, 116 (1868)	—	—	—	—	—	Q.	N.A.	B.fl.IV,313	M.fr.VI,116.
A. ophioxyloides, F. v. M., fragm. I, 57 (1858)	—	—	—	—	—	—	N.A.	B.fl.IV,313	M.fr.VI,117.
A. villosa, Blume, Bijdr. 1038 (1826)	—	—	—	—	—	Q.	—	B.fl.IV,313	M.fr.VI,117.
A. linearis, Bentham, Fl. Austr. IV, 314 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,314	[117.
A. constricta, F. v. M., fragm. I, 57 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,314	M.fr.III,163;IV,170;VI,

ICHNOCARPUS, R. Brown in Mem. Wern. Soc. I, 75 (1809).

I. frutescens, R. Brown in Mem. Wern. Soc. I, 75 (1809) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,315	M.fr.VI,118.

WRIGHTIA, R. Brown in Mem. Wern. Soc. I, 75 (1809). (Balfouria.)

W. pubescens, R. Brown, prodr. 467 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,316	
W. saligna, F. v. M. in Benth. Fl. Austr. IV, 316 (1869) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,316	M.fr.V,161;VI,118.
W. Cunninghamii, Bentham, Fl. Austr. IV, 317 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,317	

PARSONSIA, R. Brown in Mem. Wern. Soc. I, 64 (1809).

P. lanceolata, R. Brown, prodr. 466 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,318	M.fr.VI,126.
P. velutina, R. Brown, prodr. 466 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,318	M.fr.VI,130.
P. Leichhardtii, F. v. M., fragm. VI, 128 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,319	M.fr.VI,128.
P. ventricosa, F. v. M. in Transact. phil. Inst. Vict. II, 71 (1857)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,319	M.fr.VI,130.

LYONSIA, R. Brown in Mem. Wern. Soc. I, 66 (1809).

L. lilacina, F. v. M. in Bentham Fl. Austr. IV, 321 (1869) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,321	M.fr.VI,127.
L. induplicata, F. v. M. in Benth. Fl. Austr. IV, 321 (1869) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,321	M.fr.VI,129.
L. straminea, R. Brown, prodr. 466 (1810)...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,321	M.fr.VI,129.
L. reticulata, F. v. M., Rep. Burdek. Exped. 16 (1860)... ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,321	M.fr.VI,129.
L. diaphanophleba, F. v. M. in Benth. Fl. Austr. IV, 322 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,322	M.fr.II,158;VI,130.
L. Langiana, F. v. M., fragm. VI, 128 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,322	M.fr.VI,128.
L. largiflorens, F. v. M. in Benth. Fl. Austr. IV, 322 (1869) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,322	
L. latifolia, Bentham, Fl. Austr. IV, 323 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,323	
L. oblongifolia, Bentham, Fl. Austr. IV, 323 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,323	
L. eucalyptifolia, F. v. M. in Benth. Fl. Austr. IV, 323 (1869)...	—	—	—	—	—	Q.	—	B.fl.IV,323	M.fr.II,159;VI,130

ASCLEPIADEAE.

N. J. Jacquin, Misc. Austr. I, 35 (1778).

GYMNANTHERA, R. Brown in Mem. Wern. Soc. I, 58 (1809).

G. nitida, R. Brown, prodr. 464 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,326	M.fr.V,160.

SECAMONE, R. Brown in Mem. Wern. Soc. I, 55 (1809).

S. elliptica, R. Brown, prodr. 464 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,327	M.fr.V,161.
S. ovata, R. Brown, prodr. 464 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,328	

VINCETOXICUM, Dodoens, stirp. hist. pemptades, 704 (1583). (Oxystelma partly, Cynoctonum.)

V. ovatum, Bentham, Fl. Austr. IV, 330 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,330	
V. elegans, Bentham, Fl. Austr. IV, 330 (1869)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,330	M.fr.V,160.
V. carnosum, Bentham, Fl. Austr. IV, 331 (1869)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,331	
V. leptolepis, Bentham, Fl. Austr. IV, 331 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,331	

CYNANCHUM, Linné, gen. plant. 63 (1737).

C. erubescens, R. Brown, prodr. 463 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,332	
C. floribundum, R. Brown, prodr. 463 (1810)	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,332	M.fr.V,160,
C. pedunculatum, R. Brown, prodr. 463 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,333	M.fr.V,160.
C. puberulum, F. v. M. in Benth. Fl. Austr. IV, 333 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,333	

SARCOSTEMMA, R. Brown in Mem. Wern. Soc. I, 50 (1809).

S. Australe, R. Brown, prodr. 463 (1810)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,328	M.fr.V,160;XI,80.

DAEMIA, R. Brown in Mem. Wern. Soc. I, 50 (1809). (Pentatropis, Rhyncharrhena.)

D. linearis, F. v. M.; Pentatropis, Decaisne in De Cand., prodr. VIII, 536 (1833)	W.A.	—	—	—	—	—	—	B.fl.IV,329	
D. atropurpurea; Rycharrh., F. v. M., fr. I, 128 (1859) ...	—	—	—	—	—	Q.	—	B.fl.IV,329	M.fr.I,128.
D. quinquepartita; Rycharrh., F. v. M., fr. I, 128 (1859) ..	—	—	—	V.	N.S.W.	—	—	B.fl.IV,329	M.fr.I,128;XI,80.
D. Kempeana, F. v. M. in Wing's S. Sc. Record, Aug. (1882) ...	—	S.A.	—	—	—	—	—	—	

GYMNEMA, R. Brown in Mem. Wern. Soc. I, 33 (1809). (Bidaria.)

G. sylvestre, R. Brown in Mem. Wern. Soc. I, 33 (1809) ...	—	—	—	—	—	—	N.A.	B.fl.IV,343	
G. pleiadenium, F. v. M., fragm. XI, 77 (1879)	—	—	—	—	—	Q.	—	—	M.fr.XI,77.
G. Muelleri, Bentham, Fl. Austr. IV, 343 (1869)...	—	—	—	—	—	—	N.A.	B.fl.IV,343	
G. brevifolium, Bentham, Fl. Austr. IV, 343 (1809)	—	—	—	—	—	Q.	—	B.fl.IV,343	
G. trinerve, R. Brown, prodr. 462 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,343	
G. stenophyllum, A. Gray in Proc. Am. Ac. of Sc. V, 335 (1864)	—	—	—	—	—	—	N.A.	B.fl.IV,344	

GONGRONEMA, Decaisne in De Candolle, prodr. VIII, 624 (1844).

G. micradenia, Bentham & J. Hooker, gen. plant. II, 770 (1876)	—	—	—	—	—	Q.	—	B.fl.IV,344	M.fr.XI,78.

TYLOPHORA, R. Brown in Mem. Wern. Soc. I, 28 (1809).

T. erecta, F. v. M. in Benth. Fl. Austr. IV, 334 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,334	M.fr.XI,79.
T. macrophylla, Bentham, Fl. Austr. IV, 334 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,334	
T. grandiflora, R. Brown, prodr. 460 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,334	M.fr.XI,80.
T. floribunda, Bentham, Fl. Austr. IV, 335 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,335	
T. calcarata, Bentham, Fl. Austr. IV, 335 (1869)...	—	—	—	—	—	Q.	—	B.fl.IV,335	
T. barbata, R. Brown, prodr. 460 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,335	M.fr.V,160;XI,79.
T. Woollsii, Bentham, Fl. Austr. IV, 335 (1869)	—	—	—	—	N.S.W.	—	—	B.fl.IV,335	
T. paniculata, R. Brown, prodr. 460 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,336	
T. enervis, F. v. M., fragm. IX, 70 (1875)...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,70.
T. flexuosa, R. Brown, prodr. 460 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,336	M.fr.XI,79.
T. biglandulosa, A. Gray in Proc. of Am. acad. sc. vol. V (1864)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.

MARSDENIA, R. Brown in Mem. Wern. Soc. I, 28 (1809). (Leichhardtia.)

M. cinerascens, R. Brown, prodr. 461 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,337	
M. rhyncholepis, F. v. M., fragm. XI, 78 (1879)	—	—	—	—	—	Q.	—	—	M.fr.XI,78.
M. flavescens, Cunningham in Bot. Mag., t. 3289 (1833) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,337	M.fr.V,160.
M. cymulosa, Bentham, Fl. Austr. IV, 338 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,338	
M. velutina, R. Brown, prodr. 461 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,338	
M. Hullsii, F. v. M. in Benth. Fl. Austr. IV, 338 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,338	
M. araujacea, F. v. M., fragm. VI, 135 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,339	M.fr.VI,135.
M. tubulosa, F. v. M., fragm. IX, 71 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,71.
M. rostrata, R. Brown, prodr. 461 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,339	M.fr.V,160;VI,135.
M. Fraseri, Bentham, Fl. Austr. IV, 339 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,339	
M. longiloba, Bentham, Fl. Austr. IV, 340 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,340	M.fr.XI,79.
M. suaveolens, R. Brown, prodr. 461 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,340	
M. leptophylla, F. v. M. in Benth. Fl. Austr. IV, 340 (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,340	
M. Leichhardtiana, F. v. M., fragm. V, 160 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,341	M.fr.V,160.
M. viridiflora, R. Brown, prodr. 461 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,341	M.fr.V,160.
M. coronata, Bentham, Fl. Austr. IV, 341 (1810)...	—	—	—	—	—	Q.	—	B.fl.IV,341	
M. microlepis, Bentham, Fl. Austr. IV, 342 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,342	

THOZETIA, F. v. M. in Benth. Fl. Austr. IV, 347 (1869).

T. racemosa, F. v. M. in Benth. Fl. Austr. IV, 347 (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,347	

HOYA, R. Brown in Mem. Wern. Soc. I, 26 (1809).

H. carnosa, R. Brown in Mem. Wern. Soc. I, 26 (1809)... ...	—	—	—	—	—	Q.	—	B.fl.IV,346	
H. australis, R. Brown in Transact. hort. Soc. VII, 28 (1826) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,346	M.fr.XI,80.
H. Nicholsoniae, F. v. M., fragm. V, 159 (1866)	—	—	—	—	—	Q.	—	B.fl.IV,347	M.fr.V,159;IX,70,190.

DISCHIDIA, R. Brown in Mem. Wern. Soc. I, 32 (1809).

D. nummularia, R. Brown, prodr. 461 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,345	M.fr.V,161.
D. Timorensis, Decaisne in nouv. annal. du Mus. III, 377, t. 17 (1835)	—	—	—	—	—	Q.	N.A.	—	

MICROSTEMMA, R. Brown in Mem. Wern. Soc. I, 25 (1809).

M. tuberosum, R. Brown, prodr. 459 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,345	M.fr.I,58.
M. glabriflorum, F. v. M., fragm. I, 58 (1858)	—	—	—	—	—	—	N.A.	B.fl.IV,345	M.fr.I,58.

CEROPEGIA, Linné, gen. plant. 65 (1737).

C. Cumingiana, Decaisne in De Cand., prodr. VIII, 643 (1844)	—	—	—	—	—	Q.	—	B.fl.IV,348	M.fr.V,159.

CONVOLVULACEAE.

A. L. de Jussieu, gen. plant. 132 (1789), from B. de Jussieu (1759).

ERYCIBE, Roxburgh, pl. Corom. III, 31, t. 159 (1798).

E. paniculata, Roxburgh, pl. Corom. III, 31, t. 159 (1798) ...	—	—	—	—	—	Q.	—	B.fl.IV,411	M.fr.VI,100.

LEPISTEMON, Blume, Bijdrag. 722 (1825).

L. urceolatus, F. v. M.; L. Fitzalani, F. v. M., fragm. X, 111 (1877)—	—	—	—	—	—	Q.	—	B.fl.IV,427	M.fr.X,111.

IPOMOEA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 47 (1737). (Pharbitis, Batatas, Calonyction, Aniseia, Skinnera.)

I. paniculata, R. Brown, prodr. 486 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,414	M.fr.VI,98; X,111.
I. Davenporti, F. v. M., fragm. VI, 97 (1868)	—	S.A.	—	—	—	—	N.A.	B.fl.IV,415	M.fr.VIII,17;X,111.
I. bona nox, Linné, spec. plant. edit. secund. 228 (1762) ...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IX,74;X,113;X,73.
I. palmata, Forskael, Fl. Aegypt. Arabic. 43 (1775)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,415	M.fr.VI,98; X,111.
I. quinata, R. Brown, prodr. 486 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,415	M.fr.VI,98; X,112.
I. diversifolia, R. Brown, prodr. 487 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,416	M.fr.X,112.

I. dissecta, Willdenow, phytogr. 5, t. 2 (1794)	—	—	—	—	—	Q.	N.A.	B.fl.IV,416	M.fr.VI,98; X,112.
I. sinuata, Ortega, decad. nov. pl. hort. Matrit. VII, 84 (1799)	—	—	—	—	—	Q.	—	—	M.fr.VIII,17; X,113.
I. hederacea, N. J. Jacquin, Collectan. I, 124 (1786)	—	—	—	—	—	Q.	N.A.	B.fl.IV,416	M.fr.VI,99; X,112.
I. congesta, R. Brown, prodr. 485 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,417	M.fr.VI,99; X,112.
I. purpurea, Roth, Catalect. bot. I, 36 (1797)	—	—	—	—	—	Q.	—	B.fl.IV,417	M.fr.VI,99; X,112.
I. peltata, Choisy, Convolvulac. orient. 70 (1834)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,418	
I. Calobra, Hill & F. v. M., fragm. XI, 73 (1879)	—	—	—	—	—	Q.	—	—	M.fr.XI,73,137.
I. alata, R. Brown, prodr. 484 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,418	M.fr.X,112.
I. Turpethum, R. Brown, prodr. 485 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,418	M,fr.VI,98.
I. longiflora, R. Brown, prodr. 484 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,418	M.fr.VI,98; IX,74; X,112.
I. costata, F. v. M. in Benth, Fl. Austr. IV, 419 (1869)	W.A	S.A.	—	—	—	—	N.A.	B.fl.IV,419	M.fr.X,112.
I. Pes Caprae, Roth, nov. spec. plant. 109	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,419	M.fr.VI,97; X,112.
I. carnosa, R. Brown, prodr. 485 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,420	
I. reptans, Poiret, Encycl. méth. suppl. III, 460 (1813)	W.A.	—	—	—	—	Q.	N.A.	B.fl.IV,420	M.fr.VI,97.
I. graminea, R. Brown, prodr. 485 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,421	
I. velutina, R. Brown, prodr. 485 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,421	
I. abrupta, R. Brown, prodr. 485 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,421	M.fr.X,112.
I. denticulata, Choisy in De Cand. prodr. IX, 379 (1845)	—	—	—	—	—	Q.	N.A.	B.fl.IV,421	M.fr I,185; X,112.
I. gracilis, R. Brown, prodr. 484 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,422	M.fr.I,185; X,112.
I. sepiaria, Koenig in Roxb. Fl. Indic. ed. Carey II, 90 (1824)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,422	
I. Muelleri, Bentham, Fl. Austr. IV, 423 (1869)	W.A.	—	—	—	—	—	N.A.	B.fl.IV,423	M.fr.X,112.
I. cymosa, Roemer & Schultes, syst. veg. IV, 241 (1819)	—	—	—	—	—	Q.	—	B.fl.IV,423	
I. linifolia, Blume, Bijdrag. 721 (1825)	—	—	—	—	—	Q.	—	B.fl.IV,423	
I. chryseides, Ker in Bot. Regist. t. 270 (1818)	—	—	—	—	—	Q.	—	B.fl.IV,423	M.fr.VI,98.
I. flava, F. v. M. in Benth. Fl. Austr. IV, 424 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,424	
I. obscura, Ker in Bot. Regist. t. 239 (1817)	—	—	—	—	—	Q.	—	B.fl.IV,424	
I. incisa, R. Brown, prodr. 486 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,424	M.fr.I,186.
I. uniflora, Roemer & Schultes, syst. IV, 247 (1819)	—	—	—	—	—	Q.	—	B.fl.IV,425	M.fr.VI,99.
I. angustifolia, N. J. Jacquin, Collect. II, 367 (1788)	—	—	—	—	—	Q.	N.A.	B.fl.IV,425	M.fr.VI,97; X,112.
I. plebeja, R. Brown, prodr. 484 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,426	M.fr.VI,99; X,112.
I. eriocarpa, R. Brown, prodr. 484 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,426	M.fr.VI,98; X,112.
I. heterophylla, R. Brown, prodr. 487 (1810)	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,426	M.fr.VI,98; X,112.
I. erecta, R. Brown, prodr. 487 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,427	M.fr.VI,98.
CONVOLVULUS, Tournefort, inst. 82, t. 17 (1700). (Calystegia.)									
C. erubescens, Sims, Bot. Magaz. t. 1067 (1808)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,429	M.fr.VI,99; X,113.
C. parviflorus, Vahl, symbol. bot. III, 29 (1794)	—	—	—	—	—	Q.	N.A.	B.fl.IV,430	M.fr.VI,99; X,113.
C. marginatus, Poiret, Encycl. méth. suppl. II (1811)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,430	M.fr.VI,99; X,113.
C. sepium, Linné, spec. plant. 153 (1753)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,430	M.fr.VI,99; X,113.
POLYMERIA, R. Brown, prodr. 488 (1810).									
P. marginata, Bentham, Fl. Austr. IV, 432 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,432	
P. longifolia, Lindley in Mitch. Trop. Austr. 398 (1848)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.IV,432	M.fr.X,114.
P. angusta, F. v. M., fragm. VI, 100 (1868)	—	—	—	—	—	—	N.A.	B.fl.IV,432	M.fr.VI,100.
P. distigma, Bentham, Fl. Austr. IV, 433 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,433	
P. occidentalis, F. v. M., inedit.	W.A.	—	—	—	—	—	—	B.fl.IV,433	
P. calycina, R. Brown, prodr. 488 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,433	M.fr.VI,100.
P. ambigua, R. Brown, prodr. 488 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,433	
PORANA, Burmann, Fl. Ind. 51, t. 21 (1768). (Duperreya.)									
P. sericea, F. v. M., fragm. VI, 100 (1868)	W.A.	—	—	—	—	—	N.A.	B.fl.IV,434	M.fr.II,22.
BREWERIA, R. Brown, prodr. 487 (1810).									
B. linearis, R. Brown, prodr. 488 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,435	
B. media, R. Brown, prodr. 488 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,436	M.fr.X,113.
B. brevifolia, Bentham, Fl. Austr. IV, 436 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,436	
B. rosea, F. v M., fragm. I, 233 (1858)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,436	M.fr.VI,100; X,113.
B. pannosa, R. Brown, prodr. 488 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,436	M.fr.VI,100; X,113.
EVOLVULUS, Linné, spec. plant. ed. sec. 391 (1763).									
E. linifolius, Linné, spec. plant. ed. sec. 392 (1762)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,437	M.fr.VI,100; X,113.
DICHONDRA, R. & G. Forster, char. gen. 39, t. 20 (1776).									
D. repens, R. & G. Forster, char. gen. 39, t. 20 (1776)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,438	M.fr.VI,100; X,113.
CRESSA, Linné, Amoen. Acad. I, 121 (1747).									
C. Cretica, Linné, spec. plant. 223 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,437	M.fr.VI,100.
WILSONIA, R. Brown, prodr. 490 (1810).									
W. humilis, R. Brown, prodr. 490 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,439	M.fr.VI,101; X,113.
W. rotundifolia, Hooker, icon. plant. t. 410 (1842)	W.A.	S.A.	—	V.	N.S.W.	—	-	B.fl.IV,440	M.fr.VI,101.
W. Backhousi, J. Hooker in Lond. Journ. VI, 275 (1842)	W.A.	S.A.	T.	V.	N.S.W.	—	-	B.fl.IV,440	M.fr.VI,101; VIII,17.
CUSCUTA, Tournefort, inst. app. 652, t. 422 (1700), from C. Bauhin (1623).									
C. Chinensis, Lamarck, Encycl. méth. II, 229 (1786)	—	—	—	—	—	Q.	—	B.fl.IV,441	M.fr.X,113.
C. Australis, R. Brown, prodr. 491 (1810)	—	S.A.	—	V.	N.S.W.	Q	-	B.fl IV,441	M.fr.I,100; X,113.
C. Tasmanica, Engelmann in Transact. Acad. St. Louis I, 512 (1860)	—	—	T.	V.	—	-	—	B fl.IV,441	

SOLANACEAE.

Haller, enum. stirp. Helv. praef. 34 (1742).

PHYSALIS, Linné, syst. nat. 8 (1735); Linné, gen. plant. 51 (1737).									
P. minima, Linné, spec. plant. 183 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,466	M.fr.VI,144.
SOLANUM, Tournefort, inst. 148, t. 62 (1700), from Celsus.									
S. nigrum, Linné, spec. plant. 186 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,446	M.fr.VI,145.
S. vescum, F. v. M. in Transact. Vict. Inst. 69 (1855)	—	—	T.	V.	N.S.W.	—	—	—	

S. aviculare, G. Forster, prodr. 18 (1786)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,447	M.fr.VI,144.
S. simile, F. v. M. in Transact. phil. Soc. Vict. I, 19 (1854) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,448	M.fr.I,123;VI,144.
S. fasciculatum, F. v. M., fragm. I, 123 (1859)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VI,144.
S. Bauerianum, Endlicher, prodr. fl. Norfolk, 54 (1833)... ...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,74.
S. viride, R. Brown, prodr. 445 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,449	M.fr.VI,145.
S. tetrandrum, R. Brown, prodr. 445 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,449	
S. Shanesii, F. v. M., fragm. VI, 144 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,448	M.fr.VI,144.
S. verbascifolium, Linné, spec. plant. 184 (1753)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,449	M.fr.VI,145.
S. discolor, R. Brown, prodr. 445 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,450	M.fr.II,166;VI,147.
S. stelligerum, Smith, Exot. Bot. II, 57, t. 88 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,450	M.fr.VI,147.
S. parvifolium, R. Brown, prodr. 446 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,451	M.fr.II,164;VI,147.
S. ferocissimum, Lindley in Mitch. Three Exped. II, 58 (1838)...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.IV,451	
S. sporadotrichum, F. v. M. in Melb. Chemist, Oct. (1882) ...	—	—	—	—	—	Q.	—	—	
S. defensum, F. v. M., fragm. V, 193 (1866)	—	—	—	—	—	Q.	—	B.fl.IV,451	M.fr.V,193.
S. violaceum, R. Brown, prodr. 445 (1810)...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,452	
S. tetrathecum, F. v. M., fragm. II, 165 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,453	M.fr.II,165.
S. elachophyllum, F. v. M., fragm. II, 164 (1861)...	—	—	—	—	—	Q.	—	B.fl.IV,453	M.fr.II,164.
S. orbiculatum, Dunal in Poir. Encycl. méth. XIII, 762 (1817)...	W.A.	S.A.	—	—	—	—	—	B.fl.IV,453	M.fr.VI,145.
S. oligacanthum, F. v. M. in Transact. phil. Soc. Vict. I, 19 (1854)	—	S.A.	—	—	N.S.W.	—	—	B.fl.IV,454	M.fr.VI,145.
S. esuriale, Lindley in Mitch. Three Exped. II, 43 (1838) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,454	
S. chenopodinum, F. v. M., fragm. II, 165 (1861)...	—	S.A.	—	—	N.S.W.	—	N.A.	B.fl.IV,454	M.fr.VI,146.
S. Sturtianum, F. v. M. in Transact. phil. Soc. Vict. I, 19 (1854)	—	S.A.	—	—	N.S.W.	—	N.A.	B.fl.IV,454	M.fr.VI,146.
S. furfuraceum, R. Brown, prodr. 446 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,455	
S. dianthophorum, Dunal, hist. des Solan. 183 (1813)	—	—	—	—	—	Q.	—	B.fl.IV,455	M.fr.VI,145.
S. Dallachyi, Bentham, Fl. Austr. IV, 456 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,456	M.fr.VI,145.
S. densivestitum, F. v. M. in Benth. Fl. Austr. IV, 456 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,456	
S. nemophilum, F. v. M., fragm. II, 161 (1861)	—	—	—	—	—	Q.	—	B.fl.IV,456	M.fr.II,161.
S. Oldfieldii, F. v. M., fragm. II, 161 (1861)	W.A.	—	—	—	—	—	—	B.fl.IV,457	M.fr.VI,145.
S. semiarmatum, F. v. M., fragm. II, 163 (1861)...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,457	M.fr.VI,146.
S. armatum, R. Brown, prodr. 446 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,458	
S. Hystrix, R. Brown, prodr. 446 (1810)	—	S.A.	—	—	—	—	—	B.fl.IV,458	
S. cataphractum, Cunningham in Benth. Fl. Austr. IV, 459 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,459	
S. pungetium, R. Brown, prodr. 446 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,459	M.fr.VI,146.
S. eremophilum, F. v. M. in Schlecht. Linnaea XXV, 432 (1852)	—	S.A.	—	—	N.S.W.	—	—	B.fl.IV,459	M.fr.VI,146.
S. campanulatum, R. Brown, prodr. 446 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,460	M.fr.VI,146.
S. adenophorum, F. v. M., fragm. II, 162 (1861)...	—	—	—	—	—	Q.	—	B.fl.IV,460	M.fr.II,162.
S. cinereum, R. Brown, prodr. 446 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,460	
S. lacunarium, F. v. M. in Transact. phil. Soc. Vict. I, 18 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,461	M.fr.VI,146.
S. petrophilum, F. v. M. in Linnaea XXV, 433 (1852)	—	S.A.	—	—	N.S.W.	—	—	B.fl.IV,461	M.fr.VI,146.
S. diversiflorum, F. v. M., fragm. VI, 146 (1868)...	—	—	—	—	—	—	N.A.	B.fl.IV,461	M.fr.VI,146.
S. carduiforme, F. v. M., fragm. II, 163 (1861)	—	—	—	—	—	—	N.A.	B.fl.IV,462	M.fr.II,163.
S. melanospermum, F. v. M., fragm. II, 163 (1861)	—	—	—	—	—	—	N.A.	B.fl.IV,462	M.fr.II,163.
S. horridum, Dunal, Solan. synops. 28 (1816)	—	—	—	—	—	—	N.A.	B.fl.IV,463	
S. echinatum, R. Brown, prodr. 447 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,463	M.fr.VI,145.
S. lasiophyllum, Dunal in Poir. Encycl. méth. XIII, 764 (1817)	W.A.	—	—	—	—	—	N.A.	B.fl.IV,463	M.fr.VI,145.
S. ellipticum, R. Brown, prodr. 446 (1810)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,464	M.fr.VI,145.
S. quadriloculatum, F. v. M., fragm. II, 161 (1861)	—	—	—	—	—	—	N.A.	B.fl.IV,464	M.fr.II,161.
S. phlomoides, Cunningham in Benth. Fl. Austr. IV, 464 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,464	
S. Cunninghamii, Bentham, Fl. Austr. IV, 465 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,465	
LYCIUM, Linné, gen. plant. 57 (1737), from Celsus.									
L. australe, F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854)...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,467	M.fr.I,83,243;II,179.
DATURA, Linné, gen. plant. 48 (1737).									
D. Leichhardtii, F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854)	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,468	M.fr.VI,144.
NICOTIANA, Tournefort, inst. 117, t. 41 (1700), from C. Bauhin (1623).									
N. suaveolens, Lehmann, gener. Nicot. hist. 43 (1818)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,469	M.fr.VI,144.
ANTHOTROCHE, Endlicher, nov. stirp. mus. Vindob. dec. 6 (1839).									
A. pannosa, Endlicher, nov. stirp. mus. Vind. dec. 7 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,467	M.fr.VI,142.
A. Blackii, F. v. M., fragm. VIII, 232 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,232.
A. Walcottii, F. v. M., fragm. I, 123 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,468	M.fr.I,123.
ANTHOCERCIS, Labillardière, Nov. Holl. pl. spec. II, 19 (1806). (Cyphanthera, Eadesia.)									
A. viscosa, R. Brown, prodr. 448 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,475	M.fr.VI,143.
A. litorea, Labillard., Nov. Holl. pl. spec. II, 19, t. 158 (1806)	W.A.	—	—	—	—	—	—	B.fl.IV,476	M.fr.VI,143;XI,123.
A. gracilis, Bentham in De Cand., prodr. X, 192 (1846)... ...	W.A.	—	—	—	—	—	—	B.fl.IV,476	
A. genistoides, Miers, Illustr. of plants II, App. 27, t. 83 (1857)	W.A.	—	—	—	—	—	—	B.fl.IV,476	M.fr.I,122;VI,143.
A. anisantha, Endlicher in Ann. Wien. Mus. II, 201 (1838) ...	W.A.	S.A.	—	—	—	—	—	B.fl.IV,477	M.fr.VI,143.
A. intricata, F. v. M., fragm. I, 211 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,477	M.fr.I,211.
A. arborea, F. v. M., fragm. I, 212 (1859)...	W.A.	—	—	—	—	—	—	B.fl.IV,477	M.fr.I,212;VI,143.
A. angustifolia, F. v. M. in Transact. phil. Soc. Vict. I, 21 (1854)	—	S.A.	—	—	—	—	—	B.fl.IV,478	
A. fasciculata, F. v. M., fragm. I, 122 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,478	M.fr.I,122;VI,143.
A. microphylla, F. v. M., fragm. I, 179 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,478	M.fr.I,179;VI,143.
A. myosotidea, F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,478	M.fr.I,179;VI,143.
A. scabrella, Bentham in De Cand., prodr. X, 192 (1846) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,479	
A. albicans, Cunningham in Field's N.S.W. 335, t. 2 (1825) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,479	M.fr.VI,143.
A. Odgersii, F. v. M., fragm. X, 19 (1876)...	—	—	—	—	—	—	N.A.	—	M.fr.X,19.
A. Tasmanica, J. Hooker, Fl. Tasman. I, 289, t. 92 (1860) ...	—	—	T.	—	—	—	—	B.fl.IV,479	M.fr.VI,143.
A. Eadesii, F. v. M., fragm. II, 139 (1861)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,480	M.fr.VI,142.
A. racemosa, F. v. M., fragm. I, 211 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,480	M.fr.I,211.

DUBOISIA, R. Brown, prodr. fl. Nov. Holl. 448 (1810).

D. myoporoides, R. Brown, prodr. 448 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,474	M.fr.VI,144.
D. Hopwoodi, F. v. M., fragm. VIII, 232 (1874)	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.IV,480	M.fr.II,138;VI,143;X,20.
D. Leichhardti, F. v. M. in Wing's S. Sc. Record, II, 222 (1882)	—	—	—	—	N.S.W.	—	—	B.fl.IV,480	M.fr.VI,142.

SCROPHULARINAE.

Mirbel, Elém. de Botan. II, 879 (1815).

MIMULUS, Linné in Act. Upsal. 82 (1741). (Uvedalia.)

M. Uvedaliae, Bentham in De Cand. X, 369 (1846)	—	—	—	—	—	Q.	N.A.	B.fl.IV,482	M.fr.VI,103;IX,167.
M. debilis, F. v. M. in Transact. phil. Inst. Vict. I, 62 (1857) ...	—	—	—	—	—	—	N.A.	—	M.fr.VI,103.
M. gracilis, R. Brown, prodr. 439 (1810)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,482	M.fr.VI,103;IX,167.
M. repens, R. Brown, prodr. 439 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,482	M.fr.VI,103;IX,166.
M. prostratus, Bentham in De Cand. X, 373 (1846)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,483	M.fr.IX,167.

MAZUS, Loureiro, Fl. Cochinch. II, 385 (1790).

M. Pumilio, R. Brown, prodr. 439 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,484	M.fr.VI,104;IX,167.

ADENOSMA, R. Brown, prodr. fl. Nov. Holl. 442 (1810). (Pterostigma.)

A. coerulea, R. Brown, prodr. 443 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,484	
A. Muelleri, Bentham, Fl. Austr. IV, 485 (1869)...	—	—	—	—	—	—	N.A.	B.fl.IV,485	

STEMODIA, Linné, syst. ed. dec. 1118 (1759). (Morgania, Limnophila.)

S. lythrifolia, F. v. M. in Benth. Fl. Austr. IV, 486 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,486	
S. grossa, Bentham, Fl. Austr. IV, 486 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,486	
S. viscosa, Roxburgh, pl. of Coromand. II, 33, t. 163 (1798) ...	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,486	
S. linophylla, F. v. M., fragm. X, 88 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,88.
S. debilis, Bentham, Fl. Austr. IV, 487 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,487	
S. pedicellaris, F. v. M., fragm. VIII, 231 (1874)...	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,231.
S. Morgania, F. v. M., fragm. X, 89 (1876)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,488	M.fr.VI,104.
S. sessiliflora, F. v. M., Limnophila, Blume, Bijdr. 750 (1826) ...	—	—	—	—	—	—	N.A.	—	
S. gratioloides, F. v. M., fragm. X, 89 (1876)	—	—	—	—	—	Q.	N.A.	B.fl.IV,489	M.fr.X,89.
S. punctata, F. v. M.; Limnophila, Blume, Bijdr. 750 (1826) ...	—	—	—	—	—	Q.	—	B.fl.IV,490	
S. hirsuta, Heyne in Wallich, numer. list. 3930 (1828)	—	—	—	—	—	—	N.A.	B.fl.IV,490	
S. tenuiflora, Bentham, Scrophul. Indic. 23 (1835)	—	—	—	—	—	—	N.A.	B.fl.IV,490	

BRAMIA, Lamarck, Encycl. méthod. I, 459 (1783). (Herpestis.)

B. floribunda, F. v. M., fragm. IX, 167 (1875)	—	—	—	—	—	Q.	N.A.	B.fl.IV,491	M.fr.VI,104.
B. Indica, Lamarck, Encycl. méth. I, 459 (1783)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,491	M.fr.IX,167.

GRATIOLA, Ruppius, Fl. Jenens. 241 (1718), from Dodoens (1556).

G. pedunculata, R. Brown, prodr. 435 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,492	M.fr.VI,108;IX,167.
G. Peruviana, Linné, spec. plant. 17 (1753)..	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,493	M.fr.VI,103;IX,167.
G. nana, Bentham in De Cand. prodr. X, 404 (1846)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,493	M.fr.IX,167.

DOPATRIUM, Hamilton in Benth. Scroph. Ind. 4 (1835).

D. junceum, Hamilton in Benth. Scroph. Ind. 4 (1745)	—	—	—	—	—	Q.	—	B.fl.IV,494	

ARTANEMA, D. Don in Sweet, Fl. gard. t. 234 (1829).

A. fimbriatum, D. Don in Sweet, Fl. Gard. t. 234 (1829) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,495	M.fr.VI,104;IX,167.

LINDERNIA, Allioni, rarior. Pedemont. stirp. specim. t. 5 (1755). (Vandellia, Bonnaya, Ilysanthes, Torenia partly, Tittmannia partly.)

L. crustacea, F. v. M.; Vandellia, Bentham, Scroph. Indic. 35 (1835)	—	—	—	—	N.S.W.	Q.	N.A.	—	
L. pubescens, F.v.M.; Vandellia, Benth. in De Cand. pr. X, 415(1846)	—	—	—	—	—	—	N.A.	B.fl.IV,496	M.fr.VI,102.
L. alsinoides, R. Brown, prodr. 441 (1810)...	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,496	
L. scapigera, R. Brown, prodr. 441 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,497	M.fr.VI,102.
L. subulata, R. Brown, prodr. 441 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,497	M.fr.VI,102.
L. lobelioides; Vandellia, F. v. M. in Transact. phil. Inst. Vict. III, 61 (1858)	—	—	—	—	—	—	N.A.	B.fl.IV,497	M.fr.VI,102.
L. clausa, F. v. M., fragm. VI, 102 (1868)	—	—	—	—	—	—	N.A.	B.fl.IV,498	
L. veronicifolia, F. v. M., fragm. VI, 101 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,498	M.fr.VI,101.
L. serrata, F.v.M.; Gratiola, Roxburgh. fl. Ind. ed. Car. I, 140 (1820)	—	—	—	—	—	Q.	—	—	

HEMIARRHENA, Bentham, Fl. Austr. IV, 518 (1869).

H. plantaginea, Bentham in Hook. icon. pl. t. 1059 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,518	M.fr.VI,102.

PEPLIDIUM, Delile, Fl. Aegypt. 148 (1813).

P. humifusum, Delile, Fl. Aegypt. 148 (1813)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,500	M.fr.VI,105.
P. Muelleri, Bentham, Fl. Austr. IV, 500 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,500	

MICROCARPAEA, R. Brown, prodr. 435 (1810).

M. muscosa, R. Brown, prodr. 435 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,501	M.fr.VI,104.

GLOSSOSTIGMA, Arnott in nov. act. acad. Caes. Leop. Carol. XVIII, 355 (1836). (Tricholoma.)

G. spathulatum, Arnott in nov. act. Leop. XVIII, 355 (1836) ...	—	—	—	—	—	Q.	—	B.fl.IV,501	
G. Drummondii, Bentham in De Cand. prodr. X, 426 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.IV,502	M.fr.VI,104.
G. elatinoides, Bentham in J. Hook. Fl. New Zeal. I, 189 (1853)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,502	

LIMOSELLA, Lindern, opusc. plant. Argentorat. 156, t. 5 (1728).

L. aquatica, Linné, spec. plant. 631 (1753)...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,502	M.fr.VI,105;IX,165.
L. Curdieana, F. v. M., fragm. IX, 166 (1875)	—	S.A.	—	V.	N.S.W.	—	—	—	M.fr.IX,166.

SCOPARIA, Linné, syst. nat. ed. sext. 87 (1748).

S. dulcis, Linné, spec. plant. 116 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.IV,504	M.fr.VI,104.

OURISIA, Commerçon in A. L. de Jussieu, gen. pl. 100 (1789).

O. integrifolia, R. Brown, prodr. 439 (1810)	—	—	T.	—	—	—	—	B.fl.IV,512	M.fr.VI,105;IX,168.

VERONICA, Tournefort, inst. 143, t. 60 (1700), from Fuchs (1543). (Pygmaea.)

V. densifolia, F. v. M., fragm. II, 137, t. 63 (1861)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,505	M.fr.VI,103.
V. formosa, R. Brown, prodr. 434 (1810)	—	—	T.	—	—	—	—	B.fl.IV,506	M.fr.VI,102;IX,167.

V. decorosa, F. v. M. in Schlechtend., Linnaea XXV, 430 (1852)	—	S.A.	—	—	—	—	—	B.fl.IV,506	
V. perfoliata, R. Brown, prodr. 434 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,507	M.fr.VI,102;IX,167.
V. Derwentia, Littlejohn in Andr. Bot. Reposit. t. 531 (1808) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,507	M.fr.VI,102;IX,168.
V. arenaria, Cunningham in De Cand. prodr. X, 463 (1846) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,507	M.fr.VI,102;IX,167.
V. nivea, Lindley, Bot. Regist. Misc. 42 (1842)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,508	M.fr.VI,102;IX,167.
V. gracilis, R. Brown, prodr. 435 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,508	M.fr.VI,103;IX,168.
V. distans, R. Brown, prodr. 435 (1810)	W.A.	S.A.	T.	V.	—	—	—	B.fl.IV,509	M.fr.VI,102.
V. calycina, R. Brown, prodr. 435 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,509	M.fr.VI,103.
V. plebeja, R. Brown, prodr. 435 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,510	M.fr.VI,102.
V. notabilis, F. v. M. in Benth. Fl. Austr. IV, 511 (1869) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,511	M.fr.IX,168.
V. serpillifolia, Linné, spec. plant. 12 (1753)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,511	M.fr.VI,103.
V. peregrina, Linné, spec. plant. 14 (1753)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,511	M.fr.VI,103;IX,168.
RAMPHICARPA, Bentham in Hook. Comp. to the Bot. Mag. I, 368 (1835).									
R. longiflora, Bentham in Hook. Comp. Bot. Mag. I, 368 (1835)	—	—	—	—	—	Q.	N.A.	B.fl.IV,518	M.fr.VI,104;IX,168.
CENTRANTHERA, R. Brown, prodr. fl. Nov. Holland. 438 (1810).									
C. hispida, R. Brown, prodr. 438 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,513	M.fr.VI,104;IX,168.
SOPUBIA, Hamilton in D. Don, prodrom. flor. Nepalens. 88 (1825).									
S. trifida, Hamilton in D. Don, prodr. fl. Nep. 88 (1825) ...	—	—	—	—	—	Q.	—	B.fl.IV,513	
BUECHNERA, Linné, hort. Cliffort. 501 (1737). (Buchnera, Striga.)									
B. tetragona, R. Brown, prodr. 437 (1810)...	—	—	—	—	—	Q.	N.A.	B.fl.IV,514	M.fr.IX,168.
B. urticifolia, R. Brown, prodr. 437 (1810)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,514	M.fr.VI,104.
B. linearis, R. Brown, prodr. 437 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,515	
B. tenella, R. Brown, prodr. 437 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,515	
B. gracilis, R. Brown, prodr. 437 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,515	M.fr.VI,104.
B. ramosissima, R. Brown, prodr. 438 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,515	
B. coccinea, Bentham, Scrophul. Ind. 40 (1835)	—	—	—	—	—	Q.	—	B.fl.IV,516	
B. parviflora, R. Brown, prodr. 438 (1810)...	—	—	—	—	—	Q.	—	B.fl.IV,517	M.fr.VI,104.
B. multiflora, F.v.M.; Striga, Benth. in De Cand., prodr. X, 501 (1836)	—	—	—	—	—	—	N.A.	B.fl.IV,517	
B. curviflora, R. Brown, prodr. 438 (1810)...	—	—	—	—	—	Q.	N.A.	B.fl.IV,517	M.fr.VI,104.
EUPHRASIA, Tournefort, instit. 174, t. 78 (1700), from Matthaeus (1478).									
E. Brownii, F. v. M., fragm. V, 88 (1865)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,521	M.fr.IX,168.
E. scabra, R. Brown, prodr. 437 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,521	M.fr.IX,168.
E. cuspidata, J. Hooker, Fl. Tasman. I, 298 (1860) ...	—	—	T.	—	—	—	—	B.fl.IV,522	
E. antarctica, Bentham in De Cand., prodr. X, 555 (1846) ...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,522	

OROBANCHEAE.

A. L. de Jussieu in Ann. Mus. V, 25 (1804).

OROBANCHE, Tournefort, inst. 175, t. 81 (1700), from l'Ecluse (1583).									
O. Australiana, F. v. M., inedit.	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,533	

LENTIBULARINAE.

L. C. Richard in Bulliard, Fl. Paris. ed. sec. I, 26 (1796).

UTRICULARIA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 5 (1737).									
U. stellaris, Linné, fil. suppl. plant. 86 (1781)	—	—	—	—	—	Q.	N.A.	B.fl.IV,525	M.fr.VI,161.
U. flexuosa, Vahl, enum. plant. I, 198 (1804)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,525	
U. exoleta, R. Brown, prodr. 430 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,526	M.fr.VI,161.
U. tubulata, F. v. M., fragm. IX, 48 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,48.
U. albiflora, R. Brown, prodr. 431 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,526	
U. pygmaea, R. Brown, prodr. 432 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,526	
U. fulva, F. v. M. in Transact. phil. Inst. Vict. III, 63 (1858)...	—	—	—	—	—	—	N.A.	B.fl.IV,527	M.fr.VI,160.
U. chrysantha, R. Brown, prodr. 432 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,527	M.fr.VI,161.
U. bifida, Linné, spec. plant. 18 (1753)	—	—	—	—	—	Q.	—	B.fl.IV,527	
U. cyanea, R. Brown, prodr. 431 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,527	M.fr.VI,161.
U. lateriflora, R. Brown, prodr. 431 (1810)...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,528	M.fr.VI,162.
U. simplex, R. Brown, prodr. 431 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,528	M.fr.VI,162.
U. monantha, J. Hooker, Fl. Tasman. I, 299 (1860)	—	—	T.	—	—	—	—	B.fl.IV,528	
U. dichotoma, Labillard., Nov. Holl. pl. spec. I, 11, t. 8 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,529	M.fr.VI,161.
U. volubilis, R. Brown, prodr. 430 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,529	M.fr.VI,160.
U. Hookeri, Lehmann, nov. stirp. pugill. VIII, 47 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.IV,530	M.fr.VI,162.
U. violacea, R. Brown, prodr. 431 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,530	M.fr.VI,160.
U. Menziesii, R. Brown, prodr. 431 (1810)...	W.A.	—	—	—	—	—	—	B.fl.IV,530	M.fr.VI,160.
U. biloba, R. Brown, prodr. 432 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV.531	
U. limosa, R. Brown, prodr. 432 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,531	
U. Baueri, R. Brown, prodr. 431 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,531	
POLYPOMPHOLYX, Lehmann in der Bot. Zeitung, Halle, 109 (1844). (Tetralobus.)									
P. multifida, F. v. M., fragm. VI, 162 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,532	M.fr.VI,162.
P. tenella, Lehmann, nov. stirp. pugill. VIII, 50 (1844)... ...	W.A.	S.A.	T.	V.	—	—	—	B.fl.IV,532	M.fr.VI,162.

GESNERIACEAE.

Humboldt, Bonpland & Kunth, nov. gen. et spec. pl. II, 392 (1817).

FIELDIA, Cunningham in Field's New South Wales 363, t. 2 (1825).									
F. australis, Cunningham in Field's N.S.W. 364, t. 2 (1825) ...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,535	M.fr.IX,191.
NEGRIA, F. v. M., fragm. VII, 151 (1871). (Rhabdothamnus partly.)									
N. rhabdothamnoides, F. v. M., fragm. VII, 152 (1871)... ...	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,152;XI,133.
BAEA, Commerçon in Lamarck, Encycl. méth. I, 401 (1783). (Streptocarpus.)									
B. hygroscopica, F. v. M., fragm. IV, 146 (1864)...	—	—	—	—	—	Q.	—	B.fl.IV,535	M.fr.IV,146.

BIGNONIACEAE.

Ventenat, Tabl. du règn. végét. II, 402 (1799).

TECOMA, A. L. de Jussieu, gen. plant. 139 (1789).

T. australis, R. Brown, prodr. 471 (1810)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,537	
T. Hillii, F. v. M., fragm. X, 101 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,101;XI,136.
T. jasminoides, Lindley, Bot. Regist. t. 2002 (1838)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,537	

DOLICHANDRONE, Fenzl in Ann. nat. hist. third ser. X, 31 (1862). (Spathodea partly, Dolichandra partly.)

D. heterophylla, F. v. M., fragm. IV, 149 (1864)..	—	—	—	—	—	Q.	N.A.	B.fl.IV,539	M.fr.IV,149.
D. filiformis, Seemann, Journ. of Bot. I, 226 (1863)	—	—	—	—	—	—	N.A.	B.fl.IV,539	

HAUSSMANNIA, F. v. M., fragm. IV, 148 (1864).

H. jucunda, F. v. M., fragm. IV, 148 (1864)	—	—	—	—	—	Q.	—	B.fl.IV,540	M.fr.IV,148.

DIPLANTHERA, R. Brown, prodr. Fl. Nov. Holl. 449 (1810). (Deplanchea, Bulweria.)

D. tetraphylla, R. Brown, prodr. 449 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,540	

ACANTHACEAE.

A. L. de Jussieu, gen. plant. 102 (1789), from B. de Jussieu (1759).

THUNBERGIA, Retzius, Physiogr. Saellsk. Handl. I, 163 (1776).

T. Arnhemica, F. v. M., fragm. IX, 73 (1875)	—	—	—	—	—	—	N.A.	—	M.fr.IX,73.
T. Powelli, F. v. M. in Wing's South. Sc. Rec. II, 34 (1882) ...	—	—	—	—	—	—	N.A.	—	

NELSONIA, R. Brown, prodr. fl. Nov. Holl. 485 (1810).

N. campestris, R. Brown, prodr. 481 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,543	M.fr.VI,88;XI,18.

EBERMAYERA, Nees in Wallich, pl. Asiat. rar. III, 75 (1832).

E. glauca, Nees in De Candolle, prodr. XI, 73 (1847)	—	—	—	—	—	Q.	N.A.	B.fl.IV,544	M.fr.VI,88;XI,18

HYGROPHILA, R. Brown, prodr. Fl. Nov. Holl. 479 (1810).

H. angustifolia, R. Brown, prodr. 479 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,544	M.fr.VI,88.

STROBILANTHES, Blume, Bijdr. tot de fl. van Nederl. Ind. 796 (1826).

S. Tatei, F. v. M. in transact. Roy. Soc. S. Austr. V, Aug. (1882)	—	—	—	—	—	—	N.A.	—	

RUELLIA, Plumier, nov. gen. 12, t. 2 (1703). (Dipteracanthus.)

R. bracteata, R. Brown, prodr. 479 (1810)...	—	—	—	—	—	Q.	N.A.	B.fl.IV,546	M.fr.VI,91.
R. primulacea, F. v. M. in Benth. Fl. Austr. IV, 546 (1869) ...	—	S.A.	—	—	—	Q.	—	B.fl.IV,546	M.fr.XI,18.
R. corynotheca, F. v. M. in Benth. Fl. Austr. IV, 546 (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,546	M.fr.XI,18.
R. australis, R. Brown, prodr. 479 (1810)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.IV,547	M.fr.VI,91.
R. spiciflora, F. v. M. in Benth. Fl. Austr. IV, 547 (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,547	
R. acaulis, R. Brown, prodr. 479 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,547	

ACANTHUS, Tournefort, inst. 176, t. 81 (1700), from Theophrastos. (Dilivaria.)

A. ilicifolius, Linné, spec. plant. 939 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.IV,548	M.fr.X,18.

JUSTICIA, Houston in Linné, gen. plant. 4 (1737). (Rostellaria, Rostellularia.)

J. procumbens, Linné, spec. plant. 15 (1753)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,549	M.fr.VI,91;XI,18.
J. Bonneyana, F. v. M. in Wing's S. Sc. Record II, 74 (1882) ...	—	S.A.	—	—	N.S.W.	—	—	—	M.fr.X,73.
J. hygrophiloides, F. v. M., fragm. VI, 89 (1868)...	—	—	—	—	—	Q.	—	B.fl.IV,550	M.fr.VI,89.
J. Kempeana, F. v. M., fragm. XI, 101 (1881)	W.A.	S.A.	—	—	—	—	N.A.	—	M.fr.XI,101.
J. cavernarum, F. v. M., fragm. VI, 91 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,550	M.fr.VI,91.
J. eranthemoides, F. v. M., fragm. VI, 90 (1868)...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,551	M.fr.VI,90.

GRAPTOPHYLLUM, Nees in Wall. pl. Asiat. rar. III, 102 (1832). (Earlia, Thyrsacanthus.)

G. Earlii, F. v. M., fragm. VI, 87 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,551	M.fr.VI,87.
G. ilicifolium, F. v. M. in Benth. Fl. Austr. IV, 552 (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,552	
G. spinigerum, F. v. M., fragm. XI, 17 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,17.

DICLIPTERA, A. L. de Jussieu in Ann. du Mus. IX, 267 (1807). (Brochosiphon.)

D. glabra, Decaisne in Nouv. Annal. du Mus. III, 388 (1834) ...	—	—	—	—	—	—	N.A.	B.fl.IV,552	M.fr.VI,88;XI,18.
D. spicata, Decaisne in Nouv. Annal. du Mus. III, 389 (1834)...	—	—	—	—	—	Q.	—	B.fl.IV,553	M.fr.VI,89.
D. Burmanni, Nees in Wallich, pl. Asiat. rar. III, 112 (1832) ...	—	—	—	—	—	Q.	—	—	M.fr.VII,62.

HYPOESTES, Solander in R. Brown, prodr. 474 (1810).

H. floribunda, R. Brown, prodr. 474 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,554	M.fr.XI,18.

ERANTHEMUM, Linné, Fl. Zeyl. 6 (1747).

E. variabile, R. Brown, prodr. 477 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,555	M.fr.VI,91;XI,18.
E. tenellum, Bentham, Fl. Austr. IV, 555 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,555	

HYDROPHYLLEAE.

R. Brown in Bot. Regist. t. 242 (1817).

HYDROLEA, Linné, spec. plant. ed. sec. 328 (1762).

H. Zeilanica, Vahl, symbol. botan. II, 46 (1791)	—	—	—	—	—	—	N.A.	B.fl.IV,382	M.fr.V,192.
H. spinosa, Linné, spec. plant. ed. sec. 328 (1762)	—	—	—	—	—	Q.	—	B.fl.IV,383	M.fr.V,192.

ASPERIFOLIAE.

Haller, enum. stirp. Helv. Praef. 34 (1742).

CORDIA, Plumier, gen. plant. 13, t. 14 (1703).

C. subcordata, Lamarck, Illustr. des genr. I, 421 (1791)... ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,385	M.fr.VI,114;IX,122.[122.
C. aspera, G. Forster, florul. ins. austr. prodr. 18 (1786)... ...	—	—	—	—	—	Q.	—	B.fl.IV,386	M.fr.V,193;VI,114;IX,
C. Myxa, Linné, spec. plant. 190 (1753)	—	—	—	—	—	Q.	—	B.fl.IV,386	M.fr.I,59;VI,114.

EHRETIA, P. Browne, Civ. and Nat. Hist. of Jamaic. 168, t. 16 (1756).

E. acuminata, R. Brown, prodr. 497 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,387	M.fr.V,21;IX,122;XI,124
E. pilosula, F. v. M., fragm. V, 20 (1865)...	—	—	—	—	—	Q.	—	B.fl.IV,388	M.fr.V,20.
E. saligna, R. Brown, prodr. 497 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,388	M.fr.V,21;IX,122.

E. membranifolia, R. Brown, prodr. 497 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,388	M.fr.V,21;VI,115;IX,122
E. laevis, Roxburgh, Pl. Corom. I, 42, t. 56 (1795)	—	—	—	—	—	Q.	—	B.fl.IV,389	
TOURNEFORTIA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 55 (1737.)									
T. argentea, Linné fil. suppl. plant. 133 (1781)	—	—	—	—	—	Q.	N.A.	B.fl.IV,389	M.fr.IV,95;VI,116.
T. mollis, F. v. M., fragm. I, 59 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.IV,390	M.fr.VI,116.
T. sarmentosa, Lamarck, Illustr. des genr. I, 416 (1791) ...	—	—	—	—	—	Q.	—	B.fl.IV,390	M.fr.IV,95;VI,116.
COLDENIA, Linné, amoen. acad. ed. prim. 119 (1747). (Lobophyllum.)									
C. procumbens, Linné, spec. plant. 125 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.IV,391	M.fr.VI,115.
HELIOTROPIUM, Tournefort, inst. 138, t. 57 (1700), from Theophr., Diosc. & Plinius. (Tiaridium, Heliophytum.)									
H. Indicum, Linné, spec. plant. 130 (1753)	—	—	—	—	—	—	N.A.	—	M.fr.XI,124.
H. pleiopterum, F. v. M., fragm. IX, 121 (1875)...	—	S.A.	—	—	—	—	—	—	M.fr.IX,121.
H. curassavicum, Linné, spec. plant. 130 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,393	M.fr.VI,116;IX,121.
H. Europaeum, Linné, spec. plant. 130 (1753)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,394	M.fr.VI,116. [124.
H. undulatum, Vahl, symbol. bot. I, 13 (1795)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,394	M.fr.VI,116;IX,121;XI,
H. asperrimum, R. Brown, prodr. 493 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,394	M.fr.VI,116;IX,121.
H. crispatum, F. v. M. in Benth. Fl. Austr. IV, 395 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,395	
H. fasciculatum, R. Brown, prodr. 494 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,395	
H. vestitum, Bentham, Fl. Austr. IV, 395 (1869)	—	—	—	—	N.S.W.	—	—	B.fl.IV,395	
H. brachygyne, Bentham, Fl. Austr. IV, 396 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,396	
H. epacrideum, F. v. M. in Benth. Fl. Austr. IV, 396 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,396	
H. ovalifolium, Forskael, Fl. Aegypt. Arab. 47 (1775)	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,396	M.fr.VI,115.
H. strigosum, Willdenow, sp. pl. I, 743 (1797)	—	—	—	—	—	Q.	N.A.	B.fl.IV,397	
H. prostratum, R. Brown, prodr. 494 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,397	
H. bracteatum, R. Brown, prodr. 493 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,397	
H. pauciflorum, R. Brown, prodr. 493 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,398	
H. conocarpum, F. v. M. in Benth. Fl. Austr. 398 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,398	
H. filaginoides, Bentham, Fl. Austr. IV, 398 (1869)	—	S.A.	—	—	—	—	—	B.fl.IV,398	
H. ventricosum, R. Brown, prodr. 494 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,399	
H. tenuifolium, R. Brown, prodr. 494 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,399	
H. paniculatum, R. Brown, prodr. 494 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,399	
H. Cunninghamii, Bentham, Fl. Austr. IV, 400 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,400	
H. diversifolium, F. v. M. in Benth. Fl. Austr. IV, 400 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,400	
HALGANIA, Gaudichaud in Freyc. in voy. Botan. 448 (1826).									
H. solanacea, F. v. M. in Hook. Kew Misc. IX, 21 (1857) ...	—	—	—	—	—	—	N.A.	B.fl.IV,401	M.fr.VI,115.
H. litoralis, Gaudichaud in Freyc. voy. Bot. 449, t. 59 (1826) ...	W.A.	—	—	—	—	—	—	B.fl.IV,401	M.fr.VI,115.
H. corymbosa, Lindley, Bot. Regist. XXV, App. XL (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,402	
H. sericiflora, Bentham, Fl. Austr. IV, 402 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,402	
H. cyanea, Lindley, Bot. Regist. XXV, App. XL (1839) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	—	M.fr.VI,115;IX,122.
H. lavandulacea, Endlicher in Ann. Wien. Mus. II, 205 (1838)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,402	M.fr.I,209;VI,115;XI,125
H. integerrima, Endlicher in Ann. Wien. Mus. II, 205 (1838) ...	W.A.	—	—	—	—	—	—	B.fl.IV,402	
POLLICHIA, Medikus, Botan. Beobach. 247 (1783). (Trichodesma.)									
P. Zeylanica, F. v. M.; Trichodesma, R. Brown, prodr. 496 (1810)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,404	M.fr.XI,125.
P. latisepalea; Trichodesma, F. v. M., fragm. X, 102 (1877) ...	—	—	—	—	—	—	—	—	M.fr.X,121.
MYOSOTIS, Ruppius, Fl. Jenens. 9 (1718), from Dioscorides and Plinius. (Exarrhena.)									
M. australis, R. Brown, prodr. 495 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,405	M.fr.VI,116;IX,122.
M. suaveolens, Poiret, Encycl. méth. suppl. IV, 44 (1816) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,406	M.fr.VI,115;IX,122.
ERITRICHUM, Schrader in commentat. Goetting. IV, 186 (1820). (Eritrichium.)									
E. Australasicum, A. De Candolle, prodr. X, 134 (1846) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,406	M.fr.XI,125.
LAPPULA, Rivinus in Rupp. Fl. Jenens. (1718), from Dalechamp. (Echinospermum.)									
L. concava; Echinospermum, F. v. M., fragm. II, 139 (1861) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,407	M.fr.VI,116;IX,121.
ROCHELIA, Reichenbach in Fl. Regensb. bot. Zeit. 1, 243 (1824).									
R. Maccoya, F. v. M. in Benth. Fl. Austr. IV, 408 (1869) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,408	M.fr.I,127;IX,121.
CYNOGLOSSUM, Tournefort, inst. 139, t. 57 (1700), from Dioscorides and Nicandros. [125.									
C. latifolium, R. Brown, prodr. 495 (1810)...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,408	M.fr.VI,115;IX,122;XI,
C. suaveolens, R. Brown, prodr. 495 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,409	M.fr.VI,115;IX,122.
C. australe, R. Brown, prodr. 495 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,409	M.fr.VI,115;IX,122.
C. Drummondii, Bentham, Fl. Austr. IV, 409 (1869)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,409	M.fr.IX,122.

LABIATAE.

Adanson, Famill. des plant. II, 180 (1763), from B. de Jussieu (1759).

OCIMUM, Tournefort, inst. 203, t. 96 (1700), from Theophrastos and Dioscorides. (Ocymum.)									
O. sanctum, Linné, mantiss. plant. 85 (1767)	—	—	—	—	—	Q.	N.A.	B.fl.V,74	M.fr.IV,46.
MOSCHOSMA, Reichenbach, conspect. regn. vegetab. 115 (1828). (Basilicum 1802.)									
M. polystachya, Bentham in Wall. pl. As. rar. II, 13 (1831) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,75	M.fr.VI,108.
ORTHOSIPHON, Bentham, Labiatar. gen. et spec. 25 (1832).									
O. stamineus, Bentham in Wall. pl. As. rar. II, 15 (1831) ...	—	—	—	—	—	Q.	N.A.	B.fl.V,76	M.fr.VI,108;IX,161.
PLECTRANTHUS, L'Héritier, stirp. nov. I, 85 (1785).									
P. longicornis, F. v. M., fragm. V, 51 (1865)	—	—	—	—	—	Q.	—	B.fl.V,77	M.fr.V,51.
P. parviflorus, Willdenow, hort. Berol. I, t. 65 (1806)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,78	M.fr.VI,108;IX,161.
P. congestus, R. Brown, prodr. 506 (1810)...	—	—	—	—	—	Q.	—	B.fl.V,78	M.fr.VI,108.
P. foetidus, Bentham, labiat. 35 (1832)	—	—	—	—	—	Q.	—	—	M.fr.IX,161;XI,135.
COLEUS, Loureiro, Fl. Cochinch. II, 372 (1790).									
C. scutellarioides, Bentham in Wall. pl. Asiat. rar. II, 16 (1831)	—	—	—	—	—	Q.	N.A.	B.fl.V,79	M.fr.VI,108.
POGOSTEMON, Desfontaines in Mém. du Mus. II, 154, t. 6 (1815). (Dysophylla.)									
P. verticillatus, Hasskarl, catal. hort. Bogor. 131 (1844) ...	—	—	—	—	—	Q.	N.A.	B.fl.V,81	M.fr.V,200;IX,161.

MENTHA, Tournefort, inst. 188, t. 89 (1700), from Hippocrates and Theophrastos.

M. laxiflora, Bentham in De Cand. XII, 174 (1848)	—	—	—	V.	N.S.W.	—	—	B.fl.V,82	M.fr.VI,109;IX,162.
M. grandiflora, Bentham in Mitch. Trop. Austr. 362 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,83	
M. australis, R. Brown, prodr. 505 (1810)...	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,83	M.fr.VI,109;IX,162.
M. gracilis, R. Brown, prodr. 505 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,83	M.fr.VI,109.
M. saturejoides, R. Brown, prodr. 505 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,84	M.fr.VI,109.

LYCOPUS, Tournefort, inst. 180, t. 83 (1700), from Plinius.

L. australis, R. Brown, prodr. 500 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,85	M.fr.VI,109.

SALVIA, Linné, gen. plant. 6 (1737), from Plinius.

S. plebeja, R. Brown, prodr. 501 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,85	M.fr.VI,109;IX,161.

BRUNELLA, L'Ecluse, rar. stirp. hist. II, 42, 43 (1576), from Brunfels (1536). (Prunella.)

B. vulgaris, A. de Candolle, prodr. XII, 410 (1848)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,87	M.fr.VI,109.

SCUTELLARIA, Hermann, hort. Lugd. bot. cat. (1687), from Cortusi (1591).

S. mollis, R. Brown, prodr. 507 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.V,88	M.fr.VI,109.
S. humilis, R. Brown, prodr. 507 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,88	M.fr.VI,109.

ANISOMELES, R. Brown, prodr. 503 (1810).

A. salvifolia, R. Brown, prodr. 503 (1810)...	—	—	—	—	—	Q.	N.A.	B.fl.V,89	M.fr.VI,109;IX,162.

LEUCAS, J. Burmann, thesaur. Zeyl. 140 (1737).

L. flaccida, R. Brown, prodr. 505 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.V,90	M.fr.VI,109.

DEPREMESNILIA, F. v. M., fragm. X, 59 (1876).

D. chrysocalyx, F. v. M., fragm. X, 59 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,59.

PROSTANTHERA, Labillardière, Nov. Holl. pl. spec. 18, t. 157 (1806). (Chilodia, Cryphia, Klanderia.)

P. lasiantha, Labillard., Nov. Holl. pl. spec. II, 18, t. 15 (1806)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,93	M.fr.VI,105.
P. prunelloides, B. Brown, prodr. 508 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,94	M.fr.VI,106.
P. coerulea, R. Brown, prodr. 508 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,94	M.fr.VI,106.
P. ovalifolia, R. Brown, prodr. 509 (1810)...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,95	M.fr.VI,106.
P. melissifolia, F. v. M., fragm. I, 19 (1858)	—	—	—	V.	—	—	—	B.fl.V,95	M.fr.I,19,242.
P. incisa, R. Brown, prodr. 509 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,95	M.fr.VI,107;IX,162.
P. rotundifolia, R. Brown, prodr. 509 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,96	M.fr.VI,107;IX,162.
P. violacea, R. Brown, prodr. 509 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,96	M.fr.VI,107;IX,162.
P. incana, Cunningham in Benth. lab. 455 (1836)... ...	—	—	—	—	N.S.W.	—	—	B.fl.V,97	M.fr.IX,162.
P. hirtula, F. v. M., first gen. Report 16 (1853)	—	—	—	V.	N.S.W.	—	—	B.fl.V,97	
P. denticulata, R. Brown, prodr. 509 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.V,97	M.fr.VI,105.
P. rugosa, Cunningham in Benth. lab. 456 (1836)... ...	—	—	—	—	N.S.W.	—	—	B.fl.V,98	
P. marifolia, R. Brown, prodr. 509 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,98	M.fr.VI,105.
P. rhombea, R. Brown, prodr. 509 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,99	
P. spinosa, F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854) ...	—	S.A.	—	V.	—	—	—	B.fl.V,99	M.fr.VI,108;IX,162.
P. cuneata, Bentham in De Cand., prodr. XII, 560 (1848) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.V,99	M.fr.VI,108.
P. linearis, R. Brown, prodr. 509 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,100	M.fr.VI,106.
P. phylicifolia, F. v. M., fragm. I, 19 (1858)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,100	M.fr.I,19;IX,162.
P. decussata, F. v. M., fragm. I, 126 (1858)	—	—	—	V.	—	—	—	B.fl.V,100	M.fr.I,126.
P. empetrifolia, Sieber in Spreng. syst. cur. post. 226 (1827) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,101	
P. lithospermoides, F. v. M., fragm. VI, 107 (1868)	—	—	—	—	—	Q.	—	B.fl.V,101	M.fr.VI,107.
P. Behriana, Schlechtendal, Linnaea XX, 610 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,102	M.fr.VI,106.
P. Wilkieana, F. v. M., fragm. VIII, 230 (1874)...	—	S.A.	—	—	—	—	—	—	M.fr.VIII,230.
P. Baxteri, Cunningham in Benth. lab. 452 (1836)	W.A.	S.A.	—	—	—	—	—	B.fl.V,102	M.fr.VI,106.
P. canaliculata, F. v. M., fragm. VI, 105 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,102	M.fr.VI,105.
P. nivea, Cunningham in Benth. lab. 452 (1836)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,103	M.fr.VI,106.
P. striatiflora, F. v. M. in Schlechtend. Linnaea XXV, 425 (1852)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,103	M.fr.VI,105;IX,162.
P. saxicola, R. Brown, prodr. 509 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,104	M.fr.VI,107;IX,162.
P. debilis, F. v. M., fragm. VIII, 147 (1874)	—	—	—	V.	—	—	—	—	M.fr.VIII,147.
P. odoratissima, Bentham in Mitch. Trop. Austr. 291 (1848) ...	—	—	—	—	—	Q.	—	B.fl.V,104	
P. Eckersleyana, F. v. M., fragm. X, 17 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,17.
P. euphrasioides, Bentham in Mitch. Trop. Austr. 360 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,104	M.fr.VI,108.
P. staurophylla, F. v. M., fragm. IX, 73 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,73.
P. cryptandroides, Cunningham in Benth. lab. 453 (1836) ...	—	—	—	—	N.S.W.	—	—	B.fl.V,105	
P. eurybioides, F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)	—	S.A.	—	—	—	—	—	B.fl.V,105	M.fr.VI,105.
P. ringens, Bentham in Mitch. Trop. Austr. 363 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,106	
P. Leichhardtii, Bentham, Fl. Austr. V, 106 (1870)	—	—	—	—	—	Q.	—	B.fl.V,106	
P. Walteri, F. v. M., fragm. VII, 108 (1870)	—	—	—	V.	N.S.W.	—	—	—	M.fr.VII,108.
P. coccinea, F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,106	M.fr.VI,108;IX,162.
P. chlorantha, F. v. M. in Benth. Fl. Austr. V, 108 (1870) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,108	M.fr.IX,162;X,18.
P. calycina, F. v. M. in Benth. Fl. Austr. V, 107 (1870) ...	—	S.A.	—	—	—	—	—	B.fl.V,107	
P. Grylloana, F. v. M., fragm. X, 17 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,17.

WRIXONIA, F. v. M., fragm. X, 18 (1876).

W. prostantheroides, F. v. M., fragm. X, 18 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,18.

HEMIGENIA, R. Brown, prodr. 502 (1810). (Hemiandra, Colobandra, Atelandra.)

H. macrantha, F. v. M., fragm. I, 210 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,112	M.fr.I,210;XI,19.
H. rigida, Bentham in De Cand. prodr. XII, 563 (1848)... ...	W.A.	—	—	—	—	—	—	B.fl.V,112	
H. ramosissima, Bentham in De Cand. prodr. XII, 565 (1848) ...	W.A.	—	—	—	—	—	—	B.fl.V,113	
H. microphylla, Bentham in De Cand. prodr. XII, 565 (1848) ...	W.A.	—	—	—	—	—	—	B.fl.V,113	M.fr.VI,112;IX,162.
H. incana, Bentham in De Cand. prodr. XII, 566 (1848) ...	W.A.	—	—	—	—	—	—	B.fl.V,113	
H. canescens, Bentham in De Cand. prodr. XII, 566 (1848) ...	W.A.	—	—	—	—	—	—	B.fl.V,114	
H. podalyrina, F. v. M., fragm. VI, 112 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,114	M.fr.VI,112.
H. platyphylla, Bentham in De Cand. prodr. XII, 566 (1848) ...	W.A.	—	—	—	—	—	—	B.fl.V,115	
H. obovata, F. v. M., fragm. XI, 19 (1878)	W.A.	—	—	—	—	—	—	—	B.fl.XI,19
H. glabrescens, Bentham in De Cand. prodr. XII, 566 (1848) ...	W.A.	—	—	—	—	—	—	B.fl.V,115	

H. obtusa, Bentham in De Cand. prodr. XII, 567 (1848)... ...	W.A.	—	—	—	—	—	—	B.fl.V,115	
H. sericea, Bentham in Hueg. enum. 80 (1837)	W.A.	—	—	—	—	—	—	B.fl.V,116	M.fr.VI,112.
H. barbata, Bartling in Lehm. pl. Preiss. I, 360 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,116	
H. curvifolia, F. v. M., fragm. I, 210 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,117	M.fr.I,210.
H. scabra, Bentham, Fl. Austr. V, 117 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,117	
H. humilis, Bentham in De Cand. prodr. V, 117 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,117	
H. westringioides, Bentham in De Cand. prodr. XII, 568 (1848)	W.A.	—	—	—	—	—	—	B.fl.V,117	
H. teretiuscula, F. v. M., fragm. VI, 111 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,118	M.fr.VI,111.
H. brachyphylla, F. v. M., fragm. X, 19 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,19.
H. purpurea, R. Brown, prodr. 502 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,118	M.fr.VI,112.
H. cuneifolia, Bentham, Fl. Austr. V, 118 (1870)...	—	—	—	—	N.S.W.	—	—	B.fl.V,118	
H. Drummondii, Bentham, Fl. Austr. V, 119 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,119	
H. pimelifolia, F. v. M., fragm. VI, 112 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,119	M.fr.VI,112.
H. diplanthera, F. v. M., fragm. VI, 111 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,119	M.fr.VI,111.
H. pungens, F. v. M., fragm. XI, 20 (1878)	W.A.	—	—	—	—	—	—	B.fl.V,109	
H. leiantha, F. v. M., fragm. XI, 20 (1878)	W.A.	—	—	—	—	—	—	B.fl.V,110	
H. loganiacea, F. v. M., fragm. XI, 19 (1878)	W.A.	—	—	—	—	—	—	B.fl.V,112	M.fr.VI,113.
MICROCORYS, R. Brown, prodr. 502 (1810). (Anisandra.)									
M. longifolia, Bentham in De Cand. prodr. XII, 568 (1848) ...	W.A.	—	—	—	—	—	—	B.fl.V,121	
M. longiflora, F. v. M., fragm. VI, 113 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,121	M.fr.VI,113.
M. tenuifolia, Bentham, Fl. Austr. V, 122 (1870)...	W.A.	—	—	—	—	—	—	B.fl.V,122	
M. capitata, Bentham in De Cand. prodr. XII, 568 (1848) ...	W.A.	—	—	—	—	—	—	B.fl.V,122	
M. pimeloides, F. v. M., fragm. I, 156 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,123	M.fr.I,156.
M. subcanescens, Bentham, Fl. Austr. V, 123 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,123	
M. ericifolia, Bentham in De Cand. prodr. XII, 569 (1848) ...	W.A.	—	—	—	—	—	—	B.fl.V,123	
M. glabra, Bentham in De Cand. prodr. XII, 569 (1848)... ...	W.A.	—	—	—	—	—	—	B.fl.V,124	
M. exserta, Bentham, Fl. Austr. V, 124 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,124	
M. virgata, R. Brown, prodr. 502 (1810)	W.A.	—	—	—	—	—	—	B.fl.V,125	
M. Macrediana, F. v. M., fragm. VIII, 231 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,231.
M. barbata, R. Brown, prodr. 502 (1810)	W.A.	—	—	—	—	—	—	B.fl.V,125	
M. lenticularis, F. v. M., fragm. VI, 113 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,126	M.fr.VI,113.
M. obovata, Bentham in De Cand. prodr. XII, 569 (1848) ...	W.A.	—	—	—	—	—	—	B.fl.V,126	
M. purpurea, R. Brown, prodr. 502 (1810)...	W.A.	—	—	—	—	—	—	B.fl.V,126	
WESTRINGIA, Smith in Kongl. Svenska Vetenskaps Akad. Handlingar 171 (1797).									
W. cephalantha, F. v. M., fragm. VI, 110 (1868)...	W.A.	—	—	—	—	—	—	B.fl.V,127	M.fr.VI,110.
W. rosmariniformis, Smith in Vet. Acad. Handl. 171 (1797) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.V,128	M.fr.VI,110;IX,162.
W. Dampieri, R. Brown, prodr. 501 (1810)	W.A.	S.A.	—	—	—	—	—	B.fl.V,129	M.fr.VI,110;IX,163.
W. rigida, R. Brown, prodr. 501 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,129	M.fr.IX,163.
W. angustifolia, R. Brown, prodr. 501 (1810)	—	—	T.	—	—	—	—	—	M.fr.VI,110;IX,163.
W. senifolia, F. v. M. in Transact. phil. Soc. Vict. I, 49 (1854)	—	—	—	V.	—	—	—	B.fl.V,130	M.fr.VI,110;IX,163.
W. longifolia, R. Brown, prodr. 501 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,131	M.fr.VI,110;IX,163.
W. glabra, R. Brown, prodr. 501 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,131	M.fr.VI,110;IX,163.
W. rubiifolia, R. Brown, prodr. 501 (1810)...	—	—	T.	—	—	—	—	B.fl.V,131	M.fr.VI,110.
AJUGA, Linné, gen. plant. 167 (1737), from Scribonius (1529).									
A. australis, R. Brown, prodr. 503 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,136	M.fr.VI,109.
TEUCRIUM, Tournefort, inst. 207, t. 98 (1700), from Dioscorides.									
T. racemosum, R. Brown, prodr. 504 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,133	M.fr.VI,109;IX,162.
T. integrifolium, F. v. M. in Benth. Fl. Austr. V, 133 (1870) ...	—	S.A.	—	—	—	Q.	N.A.	B.fl.V,133	M.fr.IX,162.
T. corymbosum, R. Brown, prodr. 504 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,133	M.fr.VI,110.
T. fililobum, F. v. M. in Benth. Fl. Austr. V, 134 (1870) ...	W.A.	—	—	—	—	—	—	B.fl.V,134	M.fr.IX,162.
T. sessiliflorum, Bentham in De Cand. prodr. XI, 580 (1848) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,134	M.fr.VI,110.
T. argutum, R. Brown, prodr. 504 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,135	M.fr.VI,110;IX,162.

VERBENACEAE.

Adanson, Familles des plantes II, 195 (1763), from B. de Jussieu (1759).

SPARTOTHAMNUS, Cunningham in Loudon's hort. Brit. suppl. (1830).									
S. junceus, Cunningham in Loud. hort. Brit. 600 (1830)... ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,55	M.fr.VI,153;IX,5.
S. teucriiflorus, F. v. M. in Wing's S. Sc. Rec. II, 55 (1882) ...	—	S.A.	—	—	—	—	N.A.	—	
NESOGENES, A. de Candolle, prodr. XI, 703 (1847).									
N. euphrasioides, A. de Cand., prodr. XI, 703 (1847)	—	—	—	—	—	Q.	—	—	
LIPPIA, Houston in Linné, syst. nat. 8 (1735); Linné, gen. plant. 347 (1737). (Zapania.)									
L. nodiflora, Cl. Richard in Mich. fl. bor. Amer. II, 15 (1803) ...	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,35	M.fr.VI,151;IX,4.
L. geminata, Humboldt, Bonpland et Kunth, nov. gen. et spec. pl. II, 215 (1817)	—	—	—	—	—	Q.	—	B.fl.V,35	M.fr.VI,151.
VERBENA, Tournefort, inst. 200, t. 94 (1700), from l'Ecluse (1576).									
V. officinalis, Linné, spec. plant. 11 (1753)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,36	M.fr.VI,153;IX,5.
V. macrostachya, F. v. M., fragm. I, 60 (1858)	—	S.A.	—	—	—	Q.	—	B.fl.V,36	M.fr.I,60;IX,5.
LACHNOSTACHYS, Hooker, icon. plant. t. 414 (1842). (Walcottia, Pycnolachne.)									
L. albicans, Hooker, icon. plant. 414 (1842)	W.A.	—	—	—	—	—	—	B.fl.V,38	M.fr.V,159.
L. Cliftoni, F. v. M., fragm. IX, 3 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,3.
L. verbascifolia, F. v. M., fragm. VI, 158 (1868)...	W.A.	—	—	—	—	—	—	B.fl.V,38	M.fr.VI,158;IX,4.
L. ferruginea, Hooker, icon. plant. 415 (1842)	W.A.	—	—	—	—	—	—	B.fl.V,39	M.fr.VI,159.
L. Walcottii, F. v. M., fragm. II, 140 (1861)	W.A.	—	—	—	—	—	—	B.fl.V,39	M.fr.II,140;VI,159.
NEWCASTLIA, F. v. M. in Hook. Kew Misc. IX, 22 (1857).									
N. cladotricha, F. v. M. in Hook. Kew Misc. IX, 22 (1857) ...	—	S.A.	—	—	—	—	N.A.	B.fl.V,40	M.fr.I,184;III,21;VI,155.
N. bracteosa, F. v. M., fragm. VIII, 49 (1873)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,230;IX,4.

N. chrysotricha, F. v. M., fragm. X, 15 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,15.
N. hexarrhena, F. v. M., fragm. X, 16 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,16.
N. spodiotricha, F. v. M., fragm. III, 21, t. 21 (1862)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.V,40	M.fr.VIII,230;IX,4.
N. cephalantha, F. v. M., fragm. IX, 4 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,4.
PHYSOPSIS, Turczaninow in Bullet. de la Soc. Imper. des Natur. de Mosc. XXII, 34 (1849).									
P. spicata, Turczaninow in Bull. de Mosc. XXII, 35 (1849) ...	W.A.	—	—	—	—	—	—	B.fl.V,41	M.fr.VI,154.
MALLOPHORA, Endlicher in Annal. des Wien. Mus. II, 206 (1838). (Lachnocephalus.)									
M. globiflora, Endlicher in Ann. des Wien. Mus. II, 206 (1838)	W.A.	—	—	—	—	—	—	B.fl.V,41	M.fr.VI,154;X,15.
DICRASTYLIS, Drummond & Harvey in Hook. Kew Misc. VII, 56 (1855). (Dicrastyles.)									
D. Doranii, F. v. M., fragm. VIII, 230 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,230.
D. ochrotricha, F. v. M., fragm. IV, 161 (1864)	—	S.A.	—	—	—	—	N.A.	B.fl.V,42	M.fr.I.60;VI,155;VIII,230
D. Gilesii, F. v. M., fragm. VIII, 229 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.X,15.
D. fulva, Drummond & Harvey in Kew Misc. VII, 56 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.V,43	M.fr.I,236;VI,155.
D. reticulata, Drum. & Harvey in Hook. Kew Misc. VII, 57 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,43	M.fr.VI,155;IX,4.
D. Nicholasii, F. v. M., fragm. X, 15 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,15.
D. parvifolia, F. v. M. fragm. II, 16 (1861)	W.A.	—	—	—	—	—	—	B.fl.V,43	M.fr.VI,155;X,15.
D. Beveridgei, F. v. M., fragm. VIII, 50 (1873)	—	S.A.	—	—	—	—	—	—	M.fr.IX,4.
D. Stoechas, Drummond & Harvey in Kew Misc. VII, 57 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,44	M.fr.VI,155.
D. Lewellini, F. v. M., fragm. XI, 86 (1880)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,50;XI,137.
CHLOANTHES, R. Brown, prodr. Flor. Nov. Holl. 513 (1810). (Pithyrodia, Quoya, Dasymalla.)									
C. Stoechadis, R. Brown, prodr. 514 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,45	M.fr.VI,156.
C. parviflora, Walpers, repertor. bot. syst. IV, 58 (1848) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,46	M.fr.IX,5.
C. coccinea, Bartling in Lehm. pl. Preiss. I, 352 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,48	M.fr.VI,156;IX,5.
C. salvifolia, F. v. M.; Pityrodia, R. Brown, prodr. 513 (1810)...	—	—	—	—	—	Q.	—	B.fl.V,48	M.fr.VI,155;IX,5.
C. hemigenioides, F. v. M., fragm. VI, 156 (1868)	W.A.	—	—	—	—	—	N.A.	B.fl.V,48	M.fr.VI,156.
C. uncinata, Turczaninow in Bull. de Mosc. XXXVI, 194 (1863)	W.A.	—	—	—	—	—	—	B.fl.V,48	M.fr.VI,156.
C. Bartlingii, Lehmann, Ind. sem. hort. Hamb. (1844)	W.A.	—	—	—	—	—	—	B.fl.V,49	M.fr.VI,156.
C. verbascina, F. v. M., fragm. I, 233 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,50	M.fr.IV,80;VI,157;IX,5.
C. stachyodes, F. v. M., fragm. V, 50 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,50	M.fr.VI,158.
C. loxocarpa, F. v. M., fragm. II, 22 (1861)	W.A.	—	—	—	—	—	N.A.	B.fl.V,51	M.fr IV,80;VI,156.
C. dilatata, F. v. M., fragm. VI, 157 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,51	M.fr.VI,157.
C. cuneata, F.v.M.; Quoya, Gaudich. in Freyc. voy. 454 t. 66 (1826)	W.A.	—	—	—	—	—	—	B.fl.V,51	
C. Oldfieldii, F. v. M., fragm. I, 234 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,52	M.fr.IV,80.
C. atriplicina, F. v. M., fragm. I, 235 (1859).	W.A.	—	—	—	—	—	N.A.	B.fl.V,52	M.fr.VI,157;IX,5.
C. paniculata, F. v. M., fragm. IV, 80 (1864)	W.A.	—	—	—	—	—	—	B.fl.V,53	
C. loricata, F. v. M., fragm. X, 14 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,14.
C. halganiacea, F. v. M., fragm. X, 14 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,14.
HEMIPHORA, F. v. M., fragm. X, 13 (1876).									
H. Elderi, F. v. M., fragm. X, 13 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,13.
DENISONIA, F. v. M., fragm. I, 124 (1859).									
D. ternifolia, F. v. M., fragm. I, 124, t. 2 (1859)	—	—	—	—	—	—	N.A.	B.fl.V,54	M.fr.I,124.
CYANOSTEGIA, Turczaninow in Bull. Soc. Mosc. XXII, 35 (1849). (Bunnya.)									
C. Turczaninovii, F. v. M., fragm. VI, 154 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,54	M.fr.IX,5.
C. Bunnyana, F. v. M., fragm. V, 36 (1865)	—	—	—	—	—	—	N.A.	B.fl.V,54	M.fr.VI,153.
CALLICARPA, Linné in act. litter. et scienc. Upsal. 80 (1741).									
C. cana, Linné, mantiss. plant. alt. 198 (1771)	—	—	—	—	—	Q.	N.A.	B.fl.V,56	
C. pedunculata, R. Brown, prodr. 513 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,57	
C. longifolia, Lamarck, Encycl. méth. I, 562 (1783)	—	—	—	—	—	Q.	—	B.fl.V,57	
PREMNA, Linné, mantiss. plant. alter. 154 (1771).									
P. obtusifolia, R. Brown, prodr. 512 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.V,58	M.fr.III,36.
P. integrifolia, Linné, mant. plant. alt. 154 (1771)	—	—	—	—	—	—	N.A.	B.fl.V,59	M.fr.VI,153.
P. limbata, Bentham, Fl. Austr. V, 59 (1870)	—	—	—	—	—	Q.	—	B.fl.V,59	
P. Dallachyana, Bentham, Fl. Austr. V, 59 (1870)	—	—	—	—	—	Q.	—	B.fl.V,59	
P. acuminata, R. Brown, prodr. 512 (1810)	—	—	—	—	—	—	N.A.	B.fl.V,60	M.fr.III,36.
CLERODENDRUM, Burmann, thesaur. Zeylan. 66, t. 29 (1737). (Clerodendron.)									
C. hemiderma, F. v. M. in Benth. Fl. Austral. V, 61 (1870) ...	—	—	—	—	—	Q.	—	B.fl.V,61	M.fr.VI,151;IX,5.
C. inerme, R. Brown, prodr. 511 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,61	M.fr.VI,152.
C. Tracyanum, F. v. M. in Benth. Fl. Austr. V, 62 (1870) ...	—	—	—	—	—	Q.	—	B.fl.V,62	M.fr.V,61.
C. tomentosum, R. Brown, prodr. 510 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,62	M.fr.VI,152.
C. lanceolatum, F. v. M., fragm. III, 145 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.V,63	M.fr.III,145.
C. floribundum, R. Brown, prodr. 511 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,63	M.fr.III,145;IX,5.
C. Cunninghamii, Bentham, Fl. Austr. V, 64 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.V,64	
C. costatum, R. Brown, prodr. 511 (1810)	—	—	—	—	—	Q.	—	B.fl.V,64	
GMELINA, Linné, gen. plant. ed. sec. 526 (1742).									
G. macrophylla, Bentham, Fl. Austr. V, 65 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.V,65	M.fr.IV,128.
G. fasciculiflora, Bentham, Fl. Austr. V, 65 (1870)	—	—	—	—	—	Q.	—	B.fl.V,65	
G. Leichhardtii, F. v. M. in Benth. Fl. Austr. V, 66 (1870) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,66	M.fr.III,58,158;VI,153.
VITEX, Tournefort, inst. 603, t. 373 (1700), from Plinius.									
V. trifolia, Linné, spec. plant. 938 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,66	M.fr.III,59;VI,152;IX,5.
V. Lignum vitae, Cunningham in De Cand. prodr. XI, 692 (1847)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,67	M.fr.III,58;IX,5.
V. acuminata, R. Brown, prodr. 512 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.V,67	M.fr.V,34;VI,152,IX,5.
V. glabrata, R. Brown prodr. 512 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,68	M.fr.VI,153.
FARADAYA, F. v. M., fragm. phytogr. Austr. V, 21 (1865).									
F. splendida, F. v. M., fragm. V, 21 (1865)	—	—	—	—	—	Q.	—	B.fl.V,69	M.fr.VI,153.
AVICENNIA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 27 (1737).									
A. officinalis, Linné, spec. plant. 110 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,69	

MYOPORINAE.

R. Brown, prodr. 514 (1810).

MYOPORUM, Banks & Solander in G. Forster, prodr. 44 (1786). (Bertolonia, Andrewsia, Pogonia, Disoon.)

M. tenuifolium, G. Forster, fl. ins. Austr. prodr. 44 (1786)	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.V,4	M.fr.VI,149.
M. montanum, R. Brown, prodr. 515 (1810)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,4	M.fr.VI,149.
M. deserti, Cunningham in Hueg. enum. 78 (1837)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,5	M.fr.VI,148;VII,110.
M. laxiflorum, Bentham, Fl. Austr. V, 6 (1870)	...	...	...	—	—	—	—	—	Q.	—	B.fl.V,6	M.fr.V,23.
M. insulare, R. Brown, prodr. 515 (1810)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,6	M.fr.VI,149;VII,110.
M. viscosum, R. Brown, prodr. 516 (1810)	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,5	M.fr.VII,109.
M. serratum, R. Brown, prodr. 516 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,4	M.fr.VI,149.
M. oppositifolium, R. Brown, prodr. 516 (1810)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,7	M.fr.VI,149.
M. brevipes, Bentham, Fl. Austr. V, 6 (1870)	...	...	..	—	S.A.	—	—	—	—	—	B.fl.V,6	
M. humile, R. Brown, prodr. 516 (1810)	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,6	M.fr.VI,150.
M. debile, R. Brown, prodr. 516 (1810)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,8	M.fr.VI,149.
M. platycarpum, R. Brown, prodr. 516 (1810)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,7	M.fr.VI,150.
M. Bateae, F. v. M. in Proceed. Linn. Soc. N.S.W. VI, 792 (1881)		—	—	—	—	N.S.W.	—	—	—	M.fr.XI,139.
M. floribundum, Cunningham in Hueg. enum. 78 (1837)	...	—	—	—	V.	N.S.W.	—	—	B.fl.V,8	M.fr.I,126;VI,150.
M. Beckeri, F. v. M. in Benth. Fl. Austr. V, 7 (1870)	...	...	W.A.	—	—	—	—	—	—	B.fl.V,7	M.fr.I,126,244;VI,150.
M. salsoloides, Turcz. in Bull. de Mosc. XXXVI, 226 (1863)	...	W.A.	—	—	—	—	—	—	B.fl.V,8	M.fr.I,126;VI,150.

EREMOPHILA, R. Brown, prodr. 518 (1810). (Pholidia, Stenochilus, Eremodendron, Pseudopholidia, Sentis, Duttonia, Pholidiopsis.)

E. Mackinlayi, F. v. M., fragm. IV, 80 (1864)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,17	M.fr.IV,80.
E. Bowmani, F. v. M., fragm. II, 139 (1863)	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,18	M.fr.II,139.
E. strongylophylla, F. v. M., fragm. X, 87 (1876)	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,87.
E. leucophylla, Bentham, Fl. Austr. V, 18 (1870)	...	...	W.A.	—	—	—	—	—	—	B.fl.V,18	
E. Turtoni, F. v. M., fragm. X, 87 (1876)	...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,87.
E. Forrestii, F. v. M., fragm. VII, 49 (1869)	...	...	W.A.	—	—	—	—	—	—	B.fl.V,18	M.fr.VII,49.
E. eriocalyx, F. v. M., fragm. I, 236 (1859)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,19	M.fr.I,236.
E. Maitlandi, F. v. M. in Benth. Fl. Austr. V, 19 (1870)	...	W.A.	—	—	—	—	—	—	B.fl.V,19	
E. rotundifolia, F. v. M., fragm. I, 207 (1859)	...	...	...	—	S.A.	—	—	—	—	—	B.fl.V,19	M.fr.I,207.
E. oppositifolia, R. Brown, prodr. 518 (1810)	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,20	
E. Mitchelli, Bentham in Mitch. Trop. Austr. 31 (1848)	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,21	M.fr.VI,148.
E. Paisleyi, F. v. M., Rep. Babb. Exped. 17 (1858)	...	...	—	S.A.	—	—	—	—	—	B.fl.V,20	
E. Sturtii, R. Brown, App. to Sturt's Exped. 22 (1849)	...	...	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,21	M.fr.VI,148.
E. exilifolia, F. v. M., fragm. X, 88 (1876)	...	...	...	—	S.A.	—	—	—	—	—	—	M.fr.X,88.
E. Dempsteri, F. v. M., fragm. X, 60 (1876)	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,60.
E. Gibsoni, F. v. M., fragm. VIII, 227 (1874)	...	...	...	—	S.A.	—	—	—	—	—	—	M.fr.VIII,227.
E. Berryi, F. v. M., fragm. VIII, 228 (1874)	...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,228;X,88.
E. Clarkei, F. v. M., fragm. I, 208 (1859)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,21	M.fr.I,208.
E. Gilesii, F. v. M., fragm. VIII, 49 (1873)	...	...	...	—	S.A.	—	—	—	—	—	—	M.fr.VIII,49.
E. Latrobei, F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,22	M.fr.I,125.
E. graciliflora, F. v. M., fragm. I, 208 (1858)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,23	M.fr.I,208.
E. longifolia, F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,23	M.fr.VI,148.
E. Drummondii, F. v. M., fragm. VI, 147 (1868)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,24	M.fr.VI,147.
E. Hughesii, F. v. M., fragm. VIII, 228 (1874)	...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,228.
E. polyclada, F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,24	
E. bignoniflora, F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,25	M.fr.VI,148.
E. Freelingii, F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,25	M.fr.VI,148.
E. Fraseri, F. v. M., fragm. XI, 51 (1878)	...	...	...	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,51.
E. Macdonnelli, F. v. M., Rep. Babb. Exped. 18 (1858)	...	...	—	S.A.	—	—	—	—	—	B.fl.V,22	M.fr.VI,150.
E. Goodwinii, F. v. M., Rep. Babb. Exped. 17 (1858)	...	...	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,25	M.fr.VI,148.
E. Elderi, F. v. M., fragm. VIII, 228 (1874)	...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,228.
E. Willsii, F. v. M., fragm. III, 21, t. 20 (1862)	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.V,26	M.fr.III,21.
E. platycalyx, F. v. M., fragm. V, 109 (1866)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,26	M.fr.V,109.
E. viscida, Endlicher, nov. stirp. dec. 51 (1839)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,26	
E. Brownii, F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,27	M.fr.VI,148.
E. subfloccosa, Bentham, Fl. Austr. V, 28 (1870)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,28	
E. Oldfieldii, F. v. M., fragm. I, 208 (1859)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,28	M.fr.I,208.
E. Duttonii, F. v. M., Rep. Babb. Exped. 16 (1858)	...	...	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,28	M.fr.VI,148.
E. maculata, F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,29	
E. denticulata, F. v. M., fragm. I, 125 (1859)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,29	M.fr.IV,170.
E. latifolia, F. v. M. in Schlechtend. Linnaea XXV, 428 (1852)	...	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.V,30	M.fr.VI,148.
E. alternifolia, R. Brown, prodr. 518 (1810)	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,30	M.fr.VI,148.
E. Dalyana, F. v. M., fragm. V, 22 (1865)	...	...	...	...	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,10	M.fr.V,22.
E. scoparia, F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,10	M.fr.V,22;VI,148.
E. Delisserii, F. v. M., fragm. V, 108, t. 42 (1865)	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.V,11	M.fr.V,189.
E. crassifolia, F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858)	—	S.A.	—	—	—	—	—	B.fl.V,11	M.fr.VI,148.
E. resinosa, F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858)	W.A.	—	—	—	—	—	—	B.fl.V,11	
E. Behrii, F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858)	—	S.A.	—	—	—	—	—	B.fl.V,12	
E. Woollsiana, F. v. M., fragm. I, 125, t. 7 (1859)	...	...	W.A.	—	—	—	—	—	—	B.fl.V,12	M.fr.VI,151.
E. Christophori, F. v. M., fragm. IX, 120 (1875)	...	...	...	—	S.A.	—	—	—	—	—	—	M.fr.IX,198.
E. brevifolia, F. v. M.; Myoporum, Bartling in pl. Preiss. I, 350 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,12	M.fr.VI,150.
E. Weldii, F. v. M., fragm. VII, 109 (1870)	...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.IX,120.
E. imbricata, F. v. M.; Pholidia, Bentham, Fl. Austr. V, 13 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,13	
E. densifolia, F. v. M., fragm. II, 160 (1861)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,13	M.fr.II,160.
E. gibbosifolia, F. v. M., Rep. Babb. Exped. 18 (1858)	...	...	—	S.A.	—	V.	—	—	—	B.fl.V,13	M.fr.VI,150.
E. divaricata, F. v. M. in Proceed. Roy. Soc. Tasm. III, 293 (1858)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,14	M.fr.IV,48;VI,150.
E. microtheca, F. v. M. in Benth. Fl. Austr. V, 14 (1870)	...	W.A.	—	—	—	—	—	—	B.fl.V,14	M.fr.II,160.
E. adenotricha, F. v. M. in Benth. Fl. Austr. V, 15 (1870)	...	W.A.	—	—	—	—	—	—	B.fl.V,14	
E. santalina, F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858)	—	S.A.	—	—	—	—	—	B.fl.V,15	
E. Youngii, F. v. M., fragm. X, 16 (1876)	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,16.

PEDALINAE.

R. Brown, prodr. Fl. Nov. Holl. 519 (1810).

JOSEPHINIA, Ventenat, Jardin de la Malmaison, 103 t. 67 (1804).

J. grandiflora, R. Brown, prodr. 520 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,556	
J. Imperatricis, Ventenat, Jard. de la Malm. t. 67 (1804) ...	—	—	—	—	—	—	N.A.	B.fl.IV,557	M.fr.VI,163.
J. Eugeniae, F. v. M. in Hook. Kew Misc. IX, 370, t. 11 (1857)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.IV,557	M.fr.VI,163.

ERICACEAE.

A. L. de Jussieu, gen. plant. 159 (1789).

PERNETTYA, Gaudichaud in Ann. des sc. nat. V, 102 (1825).

P. Tasmanica, J. Hooker in Lond. Journ. of Bot. VI, 268 (1847)	—	—	T.	—	—	—	—	B.fl.IV,140	

GAULTIERA, Kalm in Linné, amoen. acad. III, 14, fig. 6 (1751). (Gaultheria.)

G. hispida, R. Brown, prodr. 559 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,141	
G. lanceolata, J. Hooker in Lond. Journ. of Bot. VI, 267 (1847)	—	—	T.	—	—	—	—	B.fl.IV,141	
G. antipoda, G. Forster, florul. ins. Austr. pr. 34 (1786)... ...	—	—	T.	—	—	—	—	B.fl.IV,142	

WITTSTEINIA, F. v. M., fragm. II, 136 (1861).

W. vacciniacea, F. v. M., fragm. II, 136 (1861)	—	—	—	V.	—	—	—	B.fl.IV,139	M.fr.III,166.

EPACRIDEAE.

R. Brown, prodr. 535 (1810).

STYPHELIA, Solander in G. Forster, fl. ins. Austr. prodr. 13 (1786). (Epacris, Forst. (1776) partly, Stiphelia, Ardisia, Peroa, Perojoa, Ventenatia, Cyathodes, Stenanthera, Astroloma, Leucopogon, Melichrus, Acrotriche, Monotoca, Soleniscia, Stomarrhena, Pentataphrus, Mesotriche, Phanerandra, Froebelia, Pentaptelion, Androstoma, Cyathopsis, Lissanthe partly.)

S. adscendens, R. Brown, prodr. 537 (1810)	—	S.A.	T.	V.	—	—	—	B.fl.IV,146	M.fr.VI,36;XI,122.
S. longifolia, R. Brown, prodr. 537 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,147	
S. laeta, R. Brown, prodr. 537 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,147	
S. triflora, Andrews, Bot. Reposit. t. 72 (1799)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,147	M.fr.XI,122.
S. viridis, Andrews, Bot. Reposit. t. 312 (1806)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,148	
S. Hainesii, F. v. M., fragm. IV, 96, t. 28 (1864)...	W.A.	—	—	—	—	—	—	B.fl.IV,148	M.fr.VIII,55.
S. tubiflora, Smith, Specim. of the Bot. of New Holl. 45, t. 14 (1793)	—	—	—	—	N.S.W.	—	—	B.fl.IV,148	
S. tenuiflora, Lindley, Bot. Regist. XXV, App. XXV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,148	M.fr.VI,46;VIII,54.
S. melaleucoides, F. v. M., fragm. IV, 97 (1864)	W.A.	—	—	—	—	—	—	B.fl.IV,149	M.fr.VI,30.
S. pusilliflora, F. v. M., fragm. IV, 105 (1864)	—	S.A.	—	—	—	—	—	B.fl.IV,149	M.fr.I,178;VI,31.
S. Leucopogon, F. v. M., fragm. IV, 97 (1864)	W.A.	—	—	—	—	—	—	B.fl.IV,149	M.fr.III,143;VI,31.
S. lasionema, F. v. M., fragm. VI, 40 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,153	M.fr.VIII,54.
S. macrocalyx, F. v. M., fragm. VI, 37 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,153	M.fr.VI,37.
S. xerophylla, F. v. M., fragm. VI, 38 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,153	M.fr.VI,38.
S. pentapogonea, F. v. M., fragm. VI, 36 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,153	M.fr.VI,36.
S. prostrata, F. v. M.; Astroloma, R. Brown, prodr. 538 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,154	
S. tecta, Sprengel, syst. I, 657 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,154	M.fr.VI,37.
S. Candolleana, F. v. M., fragm. VI, 38 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,154	M.fr.VIII,54.
S. microdonta; Astroloma, F.v.M. in Benth. Fl. Austr. IV, 155 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,155	
S. pallida, Sprengel, syst. I, 658 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,155	M.fr.VI,37.
S. compacta, Sprengel, syst. I, 657 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,155	M.fr.VI,38.
S. humifusa, Persoon, synops. plant. I, 174 (1805)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,156	M.fr.VI,37;VIII,54.
S. Epacridis, F. v. M., fragm. VI, 38 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,156	M.fr.VI,38.
S. Drummondii, F. v. M., fragm. VI, 37 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,157	M.fr.VI,37.
S. microcalyx, F. v. M., fragm. VI, 37 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,157	M.fr.VI,37.
S. Baxteri, F. v. M., fragm. VI, 37 (1867)...	W.A.	—	—	—	—	—	—	B.fl.IV,158	M.fr.IV,97. [122.
S. Sonderi, F. v. M., fragm. VI, 36 (1867)...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,158	M.fr.VIII,54;IX,48; XI,
S. longiflora, F. v. M., fragm. VIII, 54 (1873)	W.A.	—	—	—	—	—	—	B.fl.IV,158	M.fr.VIII,54.
S. pinifolia, F. v. M., fragm. VI, 36 (1867)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,159	M.fr.VI,36.
S. procumbens, Persoon, synops. I, 174 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,162	M.fr.VI,38;VIII,55.
S. urceolata, F. v. M., fragm. VI, 38 (1868)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,162	M.fr.VIII,54.
S. Billardieri, F. v. M., fragm. VI, 43 (1867)	—	—	T.	—	—	—	—	B.fl.IV,169	M.fr.VIII,54.
S. straminea, Sprengel, syst. I, 656 (1825)...	—	—	T.	—	—	—	—	B.fl.IV,169	M.fr.VI,43.
S. Hookeri, F. v. M., fragm. VI, 44 (1867)	—	—	T.	—	—	—	—	B.fl.IV,169	M.fr.VI,44;VIII,54.
S. dealbata, Sprengel, syst. I, 659 (1825)	—	—	T.	—	—	—	—	B.fl.IV,170	M.fr.VI,43;VIII,54.
S. abietina, Labillardière, Nov. Holl. pl. spec. I, 48, t. 68 (1804)	—	—	T.	—	—	—	—	B.fl.IV,170	M.fr.VI,43.
S. Oxycedrus, Labillardière, Nov. Holl. pl. spec. I, 49, t. 69 (1804)	—	—	T.	V.	—	—	—	B.fl.IV,170	M.fr.VI,43.
S. parvifolia, F. v. M. in papers of R. S. Tasm. 86 (1874) ...	—	—	T.	—	—	—	—	B.fl.IV,171	
S. sapida, F. v. M., fragm. VI, 42 (1867)	—	—	—	—	N.S.W.	—	—	B.fl.IV,175	M.fr.VI,42;VIII,54.
S. strigosa, Smith, Specim. of the Bot. of New Holl. 49 (1793)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,176	M.fr.VI,42;IX,48;XI,122.
S. verticillata, Sprengel, syst. I, 656 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,184	M.fr.VI,43;VIII,55.
S. interrupta, Sprengel, syst. I, 656 (1825)...	W.A.	—	—	—	—	—	—	B.fl.IV,184	
S. amplexicaulis, Rudge in Transact. Linn. Soc. VIII, 292, t. 8...	—	—	—	—	N.S.W.	—	—	B.fl.IV,185	M.fr.VI,44.
S. alternifolia, Sprengel, syst. I, 655 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,185	
S. lanceolata, Smith, Specim. of the Bot. of New Holl. 49 (1793)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,185	M.fr.VI,43.
S. Richei, Labillardière, Nov. Holl. pl. spec. I, 44, t. 60 (1804)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,186	M.fr.VI,42;VIII,54.
S. australis, F. v. M., fragm. VI, 43 (1867)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,187	M.fr.IV,37.
S. capitellata, F. v. M., fragm. VI, 31 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,187	M.fr.VI,31.
S. revoluta, Sprengel, syst. I, 657 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,187	M.fr.VI,31.
S. grandiuscula, F. v. M., fragm. VI, 47 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,188	M.fr.VI,47.
S. reflexa, Sprengel, syst. I, 655 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,188	M.fr.VI,32.
S. corifolia, F. v M.; Leucopogon, Endl., nov. stirp. dec. 15 (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,188	
S. distans, Sprengel, syst. I, 655 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,189	M.fr.VI,32.

S. gibbosa, F. v. M.; Leucopogon, Stschegleew in Bull. de Mosc. XXXII, 12 (1859)	W.A.	—	—	—	—	—	B.fl.IV,189	
S. thymifolia, F. v. M.; Leucopogon, Lindley in Benth. Fl. Austr. IV, 189 (1869)	—	—	—	V.	—	—	—	B.fl.IV,189
S. cordata, F. v. M.; Leucop., Sonder in pl. Preiss. I, 313 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,190
S. Bossiaea, F. v. M., fragm. VI, 47 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,190 M.fr.VI,47.
S. hirsuta, F. v. M., fragm. VI, 31 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,191 M.fr.VI,31.
S. collina, Labillardière, Nov. Holl. pl. spec. I, 47, t. 65 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,191 M.fr.VI,45;VIII,54.
S. glacialis, F. v. M.; Leucopogon, Lindl. in Mitch. three Exped. II, 127 (1838)	—	—	—	V.	—	—	—	B.fl.IV,191
S. compacta, F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXXII, 13 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,192
S. squarrosa, F. v. M., fragm. VI, 31 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,192 M.fr.VI,31.
S. microphylla, Sprengel, syst. I, 656 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.IV,192 M.fr.VI,31.
S. tetragona, F. v. M., fragm. VI, 31 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,193 M.fr.VI,31.
S. phyllostachys, F. v. M.; Leuc., Benth., fl. Austr. IV, 193 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,193 M.fr.VI,32.
S. glabella, Sprengel, syst. I, 655 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,194 M.fr.VI,32.
S. semiopposita, F. v. M., fragm. VI, 49 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,194 M.fr.VI,49.
S. florulenta, F. v. M.; Leuc., Bentham, Fl. Austr. IV, 194 (1869)	W.A.	—	—	—	—	—	—	B.fl IV,194
S. striata, Sprengel, syst. I, 656 (1825)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,194
S. lasiostachya, F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXXII, 11 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,195
S. carinata, Sprengel, syst. I, 658 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,195
S. opponens, F. v. M., fragm. VI, 48 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,196 M.fr.VI,48.
S. oppositifolia, F. v. M. fragm. VI, 32 (1867)	W.A.	—	—	—		—	—	B.fl.IV,196 M.fr.VI,32.
S. tamariscina, Sprengel, syst. I, 656 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,197
S. bracteolaris, F. v. M.; Leuc., Benth., Fl. Austr. IV, 197 (1869)	W.A.		—	—	—	—	—	B.fl.IV,197
S. blepharophylla, F. v. M. fragm. VI, 34 (1867)...	W.A.	—	—	—	—	—	—	B.fl.IV,197 M.fr.VI,34.
S. tenuis, F. v. M.; Leucop., De Candolle prodr. VII, 744 (1833)	W.A.	—	—	—	—	—	—	B.fl.IV,197
S. gnaphalioides, F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXXII, 14 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,198
S. concurva, F. v. M., fragm. VI, 36 (1867)	—	S.A.	—	—		—	—	B.fl.IV,198 M.fr.III,144.
S. Gilbertii, F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXII, 15 (1859)	W.A.	—	—	—		—	—	B.fl.IV,198
S. gracilis, Sprengel, syst. I, 658 (1825)	W.A.	—	—	—	—		—	B.fl.IV,199
S. acicularis, F. v. M.; Leucop., Benth., Fl. Austr. IV, 199 (1869)	W.A.	—	—	—	—			B.fl.IV,199
S. cryptantha, F. v. M.; Leuc., Benth., Fl. Austr. IV, 199 (1869)	W.A.	—	—	—		—	—	B.fl.IV,199
S. gracillima, F. v. M., fragm. VI, 34 (1867)	W.A.	—	—	—			—	B.fl.IV,200 M.fr.VI,34.
S. cymbiformis, F. v. M., fragm. VI, 34 (1867)	W.A.	—	—	—	—		—	B.fl.IV,200 M.fr.VI,34.
S. apiculata, Sprengel, syst. I, 656 (1825)	W.A.	—		—	—		—	B.fl.IV,201 M.fr.IV,105;VI,31.
S. polystachya, Sprengel, syst. I, 659 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,201 M.fr.VI,31.
S. virgata, Labillardière, Nov. Holl. pl. spec. I, 46, t. 64 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,201 M.fr.VI,42.
S. pulchella, F. v. M., fragm. VI, 34 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,202 M.fr.VI,34.
S. polymorpha, F. v. M., fragm. VI, 31 (1867)	W.A.	—	—	—	—		—	B.fl.IV,202 M.fr.VI,31.
S. assimilis, F. v. M.; Leucopogon, R. Brown, prodr. 545 (1810)	W.A.	—	—	—	—	-	—	B.fl.IV,202 M.fr.IV,106;VI,32.
S. Oldfieldii, F.v.M.; Leucopogon, Bentham, Fl. Austr. IV, 203 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,203
S. cucullata, Sprengel, syst. I, 656 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,203 M.fr.VI,32.
S. sprengelioides, F.v.M.; Leuc., Sonder in pl. Preiss. I, 319 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,204 M.fr.VI,34.
S. obtusata, F. v. M.; Leuc., Sonder in pl. Preiss. I, 313 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,204
S. fimbriata, F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXXII, 17 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,204
S. ozothamnoides, Leuc., F. v. M. in Benth. fl. Austr. IV, 205 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,204 M.fr.VI,155.
S. plumuliflora, F. v. M., fragm. VI, 29 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,205 M.fr.XI,122.
S. unilateralis, F. v. M.; Leucopogon, Stschegl. in Bull. de Mosc. XXXII, 19 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,205
S. montana, F. v. M., fragm. VI, 45 (1867)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,206 M.fr.VI,45.
S. Macraei, F. v. M., fragm. VI, 46 (1867)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,206 M.fr.VI,46;VIII,54.
S. pleurandroides, F. v. M., fragm. VI, 32 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,206 M.fr.III,143.
S. linifolia, F. v. M., fragm. VI, 36 (1867)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,207 M.fr.VI,36.
S. pluriloculata, F. v. M., fragm. VI, 32 (1867)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,207 M.fr.I,37.
S. pleiosperma, F. v. M., fragm. VI, 41 (1867)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,207 M.fr.VI,41.
S. rubicunda, F. v. M., fragm. VI, 31 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,208 M.fr.IV,99.
S. attenuata, F. v. M.; Leucopogon, Cunningham in Field's N.S.W. 341 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.IV,208
S. conferta, F. v. M.; Leuc., Bentham, Fl. Austr. IV, 208 (1869)	—	—	—	—	N.S.W.	—	—	B.fl.IV,208
S. mutica, F. v. M., fragm. VI, 45 (1867)	—	—	—	—	N.S.W.	—	—	B.fl.IV,209 M.fr.VI,45.
S. ericoides, Smith, Specim. of the Bot. of New Holl. 48 (1793)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,209 M.fr.VI,45.
S. brevicuspis, F. v. M.; Leuc., Benth., Fl. Austr. IV, 210 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,210
S. propinqua, Sprengel, syst. I, 658 (1825)...	W.A.	—	—	-		—	—	B.fl.IV,210 M.fr.VI,34.
S. subulifolia, F. v. M., fragm. VI, 33 (1867)	W.A.	—	—	—		—	—	B.fl.IV,211 M.fr.IV,103.
S. Allittii, F. v. M., fragm. VI, 34 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,211 M fr.IV,103.
S. racemulosa, F. v. M., fragm. VI, 33 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,211 M.fr.VI,33.
S. pendula, Sprengel, syst. I, 657 (1825)	W.A.	—	-	—	-	—	—	B.fl.IV,212 M.fr.VI,33.
S. concinna, F. v. M.; Leuc., Bentham, Fl. Austr. IV, 212 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,212
S. margarodes, Sprengel, syst. I, 657 (1825)	—	—		—	N.S.W.	Q.	—	B.fl.IV,213 M.fr.VI,36.
S. flavescens, F. v. M., fragm. VI, 33 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,213 M.fr.IV,100.
S. blepharolepis, F. v. M., fragm. VI, 48 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,213 M.fr.VI,48.
S. esquamata, Sprengel, syst. I, 658 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.IV,214
S. rotundifolia, Sprengel, syst. I, 655 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,214
S. cordifolia, F. v. M., fragm. VIII, 54 (1873)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,214 M.fr.VI,45.
S. megacarpa, F. v. M., fragm. VI, 32 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,215 M fr.IV,102.

S. ruscifolia, Sprengel, syst. I, 656 (1825)	—	—	—	-	—	Q.	—	B.fl.IV,215	
S. imbricata, Sprengel, syst. I, 656 (1825)	—	—	—	—	—	Q.	—	B.fl.IV,215	
S. cuspidata, Sprengel, syst. I, 657 (1825)	—	—	—	—	—	Q.	—	B.fl.IV,216	
S. leptospermoides, Sprengel, syst. I, 659 (1825)	—	—	-	—	—	Q.	—	B.fl.IV,216	
S. acuminata, Sprengel, syst. I, 659 (1825)	—	—	—	—	—	—	N.A.	B.fl.IV,216	
S. flexifolia, Sprengel, syst. I, 659 (1825)	—	—	—	—	—	Q.	—	B.fl.IV,217	
S. biflora, Sprengel, syst. I, 659 (1825)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,217	M.fr.IV,104.
S. setigera, Sprengel, syst. I, 659 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.IV,217	M.fr.VI,45.
S. exolasia, F. v. M., fragm. VI, 34 (1867)...	—	—	—	—	N.S.W.	—	—	B.fl.IV,217	M.fr.VI,34.
S. Fraseri, F. v. M.; Leucop., Cunn. in Ann. nat. hist. II. 47 (1840)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,218	M.fr.IV,105;VI,46.
S. hirtella, F. v. M.; Leucopogon in Benth. Fl. Austr. IV, 218 (1869)	—	S.A.	—	—	—	—	—	B.fl.IV,218	
S. Oldfieldii, F.v.M.; Leuc. ovalifolius, Sond. in pl. Preiss. I, 324 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,218	
S. erubescens, F. v. M., fragm. VI, 33 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,219	M.fr.IV,102.
S. lissanthoides, F. v. M., fragm. VI, 33 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,219	M.fr.IV,101.
S. stricta, F. v. M.; Leucopogon, Bentham, Fl. Austr. IV, 219 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,219	
S. Mitchellii, F. v. M.; Leucopogon, Benth. Fl. Austr. IV, 220 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,220	
S. juniperina, Sprengel, syst. I, 658 (1825)...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,220	M.fr.IV,104;VI,46.
S. rufa, F. v. M., fragm. VI, 46 (1867)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,221	M.fr.VIII,54;IX,48.
S. conostephioides, F. v. M., fragm. VI, 34 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,221	M.fr.VI,34.
S. deformis, Sprengel, syst. I, 658 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.IV,221	
S. pogonocalyx; Leucop., F.v.M. in Benth. Fl. Austr. IV, 222 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,222	
S. breviflora; Leucopogon, F. v. M., fragm. IV, 102 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.IV,222	M.fr.IV,102.
S. dura, F. v. M.; Leucopogon, Bentham, Fl. Austr. IV, 222 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,222	
S. multiflora, Sprengel, syst. I, 658 (1825)...	W.A.	—	—	—	—	—	—	B.fl.IV,222	M.fr.VI,32.
S. appressa, Sprengel, syst. I, 658 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.IV,223	
S. Neo-Anglica; Leucop., F. v. M. in Benth. Fl. Austr. IV, 222 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,223	
S. obtecta, F. v. M.; Leucopogon, Benth. Fl. Austr. IV, 223 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,223	
S. crassiflora, F. v. M., fragm. VI, 40 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,224	M.fr.VI,40.
S. strongylophylla, F. v. M., fragm. VI, 33 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,224	M.fr.IV,101.
S. crassifolia, F. v. M., fragm. VI, 33 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,224	M.fr.VI,33.
S. corynocarpa, F. v. M.; Leucop., Sonder in pl. Preiss. I, 322 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,224	
S. Woodsii; Leucopogon, F. v. M., fragm. I, 178 (1859)... ...	W.A.	S.A.	—	V.	—	—	—	B.fl.IV,225	M.fr.VI,33.
S. leptantha, F. v. M.; Leucop., Bentham, Fl. Austr. IV, 225 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,225	
S. divaricata, Sprengel, syst. I, 658 (1825)...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,226	M.fr.VI,44.
S. aggregata, Sprengel, syst. I, 657 (1825)...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,227	M.fr.VI,44;VIII,55.
S. serrulata, Labillardière, Nov. Holl. pl. spec. I, 45, t. 62 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,227	M.fr.VI,44.
S. patula, Sprengel, syst. I, 657 (1825)	—	S.A.	—	—	—	—	—	B.fl.IV,227	M.fr.VI,44;XI,122.
S. ovalifolia, Sprengel, syst. I, 656 (1825)	W.A.	S.A.	—	V.	—	—	—	B.fl.IV,228	M.fr.VI,44.
S. ramiflora, Sprengel, syst. I, 659 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,228	M.fr.
S. depressa, Sprengel, syst. I, 655 (1825)	W.A.	S.A.	—	—	—	- -	—	B.fl.IV,228	M.fr.VI,44.
S. fasciculiflora, F. v. M., fragm. VIII, 55 (1873)...	—	S.A.	—	—	—	—	—	B.fl.IV,229	M.fr.VI,44.
S. elliptica, Smith, Specim. of the Bot. of New Holl. 49 (1793)...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,230	M.fr.VI,58.
S. glauca, Labillardière, Nov. Holl. pl. spec. I, 45, t. 61 (1804)	—	—	T.	—	—	—	—	B.fl.IV,230	
S. scoparia, Smith, Specim. of the Bot. of New Holl. 49 (1793)...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,231	M.fr.VI,58;XI,122.
S. ledifolia, F. v. M.; Monotoca, Cunningham in De Cand., prodr. VII, 756 (1839)	—	—	—	—	N.S.W.	—	—	B.fl.IV,231	
S. empetrifolia, F. v. M. in Papers of R. S. Tasm. 86 (1874) ...	—	—	T.	—	—	—	—	B.fl.IV,231	
S. oligarrhenoides, F. v. M., fragm. IX, 47 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,47.
S. minutiflora, F. v. M., fragm. XI, 122 (1881)	W.A.	—	—	—	—	—	—	B.fl.IV,231	M.fr.XI,122.
OLIGARRHENA, R. Brown, prodr. Fl. Nov. Holl. 549 (1810).									
O. micrantha, R. Brown, prodr. 459 (1810)...	W.A.	—	—	—	—	—	—	B.fl.IV,232	
NEEDHAMIA, R. Brown, prodr. Fl. Nov. Holl. 549 (1810).									
N. pumilio, R. Brown, prodr. 549 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,174	
BRACHYLOMA, Sonder in Lehm. pl. Preiss. I, 304 (1845). (Lobopogon, Lissanthe partly.)									
B. daphnoides, Bentham, Fl. Austr. IV, 173 (1869)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,173	M.fr.VI,42;VIII,55.
B. ciliatum, Bentham, Fl. Austr. IV, 173 (1869)	—	S.A.	T.	V.	—	—	—	B.fl.IV,173	M.fr.VI,42.
B. depressum, Bentham, Fl. Austr. IV, 173 (1869)	—	—	T.	V.	—	—	—	B.fl.IV,173	M.fr.I,36;VI,42.
B. Scortechinii, F. v. M., fragm. XI, 121 (1881)	—	—	—	—	—	Q.	—	—	M.fr.XI,121.
B. Preissii, Sonder in Lehm. pl. Preiss. I, 305 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,172	M.fr.VI,39;VIII,55.
B. concolor, F. v. M., fragm. VI, 39 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,172	M.fr.IV,98.
B. ericoides, Sonder in Schlecht. Linnaea XXVI, 247 (1853) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,172	M.fr.IV,98;VI,39;XI,46.
CONOSTEPHIUM, Bentham in Hueg. enum. 76 (1837). (Conostephiopsis.)									
C. pendulum, Bentham in Hueg. enum. 76 (1837)...	W.A.	—	—	—	—	—	—	B.fl.IV,160	M.fr.VI,40.
C. minus, Lindley, Bot. Regist. XXV, App. XXV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,160	M.fr.VI,40.
C. Roei, Bentham, Fl. Austr. IV, 160 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,160	M.fr.VIII,55.
C. Preissii, Sonder in Lehm. pl. Preiss. I, 304 (1845)	W.A.	—	—	—	—	- -	—	B.fl.IV,161	M.fr.VI,40.
C. planifolium, F. v. M.; fragm. VI, 30 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,161	M.fr.VI,30.
COLEANTHERA, Stschegleew in Bull. Soc. Mosc. XXXII, 4 (1859). (Michiea.)									
C. coelophylla, Bentham, Fl. Austr. IV, 150 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,150	
C. myrtoides, Stschegleew in Bull. de Mosc. XXXII, 4 (1859) ...	W.A.	—	—	—	—	—	—	B.fl.IV,150	M.fr.VI,80.
C. virgata, Stschegleew in Bull. de Mosc. XXXII, 5 (1859) ...	W.A.	—	—	—	—	—	—	B.fl.IV,151	
TROCHOCARPA, R. Brown, prodr. 548 (1810). (Decaspora, Pentachondra.)									
T. laurina, R. Brown, prodr. 548 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,166	M.fr.VI,57;IX,48.
T. disticha, Sprengel, syst. I, 660 (1825)	—	—	T.	—	—	—	—	B.fl.IV,166	M.fr.VI,57;VIII,55.
T. thymifolia, Sprengel, syst. I, 660 (1825)	—	—	T.	—	—	—	—	B.fl.IV,166	M.fr.VI,57;VIII,55.
T. Clarkei, F. v. M., fragm. VI, 57 (1867)...	—	—	—	V.	—	—	—	B.fl.IV,167	M.fr.VI,57.
T. Gunnii, Bentham, fl. Austr. IV, 167 (1869)	--	—	T.	—	—	—	—	B.fl.IV,167	M.fr.VIII,55.
T. parviflora, Bentham, Fl. Austr. IV, 167 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,167	

T. involucrata, F. v. M., fragm. VI, 57 (1867)	—	—	T.	—	—	—	—	B.fl.IV,164	M.fr.VIII,55.
T. pumila, F. v. M., fragm. VI, 57 (1867)...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,164	M.fr.VI,57.
T. ericifolia, F. v. M. in Papers R. S. Tasm. 86 (1874)	—	—	T.	—	—	—	—	B.fl.IV,164	
T. verticillata, F. v. M. in Papers R. S. Tasm. 86 (1874) ...	—	—	T.	—	—	—	—	B.fl.IV,164	
EPACRIS, Cavanilles, icon. et descr. pl. IV, 25, t. 344 et 345 (1797). (Archeria.)									
E. micranthera, F. v. M., fragm. VI, 72 (1867)	—	—	T.	—	—	—	—	B.fl.VI,246	M.fr.VIII,56.
E. hirtella, J. Hooker in Lond. Journ. of Bot. VI, 271 (1847) ...	—	—	T.	—	—	—	—	B.fl.VI,246	M.fr.VI,71.
E. longiflora, Cavanilles, icon. et descr. pl. IV, 25, t. 344 (1797)	—	—	—	—	N.S.W.	—	—	B.fl.IV,234	M.fr.VI,70;VIII,56.
E. reclinata, Cunningham in Benth. Fl. Austr. IV, 234 (1869) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,234	M.fr.VIII,56. [122.
E. impressa, Labillardière, Nov. Holl. pl. spec. I, 43, t. 58 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,235	M.fr.VI,70;VIII,56;XI,
E. sparsa, R. Brown, prodr. 551 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,235	M.fr.VIII,53.
E. petrophila, J. Hooker, Fl. Tasman. I, 261 (1860)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,236	
E. rigida, Sieber in Spreng. cur. poster. 64 (1827)...	—	—	—	—	N.S.W.	—	—	B.fl.IV,236	
E. coriacea, Cunningham in Cand. prodr. VII, 763 (1839) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,236	
E. crassifolia, R. Brown, prodr. 551 (1810)...	—	—	—	—	N.S.W.	—	—	B.fl.IV,237	
E. robusta, Bentham, Fl. Austr. IV, 237 (1869)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,237	
E. obtusifolia, Smith, Exot. Bot. I, 77, t. 40 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,237	M.fr.VI,71;VIII,56.
E. myrtifolia, Labillardière, Nov. Holl. pl. spec. I, 41, t. 55 (1804)	—	—	T.	—	—	—	—	B.fl.IV,238	
E. exserta, R. Brown, prodr. 551 (1810)	—	—	T.	—	—	—	—	B.fl.IV,238	
E. mucronulata, R. Brown, prodr. 552 (1810)	—	—	T.	—	—	—	—	B.fl.IV,238	
E. lanuginosa, Labillardière, Nov. Holl. pl. spec. I, 42, t. 57 (1804)	—	S.A.	T.	V.	—	—	—	B.fl.IV,238	M.fr.VIII,56;XI,122.
E. paludosa, R. Brown, prodr. 551 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,239	M.fr.VIII,56.
E. Calvertiana, F. v. M., fragm. VIII, 53 (1874)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,53.
E. heteronema, Labillardière, Nov. Holl. pl. spec. I, 42, t. 56 (1804)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,239	M.fr.VI,71;VIII,56.
E. serpillifolia, R. Brown, prodr. 240 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,240	
E. microphylla, R. Brown, prodr. 550 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,240	
E. acuminata, Bentham, Fl. Austr. IV, 240 (1869)	—	—	T.	—	—	—	—	B.fl.IV,240	
E. apiculata, Cunningham in Field, N.S. Wales, 340 (1825) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,241	
E. pulchella, Cavanilles, icon. IV, 26, t. 345 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,241	M.fr.VI,71;XI,122.
E. purpurascens, R. Brown, prodr. 550 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,241	M.fr.IV,127;VIII,56.
WOOLLSIA, F. v. M., fragm. VIII, 52 (1872). (Lysinema partly.)									
W. pungens, F. v. M., fragm. VIII, 52 (1872)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,243	M.fr.IV,126;VI,70,88; VIII,55;IX,48;XI,122.
LYSINEMA, R. Brown, prodr. 552 (1810).									
L. lasianthum, R. Brown, prodr. 552 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,243	M.fr.III,142,VI,70;VIII, 55. [122.
L. conspicuum, R. Brown, prodr. 552 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,243	M.fr.VI,70;VIII,55;XI,
L. ciliatum, R. Brown, prodr. 552 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,243	M.fr.VI,69,70;VIII,55.
L. fimbriatum, F. v. M., fragm. IV, 125 (1864)	W.A.	—	—	—	—	—	—	B.fl.IV,244	M.fr.VI,70.
L. elegans, Sonder in Lehm. pl. Preiss. I, 327 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,244	M.fr.VI,70.
PRIONOTES, R. Brown, prodr. fl. Nov. Holl. 553 (1810).									
P. cerinthoides, R. Brown, prodr. 553 (1810)	—	—	T.	—	—	—	—	B.fl.IV,246	M.fr.VI,69;XI,122.
COSMELIA, R. Brown, prodr. fl. Nov. Holl. 553 (1810).									
C. rubra, R. Brown, prodr. 553 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,247	M.fr.VI,64.
PONCELETIA, R. Brown, prodr. fl. Nov. Holl. 554 (1810).									
P. sprengelioides, R. Brown, prodr. 554 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,248	M.fr.I,39;VI,60.
P. monticola, Cunningham in De Cand. prodr. VII, 768 (1843)...	—	—	—	—	N.S.W.	—	—	B.fl.IV,249	
SPRENGELIA, Smith in Svensk. Vetens. Acad. Handling. 260 (1794). (Poiretia.)									[VIII,56;XI,122.
S. incarnata, Smith in Vetens. Ac. Handl. 260, t. 8 (1794) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,249	M.fr.I,39;VI,54;IX,48;
ANDERSONIA, R. Brown, prodr. 553 (1810). (Atherocephala, Homalostoma, Sphincterostoma.)									
A. colossea, F. v. M., fragm. VI, 63 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,251	M.fr.VI,63.
A. patricia, F. v. M., fragm. VI, 79 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,251	M.fr.VI,79.
A. grandiflora, Stschegleew in Bull. de Mosc. XXXII, 21 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,252	M.fr.VI,62.
A. setifolia, Bentham, Fl. Austr. IV, 252 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,252	M.fr.VIII,51.
A. involucrata, Sonder in Lehm. pl. Preiss. I, 331 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,252	M.fr.VI,62.
A. homalostoma, Bentham, Fl. Austr. IV, 253 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,253	M.fr.VIII,56.
A. sprengelioides, R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,253	M.fr.VI,64.
A. latiflora, F. v. M., fragm. VI, 61 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,253	M.fr.VI,61.
A. gracilis, De Candolle, prodr. VII, 767 (1843)	W.A.	—	—	—	—	—	—	B.fl.IV,254	
A. aristata, Lindley, Bot. Regist. XXV, App. XXV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,254	M.fr.VI,63.
A. macronema, F. v. M., fragm. VIII, 51 (1873)...	W.A.	—	—	—	—	—	—	—	M.fr.VIII,51.
A. parvifolia, R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,254	M.fr.IV,125;VI,62.
A. depressa, R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,255	
A. coerulea, R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,255	M.fr.VI,64.
A. subulata, Bentham, Fl. Austr. IV, 256 (1869)...	W.A.	—	—	—	—	—	—	B.fl.IV,256	
A. heterophylla, Sonder in Lehm. pl. Preiss. I, 333 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,256	M.fr.IV,124.
A. brachyanthera, F. v. M., fragm. VI, 61 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,256	M.fr.VI,61.
A. brevifolia, Sonder in Lehm. pl. Preiss. I, 332 (1845)... ...	W.A.	—	—	—	—	—	—	B.fl.IV,256	M.fr.VI,61.
A. variegata, Sonder in Lehm. pl. Preiss. I, 334 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,257	
A. micrantha, R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,257	M.fr.VI,61.
RICHEA, R. Brown, prodr. 555 (1810). (Cystenthe, Pilitis.)									
R. sprengelioides, F. v. M., fragm. VI, 68 (1867)	—	—	T.	—	—	—	—	B.fl.IV,258	M.fr.I,38;VI,68.
R. procera, F. v. M., fragm. VI, 68 (1867)	—	—	T.	—	—	—	—	B.fl.IV,259	M.fr.I,38;VI,68.
R. acerosa, F. v. M., fragm. VI, 69 (1867)...	—	—	T.	—	—	—	—	B.fl.IV,259	M.fr.I,38;VI,69.
R. Milligani, F. v. M., fragm. VI, 69 (1867)	—	—	T.	—	—	—	—	B.fl.IV,259	M.fr.I,38;VIII,56.
R. Gunnii, J. Hooker in Lond. Journ. VI, 273 (1847)	—	—	T.	V.	—	—	—	B.fl.IV,260	M.fr.VI,67;XI,122.
R. scoparia, J. Hooker in Lond. Journ. VI, 273 (1847)	—	—	T.	—	—	—	—	B.fl.IV,260	M.fr.VI,68.

R. dracophylla, R. Brown, prodr. 555 (1810)	—	—	T.	—	—	—	—	B.fl.IV,260	M.fr.VI,67.
R. pandanifolia, J. Hooker, Fl. Antarct. I, 50 (1844)	—	—	T.	—	—	—	—	B.fl.IV,261	M.fr.VI,67;IX,48.
DRACOPHYLLUM, Labillardière, Voy. II, 211, t. 40 (1798). (Epacris, Forst. 1776 partly.)									
D. Milligani, Hooker, icon. t. 845 (1852)	—	—	T.	—	—	—	—	B.fl.IV,262	M.fr.VI,65.
D. Fitzgeraldi, F. v. M., fragm. VII, 27 (1869)	—	—	—	—	N.S.W.	—	—	—	M.fr.X,119.
D. secundum, R. Brown, prodr. 556 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,262	M.fr.VI,65;VIII,56.
D. minimum, F. v. M., fragm. I, 39 (1858)	—	—	T.	—	—	—	—	B.fl.IV,265	M.fr.VI,65;VII,56.
SPHENOTOMA, Sweet, Flora Australasica, t. 44 (1828).									
S. squarrosum, G. Don, gen. syst. III, 785 (1834)	W.A.	—	—	—	—	—	—	B.fl.IV,263	M.fr.VI,66.
S. Drummondii, F. v. M.; Dracophyllum, Benth. Fl. Austr. IV, 263 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,263	
S. dracophylloides, Sonder in Lehm. pl. Preiss. I, 33 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.IV,264	M.fr.VI,65.
S. capitatum, Lindley, Bot. Regist. t. 1515 (1832)	W.A.	—	—	—	—	—	—	B.fl.IV,264	
S. gracile, Sweet, Fl. Austral. t. 44 (1828)...	W.A.	—	—	—	—	—	—	B.fl.IV,264	M.fr.VI,66.
S. parviflorum ; Dracophyllum, F. v. M. in Benth. Fl. Austr. IV, 265 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,265	M.fr.VI,66.

APETALEAE GYMNOSPERMEAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 40 (1880).

CONIFERAE.

Haller, enum. stirp. Helv. I, 145 (1742).

DAMMARA, Rumphius, herb. Amboin. II, 174, t. 54 (1741). (Agathis.)												
D. robusta, C. Moore in Transact. pharm. Soc. Vict. II, 174 (1860)	—	—	—	—	—	Q.	—	B.fl.VI,244	M.fr.XI,104.
ARAUCARIA, A. L. de Jussieu, gen. plant. 413 (1789). (Colymbea, Columbea, Eutassa, Eutacta, Altingia.)									
A. Cunninghamii, Aiton in Sweet. hort. Brit. 475 (1827) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,243	
A. Bidwilli, Hooker, Lond. Journ. of Bot. II, 503, t. 18 (1850)...	—	—	—	—	—	Q.	—	B.fl.VI,243	
A. excelsa, R. Brown in Ait. hort. Kew. sec. ed. V, 412 (1813)...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.
ATHROTAXIS, D. Don in Transact. Linn. Soc. XVIII, 171 (1839). (Arthrotaxis, Sequoia.)									
A. cupressoides, D. Don in Trans. Linn. Soc. XVIII, 173, t. 13 (1839)	—	—	T.	—	—	—	—	B.fl.VI,242	
A. laxifolia, Hooker, icon. plant. t. 573 (1843)	—	—	T.	—	—	—	—	B.fl.VI,242	
A. selaginoides, D. Don in Trans. Linn. Soc. XVIII, 172, t. 14 (1839)	—	—	T.	—	—	—	—	B.fl.VI,242	M.fr.XI,104.
CALLITRIS, Ventenat, decas generum novorum 10 (1808). (Frenela, Actinostrobus, Leichhardtia, Octoclinis.)									
C. Macleayana, F. v. M. in Rep. Burdek. Exped. 17 (1860) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,235	M.fr.XI,104.
C. Parlatorei, F. v. M. in Seemann Journ. of Bot. 267 (1866) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,235	M.fr.V,186;XI,104.
C. Roei, Bentham & J. Hooker, gen. plant. III, 424 (1880) ...	W.A.	—	—	—	—	—	—	B.fl.VI,236	
C. Drummondi, Bentham & J. Hooker, gen. plant. III, 424 (1880)	W.A.	—	—	—	—	—	—	B.fl.VI,236	
C. verrucosa, R. Brown in Mém. du Mus. Par. XIII, 74 (1826)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,236	M.fr.V,237.
C. columellaris, F. v. M., fragm. V, 198 (1866)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.V,198.
C. Muelleri, Bentham & J. Hooker, gen. plant. III, 424 (1880)...	—	—	—	—	N.S.W.	—	—	B.fl.VI,237	
C. cupressiformis, Ventenat, decas gen. nov. 10 (1808)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,237	
C. calcarata, R. Brown in Mém. du Mus. Par. XIII, 74 (1826)...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,238	
C. oblonga, L. Cl. Richard, comment. de conif. 49, t. 18 (1826)	—	—	T.	—	—	—	—	B.fl.VI,238	
C. Actinostrobus, F. v. M., Rep. Burdek. Exped. 19 (1860) ...	W.A.	—	—	—	—	—	—	B.fl.VI,240	
C. acuminata, F. v. M.; Actinostrobus, Parlatore, enum. sem. hort. Florent. 25 (1862)	W.A.	—	—	—	—	—	—	B.fl.VI,240	M.fr.XI,104.
FITZROYA, J. Hooker in Journ. of the hort. Soc. VI, 264 (1851). (Diselma.)									
F. Archeri, Bentham & J. Hooker, gen. pl. III, 426 (1880) ...	—	—	T.	—	—	—	—	B.fl.VI,240	
PHEROSPHAERA, Archer in Hooker's Kew Misc. II, 52 (1850). (Dacrydium partly.)									
P. Hookeriana, Archer in Hooker, Kew Misc. II, 52 (1850) ...	—	—	T.	—	—	—	—	B.fl.VI,245	M.fr.XI,103.
P. Fitzgeraldi, F. v. M. in J. Hooker, icon. pl. XIV, 64, t. 1383 (1882)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,102,138.
MICROCACHRYS, J. Hooker in Lond. Journ. of Bot. IV, 149 (1845). (Dacrydium partly.)									
M. tetragona, J. Hooker, Fl. Tasman. I, 358, t. 100 (1860) ...	—	—	T.	—	—	—	—	B.fl.VI,241	
DACRYDIUM, Solander in G. Forster, pl. escul. insul. ocean. austr. 80 (1786).									
D. Franklinii, J. Hooker in Lond. Journ. of Bot. IV, 152, t. 6 (1845)	—	—	T.	—	—	—	—	B.fl.VI,245	
NAGEIA, Gaertner, de fructib. t. 39 (1788). (Podocarpus.)									
N. elata, F. v. M.; Podoc., R. Brown in Mém. de Mus. XIII, 75 (1826)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,247	
N. spinulosa, F. v. M.; Podocarpus, R. Brown in Mém. du Mus. XIII, 75 (1826)	—	—	—	—	N.S.W.	—	—	B.fl.VI,247	
N. Drouyniana ; Podocarpus, F. v. M., fragm. IV, 86 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.VI,247	M.fr.VI,251;XI,104.
N. alpina, F. v. M. in papers Roy. Soc. of Tasm. 23 (1879) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,248	M.fr.XI,104.
THALAMIA, Sprengel, Anleit. zur Kenntn. der Gew. zweite Ausg. II, 218 (1817). (Phyllocladus.)									
T. asplenifolia, Sprengel, Anl. zur Kenntn. der Gew. II, 218 (1817)	—	—	T.	—	—	—	—	B.fl.VI,246	M.fr.XI,104.

CYCADEAE.

L. C. Richard in Persoon, synops. 630 (1807).

CYCAS, Linné, hort. Cliffort. 482 (1737).												
C. media, R. Brown, prodr. 348 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VI,249	M.fr.VIII,171.
C. Normanbyana, F. v. M., fragm. VIII, 169 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,169.
C. Kennedyana, F. v. M. in Melb. Chemist, March (1882) ...	—	—	—	—	—	Q.	—	—	
C. Cairnsiana, F. v. M., fragm. X, 63 (1876)	—	—	—	—	—	Q.	—	—	M.fr.X,63,121.

ENCEPHALARTOS, Lehmann, pugill. VI, 9, t. I, et 3 (1834). (Macrozamia, Lepidozamia, Catakidozamia, Zamia partly.)

E. Pauli Guilielmi, F. v. M. in Trans. Pharm. Soc. Vict. II, 91 (1859) — — — — N.S.W. Q. — B.fl. VI, 251 M.fr. I, 86, 243; IX, 124.
E. tridentatus, Lehmann, pugill. VI, 13 (1834) ... ... — — — — N.S.W. — — B.fl. VI, 253 M.fr. III, 38; XI, 125.
E. spiralis, Lehmann, pugill. VI, 13 (1834)... ... ... — — — — N.S.W. Q. — B.fl. VI, 251 M.fr. I, 41; II, 179; VIII, 173
E. Douglasii, F. v. M.; Macrozamia, Hill (inedit.) ... ... — — — — — Q. — — [167; VIII, 173; XI, 126.
E. Fraseri, Miquel in Versl. K. Akad. Wet. Amst. XV, 368 (1863) W.A. — — — — — — B.fl. VI, 252 M.fr. I, 41, 243; II, 179; III,
E. Moorei, F. v. M., fragm. XI, 125 (1881)... ... ... — — — — — Q. — — M.fr. I, 243; XI, 125.
E. Macdonnelli, F. v. M. in Versl. K. Akad. Wet. Amst. XV, 376 (1863) — S.A. — — — — — — M.fr. IX, 124; XI, 126. [124.
E. Denisonii, F. v. M. in Journ. Pharm. Soc. Vict. II, 90 (1859) — — — — N.S.W. Q. — B.fl. VI, 253 M.fr. I, 41; VIII, 173; IX,

BOWENIA, Hooker, Bot. Magaz. t. 5398 (1863).

B. spectabilis, Hooker, Bot. Mag. t. 5398 (1863) ... ... — — — — — Q. — B.fl. VI, 254 M.fr. V, 171.

MONOCOTYLEDONEAE.

Ray, meth. plant. nov. (1682).

CALYCEAE PERIGYNAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 41 (1880).

ORCHIDEAE.

Haller, enum. stirp. Helv. praefat. 33 (1742).

PHOLIDOTA, Lindley in Hooker, Exotic Flora II, t. 138 (1825).

P. imbricata, Lindley in Hook. Exot. Fl. II, t. 138 (1825) ... — — — — — — Q. — B.fl. VI, 290 M.fr. IV, 163; V, 105.

STURMIA, Reichenbach in Moessler's Handb. II, 1552 (1828). (Liparis, Richard 1818 not of Zoologists 1738.)

S. reflexa, F. v. M., fragm. II, 72 (1861) ... ... — — — — N.S.W. Q. — B.fl. VI, 273 M.fr. III, 165; V, 102.
S. angustilabris, F. v. M., fragm. IV, 164 (1864)... ... ... — — — — — Q. — B.fl. VI, 273 M.fr. IV, 164.
S. coelogynoides, F. v. M., fragm. II, 71 (1861) ... ... — — — — N.S.W. Q. — B.fl. VI, 273 M.fr. IX, 50.
S. habenarina, F. v. M., fragm. IV, 131 (1864) .. .. — — — — — Q. — B.fl. VI, 273 M.fr. V, 102; XI, 53.

OBERONIA, Lindley, gen. and spec. of orchid. pl. 15 (1830). (Titania.)

O. iridifolia, Lindley, gen. and spec. of orchid. pl. 15 (1830) ... — — — — N.S.W. Q. — B.fl. VI, 274 [88.
O. palmicola, F. v. M., fragm. II, 24 (1860) ... ... — — — — N.S.W. Q. — B.fl. VI, 274 M.fr. II, 24; VII, 30; XI, 53,

PHREATIA, Lindley, gen. and spec. of orchid. pl. 63 (1830). (Plexaure.)

P. Limenophylax, G. Reichenbach in Seem. Bonpl. 54 (1857) — — — — — Q. — B.fl. VI, 290 M.fr. X, 64.

DENDROBIUM, Swartz in nov. act. Upsal. VI, 82 (1799). (Thelychiton partly, Pedilonum, Coelandria.)

D. bigibbum, Lindley in Paxt. Flower Gard. III, 25 (1853) ... — — — — — Q. N.A. B.fl. VI, 277 M.fr. VI, 119; XI, 87.
D. dicuphum, F. v. M., fragm. VIII, 28 (1871) ... ... — — — — — — N.A. B.fl. VI, 277 M.fr. VI, 119.
D. Sumneri, F. v. M., fragm. VI, 94 (1868) ... ... — — — — — Q. — B.fl. VI, 278 M.fr. VI, 94.
D. superbiens, G. Reichenbach in Garden. Chron. 516 (1876) ... — — — — — Q. — — M.fr. XI, 22.
D. Phalaenopsis, Fitzgerald in Gardeners' Chronicle, July (1880) — — — — — Q. — —
D. undulatum, R. Brown, prodr. 332 (1810) ... ... — — — — — Q. — B.fl. VI, 278 M.fr. I, 87.
D. Johannis, G. Reichenbach in Garden. Chron. 890 (1865) ... — — — — — Q. — B.fl. VI, 279
D. speciosum, Smith, Exot. Botany I, 17, t. 10 (1804) .. ... — — — V. N.S.W. Q. — B.fl. VI, 279 M.fr. VIII, 249.
D. falconirostre, Fitzgerald in Sydney Morn. Herald, Nov. 18 (1876) — — — — N.S.W. — — — M.fr. XI, 87.
D. tetragonum, Cunningham in Bot. Regist. XXV, Misc. 33 (1839) — — — — N.S.W. Q. — B.fl. VI, 279 M.fr. I, 87; VI, 119.
D. aemulum, R. Brown, prodr. 333 (1810)... ... ... — — — — N.S.W. Q. — B.fl. VI, 280 M.fr. I, 213; XI, 87.
D. Kingianum, Bidwill in Bot. Regist. XXV, Misc. 11 (1839) ... — — — — N.S.W. Q. — B.fl. VI, 280 M.fr. III, 60.
D. brachypus, G. Reichenbach in Garcke, Linnaea XLI, 42 (1876) — — — — N.S.W. — — —
D. gracilicaule, F. v. M., fragm. I, 179 (1859) ... ... — — — — N.S.W. Q. — B.fl. VI, 281 M.fr. IV, 171.
D. Moorei, F. v. M., fragm. VII, 29 (1870) ... ... — — — — N.S.W. — — B.fl. VI, 281 M.fr. XI, 54.
D. agrostophyllum, F. v. M., fragm. VIII, 28 (1871) ... ... — — — — — Q. — B.fl. VI, 282 M.fr. VIII, 28.
D. Baileyi, F. v. M., fragm. VIII, 173 (1874) ... ... — — — — — Q. — — M.fr. VIII, 173.
D. Smilliae, F. v. M., fragm. VI, 94 (1868) ... ... — — — — — Q. — B.fl. VI, 282 M.fr. VII, 63; XI, 87.
D. ophioglossum, G. Reichenbach in Journ. L. Soc. XV, 113 (1876) — — — — — Q. — — M.fr. XI, 54.
D. canaliculatum, R. Brown, prodr. 333 (1810) ... ... — — — — — Q. — B.fl. VI, 282 M.fr. III, 126.
D. Foelschei, F. v. M. in Wing's S. Sc. Record, Oct. (1882) ... — — — — — — N.A. —
D. monophyllum, F. v. M., fragm. I, 189 (1859) ... ... — — — — N.S.W. Q. — B.fl. VI, 283 M.fr. VII, 64.
D. hispidum, A. Richard, sert. Astrol. 13, t. 5 (1833) ... ... — — — — — Q. — B.fl. VI, 283 M.fr. VII, 30.
D. cucumerinum, MacLeay in Bot. Reg., new ser. V, Misc. 58 (1842)— — — — — N.S.W. Q. — B.fl. VI, 283 M.fr. III, 59; VIII, 151.
D. pugioniforme, Cunningham in Bot. Reg. XXV, Misc. 33 (1839) — — — — N.S.W. Q. — B.fl. VI, 284 M.fr. I, 189.
D. rigidum, R. Brown, prodr. 333 (1810) ... ... — — — — — Q. — B.fl. VI, 284 M.fr. VIII, 248; XI, 87.
D. linguiforme, Swartz in Svensk. Vetensk. Ac. Handl. 247 (1800) — — — — N.S.W. Q. — B.fl. VI, 284
D. teretifolium, R. Brown, prodr. 333 (1810) ... ... — — — — N.S.W. Q. — B.fl. VI, 285 M.fr. I, 89.
D Fairfaxii, Fitzgerald & F. v. M. in Sydney Mail 360, Nov. (1872) — — — — N.S.W. — — B.fl. VI, 285 M.fr. X, 122.
D. striolatum, G. Reichenbach in Hamburg Gartenz. 313 (1857) — — T. V. N.S.W. — — B.fl. VI, 285 M.fr. I, 88; VII, 30.
D. Mortii, F. v. M., fragm. II, 93 (1860) ... ... — — — — N.S.W. Q. — B.fl. VI, 286 M.fr. II, 93.
D. Beckleri, F. v. M., fragm. V, 95 (1865) ... ... — — — — N.S.W. Q. — B.fl. VI, 286

BULBOPHYLLUM, Petit-Thouars, Orchid. Afric. t. 95 et 108 (1822). (Bolbophyllum, Thelychiton partly.)

B. Lichenastrum, F. v. M., fragm. VII, 60 (1869) ... ... — — — — — Q. — B.fl. VI, 287 M.fr. X, 119.
B. Baileyi, F. v. M., fragm. IX, 5 (1875) ... ... — — — — — Q. — — M.fr. IX, 5.
B. nematopodum, F. v. M., fragm. VIII, 30 (1873) ... ... — — — — — Q. — B.fl. VI, 287 M.fr. VIII, 130.

B. Shepherdi, F. v. M., fragm. III, 40 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,288	M.fr.I,190.
B. Taylori, F. v. M., fragm. VIII, 150 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,150.
B. aurantiacum, F. v. M., fragm. III, 39 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,288	M.fr.VII,98.
B. argyropus, G. Reichenbach in Garcke, Linneae XLI, 42 (1876)	—	—	—	—	N.S.W.	—	—	—	
B. exiguum, F. v. M., fragm. II, 72 (1861)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,288	M.fr.XI,22.
B. minutissimum, F. v. M., fragm. XI, 53 (1878)...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.V,85;XI,88.
B. Prenticei, F. v. M. in Wing's South. Sc. Record I, 173 (1881)	—	—	—	—	—	Q.	—	—	B.fl.X,139.
B. Elisae, F. v. M., fragm. VI, 120 (1868)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,289	M.fr.VIII,28.
SARCOCHILUS, R. Brown, prodr. 332 (1810). (Thrixpermum, Gunnia, Cleisostoma.)									
S. erectus; Cleisostoma, Fitzgerald, Austr. Orchids part 4 (1877)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,52,139.
S. tridentatus, G. Reichenbach in Walp. Annal. VI, 500 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,296	M.fr.I,192;II,181,VII,98.
S. Armitii, F. v. M., fragm. IX, 49 (1875)..	—	—	—	—	—	Q.	—	—	M.fr.IX,49.
S. Beckleri; Cleisostoma, F. v. M. in Benth. Fl. Austr. VI, 296 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VI,296	M.fr.VIII,248.
S. brevilabris; Cleisostoma, F. v. M., fragm. XI, 87 (1880) ...	—	—	—	—	—	Q.	—	—	M.fr.VIII,248;XI,139.
S. Macphersoni, F. v. M., fragm. VIII, 248 (1874)	—	—	—	—	—	Q.	—	B.fl.VI,297	M.fr.VII,96.
S. Hartmanni, F. v. M., fragm. VIII, 248 (1874)...	—	—	—	—	—	Q.	—	—	M.fr.VIII,248.
S. divitiflorus, F. v. M. in Benth. Fl. Austr. VI, 292 (1870) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,292	M.fr.IX,50;XI,22.
S. falcatus, R. Brown, prodr. 332 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,293	M.fr.I,89;VII,87;X,50.
S. Fitzgeraldi, F. v. M., fragm. VII, 115 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,293	M.fr.VII,115.
S. olivaceus, Lindley, Bot. Regist. Misc. XXV, 32 (1839) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,293	M.fr.I,90,191;VII,97.
S. parviflorus, Lindley, Bot. Regist. Misc. XXIV, 34 (1838) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,294	M.fr.I,89;VII,98.
S. Ceciliae, F. v. M., fragm. V, 42, t. 42 (1865)	—	—	—	—	—	Q.	—	B.fl.VI,294	M.fr.VII,98;XI,22.
S. Hillii, F. v. M., fragm. II, 94 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,295	M.fr.I,88;III,39;VII,98.
S. phyllorrhizus, F. v. M., fragm. V, 201 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,295	M.fr.VII,30.
TAENIOPHYLLUM, Blume, Bijdr. 355 (1825).									
T. Muelleri, Lindley in Benth. Fl. Austr. VI, 291 (1870) ...	—	—	—	—	—	Q.	—	B.fl.VI,291	
ORNITHOCHILUS, Wallich in Lindley, gen. and spec. of orchid. plants 242 (1833). (Saccolabium partly.)									
O. Hillli, Bentham in Journ. Linn. Soc. XVIII, 334 (1880) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,298	M.fr.VII,97.
GEODORUM, Jackson in Andrews, Botan. Reposit. t. 626 (1810). [31.									
G. pictum, Lindley, gen. and spec. of orchid. pl. 175 (1833) ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,299	M.fr.III,24;V,102;VIII,
EULOPHIA, R. Brown in Edwards, Bot. Regist. 686 (1822). (Eulophus 1822.)									
E. venosa, G. Reichenbach in Benth. Fl. Austr. VI, 300 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.VI,300	M.fr.I,61.
E. Fitzalani, F. v. M., fragm. VIII, 30 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,300	M.fr.VII,30.
DIPODIUM, R. Brown, prodr. 330 (1810).									
D. punctatum, R. Brown, prodr. 331 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VI,301	M.fr.X,64.
D. ensifolium, F. v. M., fragm. V, 42 (1865)	—	—	—	—	—	Q.	—	B.fl.VI,301	M.fr.V,42.
CYMBIDIUM, Swartz in nov. act. Upsal. VI, 70 (1799).									
C. Hillii, F. v. M. in Regel's Gartenflora, 138 (1879)	—	—	—	—	—	Q.	—	—	M.fr.XI,88.
C. canaliculatum, R. Brown, prodr. 331 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VI,302	M.fr.XI,88.
C. albuciflorum, F. v. M., fragm. I, 188 (1859)	—	—	—	—	—	Q.	—	B.fl.VI,303	M.fr.I,188.
C. suave, R. Brown, prodr. 331 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,303	M.fr.I,187.
SPATHOGLOTTIS, Blume, Bijdr. 400 (1825).									
S. Paulinae, F. v. M., fragm. VI, 95 (1868)	—	—	—	—	—	Q.	—	B.fl.VI,304	M.fr.X,64.
PHAJUS, Loureiro, Fl. Cochinch. II, 529 (1790).									
P. grandifolius, Loureiro, Fl. Cochinch. II, 529 (1790)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,304	M.fr.I,42;IV,163;X,6
CALANTHE, R. Brown in Edwards, Bot. Regist. 573 (1821).									
C. veratrifolia, R. Brown in Bot. Regist. 573 (1821)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,305	M.fr.VIII,30;IX,50.
GALEOLA, Loureiro, Fl. Cochinch. II, 520 (1790). (Erythrorchis, Ledgeria.) [XI,105.									
G. cassythoides, G. Reichenbach, Xen. Orchid. II, 77 (1869) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,307	M.fr.II,167;V,102;VIII,31;
G. foliata, F. v. M., fragm. VIII, 31 (1872)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,307	M.fr.II,167;V,102;XI,127.
EPIPOGUM, Gmelin, Fl. Sibir. I, 11, t. 2 (1747).									
E. nutans, Lindley in Journ. Linn. Soc. I, 177 (1856)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,308	M.fr.IX,50;XI,105.
GASTRODIA, R. Brown, prodr. 330 (1810).									
G. sesamoides, R. Brown, prodr. 330 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,309	M.fr.VIII,151;X,64.
POGONIA, A. L. de Jussieu, gen. plant. 65 (1789).									
P. uniflora, F. v. M., fragm. V, 201 (1866)...	—	—	—	—	—	Q.	—	B.fl.VI,310	M.fr.V,201.
P. holochila, F. v. M., fragm. V, 200 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,310	M.fr.V,200.
P. Dallachyana, F. v. M. in Benth. Fl. Austr. VI, 310 (1870) ...	—	—	—	—	—	Q.	—	B.fl.VI,310	
P. pachystomoides, F. v. M., fragm. VIII, 174 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,174.
CORYMBORCHIS, Petit-Thouars, hist. des plant. Orchid. recueill. sur les isles d'Afriq. t. 37-38 (1822). (Corymbis.)									
C. veratrifolia, Blume, Collect. des Orchidées 125, t. 42 & 43 (1858)	—	—	—	—	—	Q.	—	B.fl.VI,311	M.fr.VIII,30;X,64.
ETAERIA, Blume, Bijdr. 409 (1825). (Hetaeria, Ramphidia.)									
E. tenuis, Bentham in Journ. Linn. Soc. XVIII, 345 (1880) ...	—	—	—	—	—	Q.	—	B.fl.VI,312	
MICROSTYLIS, Nuttall, Genera of North American plants II, 196 (1818).									
M. Bernaysii, F. v. M., fragm. XI, 21 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,21.
GOODYERA, R. Brown in Aiton. hort. Kew. sec. ed. V, 197 (1813).									
G. viridiflora, Blume, Coll. des Orchid. 41, t. 9 (1858)	—	—	—	—	—	Q.	—	B.fl.VI,313	M.fr.VIII,29. [64.
G. polygonoides, F. v. M., fragm. VIII, 29 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,313	M.fr.VII,30;VIII,29;X,
SPIRANTHES, L. C. Richard in Mém. du Mus. IV, 40 (1818). (Gyrostachis, 1807).									
S. australis, Lindley, Bot. Regist. 823 (1824)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,314	M.fr.X,65.
THELYMITRA, R. & G. Forster, char. gen. 97, t. 49 (1776). (Macdonaldia.)									
T. ixioides, Swartz in Kong. Svensk. Acad. Handl. 228, t. 3 (1800)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,317	
T. circumsepta, Fitzgerald, Austral. Orchid., part 4 (1877) ...	—	—	—	—	N.S.W.	—	—	—	

T. crinita, Lindley, Bot. Regist. XXV, App. XLIX (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,319	M.fr.VI,84.
T. aristata, Lindley, gen. and spec. of orchid. pl. 521 (1840) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,319	
T. longifolia, R. & G. Forster, char. gen. 98, t. 49 (1776) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,319	
T. villosa, Lindley, Bot. Regist. XXV, App. XLIX (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,320	M.fr.V,94,214.
T. tigrina, R. Brown, prodr. 315 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,320	
T. fusco-lutea, R. Brown, prodr. 315 (1810)	W.A.	S.A.	—	—	—	—	—	B.fl.VI,321	M.fr.V,94.
T. stellata, Lindley, Bot. Regist. XXV, App. XLIX (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,321	
T. carnea, R. Brown, prodr. 314 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,321	M.fr.V,98.
T. flexusa, Endlicher, nov. stirp. decad. 23 (1839)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,322	
T. antennifera, J. Hooker, Fl. Tasman. II, 4, t. 101 (1860) ...	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,322	
T. Macmillani, F. v. M., fragm. V, 93 (1865)	—	—	—	V.	—	—	—	B.fl.VI,323	M.fr.V,93.
T. Mackibbinii, F. v. M. in Melb. Chemist 44 (1881)	W.A.	S.A.	—	V.	—	—	—	—	M.fr.XI,139.
T. variegata, Lindley in Benth. Fl. Austr. VI, 323 (1870) ...	W.A.	—	—	—	—	—	—	B.fl.VI,323	M.fr.V,97,98.
T. venosa, R. Brown, prodr. 314 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VI,323	
T. cyanea, Lindley in Benth. Fl. Austr. VI, 323 (1870)... ...	—	—	T.	—	—	—	—	B.fl.VI,323	
EPIBLEMA, R. Brown, prodr. fl. Nov. Holl. 315 (1810).									
E. grandiflorum, R. Brown, prodr. 315 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,324	M.fr.XI,21.
DIURIS, Smith in Transact. Linn. Soc. IV, 222 (1798).									
D. alba, R. Brown, prodr. 316 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,325	M.fr.V,102.
D. punctata, Smith, Exot. Bot. I, 13, t. 8 (1804)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,326	M.fr.V,102.
D. secundiflora, Fitzgerald, Austral. Orchids, part 4 (1877) ...	—	—	—	—	N.S.W.	—	—	—	
D. aurea, Smith, Exot. Bot. 15, t. 9 (1804)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,327	M.fr.V,172.
D. palustris, Lindley, gen. and spec. of orchid. pl. 507 (1840) ...	—	S.A.	T.	V.	—	—	—	B.fl.VI,327	M.fr.V,102.
D. maculata, Smith, Exot. Bot. I, 57, t. 30 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,327	M.fr.V,102.
D. pedunculata, R. Brown, prodr. 316 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,328	M.fr.V,102,173.
D. pallens, Bentham, Fl. Austr. VI, 329 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VI,329	
D. abbreviata, F. v. M. in Benth. Fl. Austr. VI, 329 (1870) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,329	
D. setacea, R. Brown, prodr. 316 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,329	M.fr.V,101.
D. emarginata, R. Brown, prodr. 316 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,330	
D. sulphurea, R. Brown, prodr. 316 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,330	M.fr.V,102,173.
D. aequalis, F. v. M. in Fitzgerald, Austral. orch. part 2 (1876)	—	—	—	—	N.S.W.	—	—	—	M.fr.X,65.
D. dendrobioides, Fitzgerald, Austral. Orchids, part 7 (1882) ...	—	—	—	—	N.S.W.	—	—	—	
D. longifolia, R. Brown, prodr. 316 (1810)...	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,331	M.fr.V,101,172.
D. pauciflora, R. Brown, prodr. 316 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,331	
ORTHOCERAS, R. Brown, prodr. fl. Nov. Holl. 317 (1810).									
O. strictum, R. Brown, prodr. 317 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,332	
CALOCHILUS, R. Brown, prodr. fl. Nov. Holl. 320 (1810).									
C. campestris, R. Brown, prodr. 320 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,315	M.fr.VIII,151.
C. Robertsoni, Bentham, Fl. Austr. VI, 315 (1870)	—	S.A.	—	V.	—	—	—	B.fl.VI,315	
C. paludosus, R. Brown, prodr. 320 (1810)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,316	
CRYPTOSTYLIS, R. Brown, prodr. 317 (1810). (Zosterostylis.)									
C. longifolia, R. Brown, prodr. 317 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,333	M.fr.XI,21.
C. ovata, R. Brown, prodr. 317 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,334	
C. erecta, R. Brown, prodr. 317 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,334	
C. leptochila, F. v. M. in Benth. Fl. Austr. VI, 324 (1870) ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,334	
PRASOPHYLLUM, R. Brown, prodr. 317 (1810). (Genoplesium.)									
P. australe, R. Brown, prodr. 318 (1810)	—	S.A.	T.	V.	—	—	—	B.fl.VI,337	
P. flavum, R. Brown, prodr. 318 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,337	
P. elatum, R. Brown, prodr. 318 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,337	M.fr.V,100.
P. brevilabre, J. Hooker, Fl. Tasman II, 11, t. 110 (1860) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,338	M.fr.V,101.
P. hians, G. Reichenbach, Beitr. zur syst. Pflanzenk. 61 (1871)...	W.A.	—	—	—	—	—	—	B.fl.VI,338	
P. patens, R. Brown, prodr. 318 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,339	M.fr.V,101.
P. fuscum, R. Brown, prodr. 318 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,339	M.fr.V,101.
P. cyphochilum, Bentham, Fl. Austr. VI, 340 (1870)	W.A.	—	—	—	—	—	—	B.fl.VI,340	
P. ovale, Lindley, Bot. Regist. XXV, App. LIV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,341	
P. macrostachyum, R. Brown, prodr. 318 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,341	
P. Fimbria, G. Reichenbach, Beitr. zur syst. Pflanzenk. 60 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,341	
P. striatum, R. Brown, prodr. 318 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,342	M.fr.V,100.
P. parvifolium, Lindley, Bot. Regist. XXV, App. LIV (1839)...	W.A.	—	—	—	—	—	—	B.fl.VI,342	M.fr.V,101.
P. gibbosum, R. Brown, prodr. 381 (1810)...	W.A.	—	—	—	—	—	—	B.fl.VI,342	M fr.V,100.
P. cucullatum, G. Reichenbach, Beitr. zur syst. Pflanzenk. 59 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,343	
P. nigricans, R. Brown, prodr. 319 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,343	M.fr.IX,50.
P. rufum, R. Brown, prodr. 319 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,344	M.fr.V,100.
P. brachystachyum, Lindley, gen. and spec. of orchid. pl. 513 (1840)	—	—	T.	—	—	—	—	B.fl.VI,344	M.fr.V,100.
P. despectans, J. Hooker, Fl. Tasman. II, 13, t. 113 (1860) ...	—	S.A.	T.	V.	—	—	—	B.fl.VI,345	M.fr.V,100.
P. fimbriatum, R. Brown, prodr. 319 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,345	M.fr.V,100.
P. Archeri, J. Hooker, Fl. Tasman. II, 14, t. 113 (1860)... ...	—	—	T.	V.	—	—	—	B.fl.VI,345	M.fr.V,100.
P. intricatum, C. Stuart in Benth. Fl. Austr. VI, 346 (1870) ...	—	—	T.	—	—	—	—	B.fl.VI,346	
P. Woollsii, F. v. M., fragm. V, 100 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.VI,346	M.fr.V,100.
MICROTIS, R. Brown, prodr. fl. Nov. Holl. 320 (1810).									
M. porrifolia, R. Brown, prodr. 320 (1810)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,347	M.fr.X,65;XI,21.
M. media, R. Brown, prodr. 321 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,348	
M. alba, R. Brown, prodr. 321 (1810)	W.A.	—	—	—		—	—	B.fl.VI,348	
M. pulchella, R. Brown, prodr. 321 (1810)...	W.A.	—	—	—	..		—	B.fl.VI,349	
M. atrata, Lindley, Bot. Regist. XXV, App. LIV (1839) ...	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,349	M.fr.I,90,244;X,65.
CORYSANTHES, R. Brown, prodr. 328 (1810). (Coryhas, Nematoceras.)									
C. unguiculata, R. Brown, prodr. 328 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,350	
C. pruinosa, R. Cunningham in N.S. Wales, Magaz. n, 1 (1833)	W.A.	S.A.	T.	V.	N.S.W.	—	—	—	M.fr.IX,49.

C. fimbriata, R. Brown, prodr. 328 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,351	M.fr.X,65.
C. bicalcarata, R. Brown, prodr. 328 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,351	
PTEROSTYLIS, R. Brown, prodr. fl. Nov. Holl. 326 (1810).									
P. ophioglossa, R. Brown, prodr. 326 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,354	
P. concinna, R. Brown, prodr. 326 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,355	M.fr.V,99;XI,105.
P. curta, R. Brown, prodr. 326 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,355	M.fr.V,99.
P. acuminata, R. Brown, prodr. 326 (1810)...	—	—	—	V.	N.S.W.	—	—	B.fl.VI,355	
P. nutans, R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,356	M.fr.V,99.
P. pedaloglossa, Fitzgerald, Austral. Orchids, part 3 (1877) ...	—	—	—	—	N.S.W.	—	—	—	
P. pedunculata, R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,356	M.fr.VIII,249;XI,105.
P. nana, R. Brown, prodr. 327 (1810)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,357	M.fr.V,99.
P. cucullata, R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	—	—	—	B.fl.VI,357	M.fr.V,99.
P. grandiflora, R. Brown, prodr. 327 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,358	M.fr.V,99;XI,105.
P. truncata, Fitzgerald, Austral. Orchids, part 4 (1878)	—	—	—	—	N.S.W.	—	—	—	
P. reflexa, R. Brown, prodr. 327 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,359	M.fr.XI,105.
P. praecox, Lindley, gen. and spec. of Orchid. pl. 388 (1840) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,359	
P. obtusa, R. Brown, prodr. 327 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,360	M.fr.IX,50;XI,53.
P. recurva, Bentham, Fl. Austr. VI, 360 (1870)	W.A.	—	—	—	—	—	—	B.fl.VI,360	
P. parviflora, R. Brown, prodr. 327 (1810)...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,361	M.fr.V,100.
P. barbata, Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,362	M.fr.VIII,151;XI,22,105.
P. turfosa, Endlicher in Lehm. pl. Preiss. II, 5 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,362	
P. mutica, R. Brown, prodr. 328 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,362	M.fr.V,100.
P. rufa, R. Brown, prodr. 327 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,363	M.fr.V,100.
P. Daintreyana, F. v. M. in Benth. Fl. Austr. VI, 360 (1870) ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,360	
P. longifolia, R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,364	M.fr.V,100.
P. vittata, Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,364	M.fr.XI,126.
CALEYA, R. Brown in Aiton, Hort. Kew. sec. ed. V, 204 (1813). (Calmana, 1810.)									
C. major, R. Brown, prodr. 329 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,366	M.fr.XI,105.
C. minor, R. Brown, prodr. 329 (1810)	—	—	T.	—	N.S.W.	—	—	B.fl.VI,366	
C. nigrita, Lindley, Bot. Reg. XXV, App. LIV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,366	
C. Sullivani, F. v. M. in Melb. Chemist 44 (1882)	—	—	—	V.	—	—	—	—	
DRAKAEA, Lindley, Bot. Regist. XXV, App. LV (1839). (Spiculaea, Arthrochilus.)									
D. ciliata, G. Reichenbach, Beitr. zur syst. Pflanzenk. 68 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,367	
D. irritabilis, G. Reichenbach, Beitr. zur syst. Pflanzenk. 68 (1871)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,368	M.fr.VIII,151;XI,53.
D. elastica, Lindley, Bot. Regist. XXV, App. LV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,368	
ACIANTHUS, R. Brown, prodr. fl. Nov. Holl. 321 (1810).									
A. caudatus, R. Brown, prodr. 321 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,369	
A. fornicatus, R. Brown, prodr. 321 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,370	
A. exsertus, R. Brown, prodr. 321 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,370	M.fr.XI,53,105.
A. viridis, J. Hooker, Fl. Tasman. II, 372 (1860)...	—	—	T.	—	—	—	—	B.fl.VI,371	M.fr.IX,50.
CYRTOSTYLIS, R. Brown, prodr. flor. Nov. Holl. 322 (1810).									
C. reniformis, R. Brown, prodr. 322 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,376	
LYPERANTHUS, R. Brown, prodr. 325 (1810). (Burnettia).									
L. nigricans, R. Brown, prodr. 325 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,374	M.fr.VII,134;X,65.
L. Burnettii, F. v. M., fragm. V, 96 (1865)	—	—	T.	V.	—	—	—	B.fl.VI,375	M.fr.V,96;VII,134.
L. Forrestii, F. v. M. in Wing's S. Sc. Rec. II, 55 (1882) ...	W.A.	—	—	—	—	—	—	—	
L. ellipticus, R. Brown, prodr. 325 (1810)...	—	—	—	—	N.S.W.	—	—	B.fl.VI,374	M.fr.VII,133.
ERIOCHILUS, R. Brown, prodr. fl. Nov. Holl. 323 (1810).									
E. autumnalis, R. Brown, prodr. 323 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,372	M.fr.XI,22,105.
E. scaber, Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,372	
E. tenuis, Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,372	
E. dilatatus, Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,372	
E. multiflorus, Lindley, Bot. Regist. XXV, App. LIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,373	M.fr.XI,105.
E. fimbriatus, F. v. M. in Wing's S. Sc. Record II, 152 (1882)...	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,377	M.fr.XI,104.
CALADENIA, R. Brown, prodr. 325 (1810). (Leptoceras, Adenochilus.)									
C. Menziesii, R. Brown, prodr. 325 (1810)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,377	M.fr.V,101;XI,105.
C. Nortoni; Adenochilus, Fitzgerald, Austral. Orch. part 2 (1876)	—	—	—	—	N.S.W.	—	—	—	M.fr.X,64.
C. Cairnsiana, F. v. M., fragm. VII, 31 (1869)	W.A.	—	—	—	—	—	—	B.fl.VI,378	M.fr.VII,31.
C. multiclavia, G. Reichenbach, Beitr. zur syst. Pflanzenk. 64 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,378	
C. discoidea, Lindley, Bot. Regist. XXV, App. LII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,382	
C. Patersoni, R. Brown, prodr. 324 (1810)...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,382	M.fr.V,101.
C. Drummondii, Bentham, Fl. Austr. VI, 383 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,383	
C. hirta, Lindley, Bot. Regist. XXV, App. LII (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,383	
C. Roei, Bentham, Fl. Austr. VI, 383 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,383	
C. Barbarossae, G. Reichenbach, Beitr. z. syst. Pflanzenk. 64 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,383	
C. flava, R. Brown, prodr. 324 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,384	M.fr.V,101;VI,83;XI,105.
C. latifolia, R. Brown, prodr. 324 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,384	M.fr.V,101;XI,105.
C. reptans, Lindley, Bot. Regist. XXV, App. LII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,385	
C. suaveolens, G. Reichenbach, Beitr. zur syst. Pflanzenk. 67 (1871)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,385	M.fr.V,98.
C. serrata, G. Reichenbach, Beitr. zur syst. Pflanzenk. 67 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,386	M.fr.V,99.
C. carnea, R. Brown, prodr. 324 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,386	M.fr.V,101.
C. congesta, R. Brown, prodr. 324 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,387	
C. aphylla, Bentham, Fl. Austr. VI, 387 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,387	
C. coerulea, R. Brown, prodr. 324 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,388	M.fr.V,101;XI,22.
C. deformis, R. Brown, prodr. 324 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,388	M.fr.V,101;XI,105.
C. sericea, Lindley, Bot. Regist. XXV, App. LII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,389	
C. gemmata, Lindley, Bot. Regist. XXV, App. LII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,389	M.fr.V,101.

C. ixioides, Lindley, Bot. Regist. XXV, App. LII (1839) ..	W.A.	—	—	—	—	—	—	B.fl.VI,389	
C. unita, Fitzgerald in Gardeners' Chronicle 461 (1882)	W.A.	—	—	—	—	—	—	—	
CHILOGLOTTIS, R. Brown, prodr. fl. 322 (1810).									
C. diphylla, R. Brown, prodr. 323 (1810) ... —	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,390	M.fr.X,116.
C. trapeziformis, Fitzgerald, Austral. Orchids, part 3 (1877) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.X,116.
C. formicifera, Fitzgerald, Austral. Orchids, part 3 (1877) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.X,116. [116;XI,22.
C. Gunnii, Lindley, gen. and spec. Orchid. 387 (1840)	—	—	T.	V.	—	—	—	B.fl.VI,391	M.fr.VIII,151; IX,50; X,
GLOSSODIA, R. Brown, prodr. 325 (1810).									
G. major, R. Brown, prodr. 326 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,392	
G. minor, R. Brown, prodr. 326 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,392	
G. Brunonis, Endlicher, nov. stirp. dec. 16 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,393	M.fr.VI,83.
G. intermedia, Fitzgerald in Gardeners' Chron. 462 (1882) ...	W.A.	—	—	—	—	—	—	—	
G. emarginata, Lindley, gen. and spec. of Orchid. pl. 424 (1840)	W.A.	—	—	—	—	—	—	B.fl.VI,393	M.fr.XI,22.
HABENARIA, Willdenow, spec. plant. IV, 44 (1805).									
H. trinervis, Wight, icon. plant. Ind. or. t. 1701 (1852)... ...	—	—	—	—	—	—	N.A.	B.fl.VI,394	
H. elongata, R. Brown, prodr. 313 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VI,394	M.fr.VII,15.
H. graminea, Lindley, gen. and spec. of orchid. pl. 318 (1840) ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,394	M.fr.VII,16.
H. ochroleuca, R. Brown, prodr. 313 (1810)	—	—	—	—	—	—	N.A.	B.fl.VI,395	M.fr.V,102.
H. xanthantha, F. v. M., fragm. VII, 16 (1869)	—	—	—	—	—	Q.	—	B.fl.VI,395	M.fr.VII,16.
APOSTASIA, Blume, Bijdr. 423 (1825). (Niemeyera.)									
A. stylidioides, G. Reichenbach in Benth. Fl. Austr. VI, 396 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,396	

SCITAMINEAE.

Linné, philos. botan. 27 (1751).

MUSA, l'Ecluse, exoticorum libri decem 229 (1605).											
M. Banksii, F. v. M., fragm. IV, 132 (1864)	—	—	—	—	—	Q.	—	B.fl.VI,261	M.fr.IV,132.
M. Fitzalani, F. v. M., fragm. IX, 188 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,188.
M. Hillii, F. v. M., fragm. IX, 169 (1875)...	—	—	—	—	—	Q.	—	—	M.fr.IX,169.
CURCUMA, Linné, gen. plant. 332 (1737), from Hermann (1687).									
C. Australasica, J. Hooker, Bot. Mag. t. 5620 (1867)	—	—	—	—	—	Q.	—	B.fl.VI,263	M.fr.VIII,26.
AMOMUM, Linné, gen. plant. 330 (1737).									
A. Dallachyi, F. v. M., fragm. VIII, 25 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,263	M.fr.VIII,25.
ELETTARIA, White and Matton in Transact. Linn. Soc. X, 249 (1809), from Rheede (1692).									
E. Scottiana, F. v. M., fragm. VIII, 24 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,264	M.fr.VIII,24.
ALPINIA, Linné, gen. plant. 332 (1737). (Hellenia.)									
A. racemigera, F. v. M., fragm. VIII, 27 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,365	M.fr.VIII,27.
A. caerulea, Bentham, Fl. Austr. VI, 265 (1873)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,365	M.fr.VIII,26.
A. arctiflora, F. v. M., fragm. VIII, 25 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,366	M.fr.VIII,25.
COSTUS, C. Bauhin, pinax, 36 (1623).									
C. Potierae, F. v. M., fragm. IV, 164 (1864)	—	—	—	—	—	Q.	—	B.fl.VI,296	M.fr.IV,164;VIII,27.
TAPEINOCHEILOS, Miquel in annal. mus. bot. Lugdun. IV, 101, t. 4 (1869).									
T. pungens, Miquel in ann. mus. Lugd. IV, 101, t. 4 (1869) ...	—	—	—	—	—	Q.	—	B.fl.VI,297	M.fr.VIII,26.

IRIDEAE.

Ventenat, Tableau du règne végétal II, 188 (1799).

DIPLARRHENA, Labillardière, Relation du voy. à la rech. de La Per. I, 275, t. 15 (1799).											
D. Moraea, Labillardière, Voy. I, 157, t. 15 (1799)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,400	M.fr.VII,94.
PATERSONIA, R. Brown, in Bot. Magaz. t. 1041 (1807). (Genosiris, 1804.)									
P. glauca, R. Brown, prodr. 304 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,402	M.fr.VII,36.
P. longiscapa, Sweet, Fl. Austral. t. 39 (1828)	—	S.A.	T.	V.	—	—	—	B.fl.VI,402	M.fr.VII,36.
P. occidentalis, R. Brown, prodr. 304 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,403	M.fr.VII,31.
P. umbrosa, Endlicher in Lehm. pl. Preiss. II, 31 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VI,404	M.fr.VII,32.
P. xanthina, F. v. M., fragm. I, 214 (1859)	W.A.	—	—	—	—	—	—	B.fl.VI,404	M.fr.I,214.
P. limbata, Endlicher in Lehm. pl. Preiss. II, 29 (1846)... ...	W.A.	—	—	—	—	—	—	B.fl.VI,404	
P. juncea, Lindley, Bot. Regist. XXV, App. LVIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,404	M.fr.VII,33.
P. Maxwelli, F. v. M. in Benth. fl. Austr. VI, 405 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VI,405	M.fr.VII,34.
P. pygmaea, Lindley, Bot. Regist. XXV, App. LVIII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,405	M.fr.VII,33.
P. sericea, R. Brown in Bot. Mag. t. 1041 (1807)...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,406	M.fr.VII,35.
P. lanata, R. Brown, prodr. 303 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,406	M.fr.VII,35.
P. rudis, Endlicher in Lehm. pl. Preiss. II, 29 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,406	M.fr.VII,35.
P. macrantha, Bentham, Fl. Austr. VI, 407 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,407	
P. glabrata, R. Brown, prodr. 304 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,407	M.fr.VII,35.
P. Drummondii, F. v. M. in Benth. Fl. Austr. VI, 407 (1873) ...	W.A.	—	—	—		—	—	B.fl.VI,407	
P. inaequalis, Bentham, Fl. Austr. VI, 408 (1873)	W.A.	—	—	—	-	—	—	B.fl.VI,408	
P. graminea, Bentham, Fl. Austr. VI, 408 (1873)...	W.A.	—	—	—	—	—	—	B.fl.VI,408	M.fr.XI,23.
P. babianoides, Bentham, Fl. Austr. VI, 408 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,408	
IRIS, Tournefort, inst. 358, t. 186, 188 (1700), from Hippocrates, Dioscorides and Plinius. (Moraea partly.)									
I. Robinsoniana, F. v. M., fragm. VII, 153 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VI,409	M.fr.VII,153.
SISYRINCHIUM, Linné, gen. plant. 273 (1737). (Libertia, Orthrosanthus, Renealmia, Nematostigma.)									
S. paniculatum, R. Brown, prodr. 305 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,413	M.fr.VII,91.
S. pulchellum, R. Brown, prodr. 305 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,413	M.fr.VII,92.
S. cyaneum, Lindley, Bot. Regist. t. 1090 (1827)...	W.A.	S.A.	—	—	—	—	—	B.fl.VI,410	M.fr.VII,92.
S. polystachyum, F. v. M.; Orthrosanthus, Benth. Fl. Austr. VI, 411 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,411	

CAMPYNEMA, Labillardière, Nov. Holl. pl. spec. I, 93, t. 121 (1804).

C. lineare, Labillardière, Nov. Holl. pl. spec. I, 93, t. 121 (1804)	—	—	T.	—	—	—	—	B.fl.VI,415	M.fr.VI,96.

BURMANNIACEAE.

Sprengel, syst. veg. I, 125 (1825).

BURMANNIA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 100 (1737).

B. disticha, Linné, spec. plant. 287 (1753)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,397
B. juncea, Solander in R. Brown, prodr. 265 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VI,397

TACCACEAE.

Presl, reliq. Haenk. I, 149 (1828).

TACCA, R. et G. Forster, char. gen. 65 (1776).

T. pinnatifida, R. & G. Forster, char. gen. 65 (1776)	—	—	—	—	—	Q.	N.A.	B.fl.VI,458

DIOSCORIDEAE.

Du Mortier, Analyse des familles des plantes 57 (1829), from R. Brown (1810).

DIOSCOREA, Plumier, nov. plant. Americ. gen. 9, t. 26 (1703). (Dioscoridea.)

D. hastifolia, Nees in Lehm. pl. Preiss. II, 33 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,461	M.fr.VII,80;XI,20.
D. transversa, R. Brown, prodr. 295 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,460	M.fr.VII,80.
D. sativa, Linné, spec. plant. 1033 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VI,461	M.fr.VII,81.

PETERMANNIA, F. v. M., fragm. II, 92 (1860).

P. cirrosa, F. v. M., fragm. II, 93 (1860)	—	—	—	—	N.S.W.	—	—	B.fl.VI,462	M.fr.II,176.

HYDROCHARIDEAE.

Lamarck and De Candolle, Flore Française, III, 265 (1805).

HYDROCHARIS, Linné, syst. nat. 8 (1735); Linné, gen. plant. 308 (1737).

H. morsus ranae, Linné, spec. plant. 1036 (1753)...	—	—	—	—	—	Q.	—	B.fl.VI,256	M.fr.VI,199.

ENHALUS, L. C. Richard in Mém. de l'Inst. II, 64 (1811).

E. Koenigii, Richard in Mém. de l'Inst. II, 64 (1811)	—	—	—	—	—	Q.	—	—	M.fr.VIII,219

HALOPHILA, Du Petit-Thouars, gen. nov. Madag. n. 6 (1806).

H. ovata, Gaudichaud, Bot. Voy. Freyc. 470, t. 40 (1826) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,182	M.fr.VIII,219.
H. spinulosa, Ascherson in Neumayer, Anleit. zu wiss. Beob. 368 (1875)—	—	—	—	—	Q.	—	B.fl.VII,183	M.fr.VIII,219.

OTTELIA, Persoon, synops. plant. I, 400 (1805). (Damasonium partly.)

O. alismoides, Persoon, synops. plant. I, 400 (1805)	—	—	—	—	—	Q.	N.A.	B.fl.VI,257	M.fr.VI,199.
O. ovalifolia, L. C. Richard in Mém. de l'Inst. II, 78 (1811) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,257	

BLYXA, Noronha in Du Petit-Thouars, gen. nov. Madag. n. 14 (1806).

B. Roxburghii, L. C. Richard, Mém. de l'Inst. II, 77, t. 5 (1811)	—	S.A.	—	—	—	Q.	N.A.	B.fl.VI,258	

VALLISNERIA, Micheli, nov. plant. gen. 12, t. 10 (1729).

V. spiralis, Linné, spec. plant. 1015 (1753)...	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VI,259	

HYDRILLA, L. C. Richard in Mém. de l'Inst. II, 61 (1811).

H. verticillata, Caspary in Bot. Zeitung, Dec. (1856)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,259	

AMARYLLIDEAE.

J. St. Hilaire, Expos. des famill. natur. des pl. I, 134, t. 21 (1805).

HAEMODORUM, Smith in Transact. Linn. Soc. 213 (1798).

H. distichophyllum, Hooker, icon. plant. t. 866 (1852)	—	—	T.	—	—	—	—	B.fl.VI,419	M.fr.VII,118;IX,51.
H. brevicaule, F. v. M., fragm. I, 64 (1858)	—	—	—	—	—	—	N.A.	B.fl.VI,419	M.fr.VII,118.
H. sparsiflorum, F. v. M., fragm. VII, 117 (1870)	W.A.	—	—	—	—	—	—	B.fl.VI,420	M.fr.VII,117.
H. spicatum, R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,420	M.fr.VII,118.
H. brevisepaleum, Bentham, Fl. Austr. VI, 420 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,420	
H. paniculatum, Lindley, Bot. Regist. XXV, App. XLIV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,420	
H. laxum, R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,421	M.fr.VII,118.
H. simplex, Lindley, Bot. Regist. XXV, App. XLIV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,421	
H. simulans, F. v. M., fragm. VII, 117 (1870)	W.A.	—	—	—	—	—	—	B.fl.VI,421	M.fr.VII,117.
H. planifolium, R. Brown, prodr. 300 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,422	M.fr.VII,118.
H. teretifolium, R. Brown, prodr. 300 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,422	M.fr.VII,118.
H. coccineum, R. Brown, prodr. 300 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VI,422	
H. subvirens, F. v. M., fragm. I, 63 (1858)	—	—	—	—	—	—	N.A.	B.fl.VI,423	M.fr.I,63.
H. ensifolium, F. v. M., fragm. I, 64 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.VI,423	M.fr.VII,118.
H. parviflorum, Bentham, Fl. Austr. VI, 423 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,423	
H. leptostachyum, Bentham, Fl. Austr. VI, 423 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,423	
H. tenuifolium, Cunningham in Benth. Fl. Austr. VI, 425 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,423	

PHLEBOCARYA, R. Brown, prodr. 301 (1810).

P. ciliata, R. Brown, prodr. 301 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,424	M.fr.VIII,23.
P. pilosissima, F. v. M. in Benth. fl. Austr. VI, 425 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VI,425	M.fr.VIII,23.
P. filifolia, F. v. M. in Benth. fl. Austr. VI, 425 (1873)... ...	W.A.	—	—	—	—	—	—	B.fl.VI,425	M.fr.VIII,23.

TRIBONANTHES, Endlicher, nov. stirp. dec. I, 27 (1839).

T. brachypetala, Lindley, Bot. Regist. XXV, App. XLIV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,426	M.fr.VIII,23.
T. uniflora, Lindley, Bot. Regist. XXV, App. XLIV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,427	M.fr.VIII,23.
T. australis, Endlicher, nov. stirp. dec. I, 27 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,427	M.fr.VIII,23.
T. variabilis, Lindley, Bot. Regist. XXV, App. XLIV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,427	
T. longipetala, Lindley, Bot. Regist. XXV, App. XLIV (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,427	M.fr.VIII,23.

CONOSTYLIS, R. Brown, prodr. 300 (1810). (Blancoa, Androstemma.)

C. breviscapa, R. Brown, prodr. 301 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,430	
C. vaginata, Endlicher in Lehm. pl. Preiss. II, 23 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VI,431	M.fr.VIII,19.
C. petrophiloides, F. v. M. in Benth. Fl. Austr. VI, 431 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,431	
C. setosa, Lindley, Bot. Regist. XXV, App. XLIV, t. 6 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,431	M.fr.VIII,19.
C. aurea, Lindley, Bot. Regist. XXV, App. XLIV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,432	M.fr.VIII,19.
C. melanopogon, Endlicher in Lehm. pl. Preiss. II, 18 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VI,432	M.fr.VIII,20.
C. setigera, R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,432	M.fr.VIII,20.
C. Psyllium, Endlicher in Lehm. pl. Preiss. II, 21 (1846) ..	W.A.	—	—	—	—	—	—	B.fl.VI,433	
C. villosa, Bentham, Fl. Austr. VI, 433 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,433	
C. Drummondii, Bentham, Fl. Austr. VI, 433 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,433	
C. involucrata, Endlicher in Lehm. pl. Preiss. II, 23 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VI,434	
C. gladiata, Bentham, Fl. Austr. VI, 434 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,434	
C. seorsiflora, F. v. M., fragm. I, 158 (1859)	W.A.	—	—	—	—	—	—	B.fl.VI,435	M.fr.VIII,19.
C. teretiuscula, F. v. M., fragm. VIII, 18 (1872)	W.A.	—	—	—	—	—	—	B.fl.VI,435	M.fr.VIII,18.
C. stylidioides, F. v. M., fragm. VIII, 17 (1872)	W.A.	—	—	—	—	—	—	B.fl.VI,435	M.fr.VIII,17.
C. prolifera, Bentham, Fl. Austr. VI, 436 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,436	
C. racemosa, Bentham, Fl. Austr. VI, 436 (1839)...	W.A.	—	—	—	—	—	—	B.fl.VI,436	
C. candicans, Endlicher, nov. stirp. dec. 20 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,436	M.fr.VI,255.
C. dealbata, Lindley, Bot. Regist. XXV, App. XLV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,436	
C. Preissii, Endlicher in Lehm. pl. Preiss. II, 18 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,437	M.fr.VIII,20.
C. bracteata, Endlicher in Lehm. pl. Preiss. II, 16 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VI,437	M.fr.VIII,20.
C. filifolia, F. v. M., fragm. VIII, 18 (1872)	W.A.	—	—	—	—	—	—	B.fl.VI,438	M.fr.VIII,18.
C. spinuligera, F. v. M. in Benth. Fl. Austr. VI, 438 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VI,438	
C. bromelioides, Endlicher in Lehm. pl. Preiss. II, 18 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VI,438	
C. aculeata, R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,438	M.fr.VIII,19.
C. laxiflora, Bentham, Fl. Austr. VI, 439 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,439	
C. cymosa, F. v. M. in Benth. Fl. Austr. VI, 439 (1873)... ...	W.A.	—	—	—	—	—	—	B.fl.VI,439	
C. serrulata, R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,439	M.fr.VIII,19.
C. caricina, Lindley, Bot. Regist. XXV, App. XLV (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,440	M.fr.VIII,20.
C. Androstemma, F. v. M., fragm. VIII, 19 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,440	M.fr.VIII,19.
C. Bealiana, F. v. M., fragm. IX, 50 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,50.
C. canescens, F. v. M., fragm. VIII, 19 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,441	M.fr.VIII,19.

AGONIZANTHOS, Labillardière, Relat. du voy. à la rech. de La Perouse I, 411 (1798). (Anigosanthus, Macropidia, Macropodia, Schwaegrichenia)

A. rufa, Labillardière, Voy. I, 411, t. 22 (1798)	W.A.	—	—	—	—	—	—	B.fl.VI,442	M.fr.VIII,21.
A. pulcherrima, Hooker, Bot. Mag. t. 4180 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,443	M.fr.VIII,21.
A. flavida, Redouté et De Candolle, les Liliacées t. 176 (1807) ...	W.A.	—	—	—	—	—	—	B.fl.VI,443	M.fr.VIII,21.
A. Preissii, Endlicher in Lehm. pl. Preiss. II, 26 (1846)... ...	W.A.	—	—	—	—	—	—	B.fl.VI,444	M.fr.VIII,23.
A. humilis, Lindley, Bot. Regist. XXV, App. XLVI (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,444	M.fr.VIII,21.
A. viridis, Endlicher in Lehm. pl. Preiss. II, 25 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,445	M.fr.VIII,22.
A. Manglesii, D. Don in Sweet, Brit. Fl. Gard. sec. ser. t. 265 (1836)	W.A.	—	—	—	—	—	—	B.fl.VI,445	M.fr.VIII,22.
A. bicolor, Endlicher in Lehm. pl. Preiss. II, 26 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,446	M.fr.VIII,22.
A. fuliginosus, Hooker, Bot. Mag. t. 4291 (1847)	W.A.	—	—	—	—	—	—	B.fl.VI,447	M.fr.VIII,20.

CURCULIGO, Gaertner, de fruct. I, 63, t. 16 (1788).

C. recurvata, Aiton, hort. Kew. sec. ed. II, 253 (1811)	—	—	—	—	—	Q.	—	B.fl.VI,448	M.fr.VIII,27.
C. ensifolia, R. Brown, prodr. 290 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,448	M.fr.VI,96.

HYPOXIS, Linné, syst. ed decim. 986 (1759).

H. hygrometrica, Labillardière, Nov. Holl. pl. spec. I, 82, t. 108 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,449	M.fr.VI,96.
H. glabella, R. Brown, prodr. 289 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,450	M.fr.VI,96.
H. occidentalis, Bentham, Fl. Austr. VI, 451 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,451	
H. marginata, R. Brown, prodr. 289 (1810)	—	—	—	—	—	—	N.A.	B.fl.VI,451	

DORYANTHES, Correa in Transact. Linn. Soc. VI, 211 (1802).

D. excelsa, Correa in Transact. Linn. Soc. VI, 211, t. 23 (1802)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,452	M.fr.IX,51.
D. Palmeri, W. Hill in Benth. Fl. Austr. VI, 453 (1873) ...	—	—	—	—	—	Q.	—	B.fl.VI,452	M.fr.IX,51.

CRINUM, Linné, gen. plant. 97 (1737).

C. venosum, R. Brown, prodr. 297 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VI,453	M.fr.III,22.
C. Asiaticum, Linné, spec. plant. 292 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VI,454	M.fr.III,23.
C. brachyandrum, Herbert in Bot. Mag. t. 2121 (1820)	—	—	—	—	—	Q.	—	—	M.fr.XI,139.
C. angustifolium, R. Brown, prodr. 297 (1810)	—	—	—	—	—	Q.	—	—	M.fr.XI,139.
C. flaccidum, Herbert in Bot. Mag. t. 2133 (1820)...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,454	M.fr.III,23.
C. uniflorum, F. v. M., fragm. III, 23 (1862)	—	—	—	—	—	Q.	N.A.	B.fl.VI,454	M.fr.VI,251.
C. pedunculatum, R. Brown, prodr. 297 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,455	M.fr.III,23.

EURYCLES, Salisbury in Transact. of the Hort. Soc. I, 337 (1812).

E. silvestris, Salisbury in Transact. Hort. Soc. I, 337 (1812) ...	—	—	—	—	—	Q.	—	B.fl.VI,456	
E. Cunninghamii, Aiton in Loud. Hort. Brit. Suppl. 588 (1832)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,456	

CALOSTEMMA, R. Brown, prodr. fl. Nov. Holl. 297 (1810).

C. album, R. Brown, prodr. 298 (1810)	—	—	—	—	—	—	N.A.	B.fl.VI,458	
C. purpureum, R. Brown, prodr. 298 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,457	
C. luteum, Sims, Bot. Mag. t. 2101 (1819)...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VI,457	

CALYCEAE HYPOGYNAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 44 (1880).

ROXBURGHIACEAE.

Lindley in Wallich, pl. Asiat. rar. III, 50 (1832).

ROXBURGHIA, Jones in Roxburgh, pl. Corom. I, 29, t. 32 (1795). (Stemona 1790.)

R. Javanica, Kunth, enum. V, 288 (1850) ... ... — — — — — Q. N.A. B.fl.VII,1 M.fr.VII,81

LILIACEAE.

Haller, enum. stirp. Helv. I, 279 (1742).

PHORMIUM, R. & G. Forster, charact. gener. 47, t. 24 (1776). (Chlamydia.)

P. tenax, R. & G. Forster, char. gen. 47, t. 24 (1776) ... ... — — — — N.S.W. — — —

SMILAX, Tournefort, inst. 654, t. 421 (1700), from Theophrastos and Dioscorides.

S. glycyphylla, Smith in White, Voy. to N.S.Wales 230 (1790) — — — — N.S.W. Q. — B.fl.VII,7 M.fr.VII,77.

S. australis, R. Brown, prodr. 293 (1810) ... ... — — — V. N.S.W. Q. N.A. B.fl.VII,7 M.fr.VII,78.

RHIPOGONUM, R. & G. Forster, char. gen. 49, t. 25 (1776). (Ripogonum.)

R. album, R. Brown, prodr. 293 (1810) ... ... — — — V. N.S.W. Q. — B.fl.VII,9 M.fr.VII,79.

R. discolor, F. v. M., fragm. VII, 78 (1870) ... ... — — — — N.S.W. Q. — B.fl.VII,9 M.fr.VII,78.

R. Fawcettianum, F. v. M. in Benth. Fl. Austr. VII, 9 (1878)... — — — — N.S.W. Q. — B.fl.VII,9 M.fr.VII,80.

R. Elseyanum, F. v. M., fragm. I, 44 (1858) ... ... — — — — N.S.W. Q. — B.fl.VII,10 M.fr.VII,80.

FLAGELLARIA, Linné, amoen. acad. 396 (1747).

F. Indica, Linné, spec. plant. 333 (1753) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,10

DRYMOPHILA, R. Brown, prodr. fl. Nov. Holl. 292 (1810).

D. cyanocarpa, R. Brown, prodr. 292 (1810) ... ... — — T. V. N.S.W. — — B.fl.VII,12 M.fr.VII,73;IX,191.

D. Moorei, Baker in Journ. Linn. Soc. XIV, 571 (1875)... ... — — — — N.S.W. — — B.fl.VII,13 M.fr.IX,190.

DIANELLA, Lamarck, Encyclop. méthod. II, t. 250 (1786).

D. Tasmanica, J. Hooker, fl. Tasman. II, 57, t. 133 (1860) ... — — T. V. N.S.W. — — B.fl.VII,14 M.fr.VI,121.

D. longifolia, R. Brown, prodr. 280 (1810)... ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,14 M.fr.VI,122.

D. revoluta, R. Brown, prodr. 280 (1810) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,15 M.fr.VI,121;XI,20.

D. caerula, Sims, Bot. Magaz. t. 505 (1799) ... ... — — — — N.S.W. Q. — B.fl.VII,16 M.fr.VI,123.

D. ensifolia, De Candolle & Redouté, Liliac. t. 1 (1802)... ... — — — — — Q. N.A. B.fl.VII,16 M.fr.VI,123.

ASPARAGUS, Tournefort, inst. 300, t. 154 (1700), from Theophrastos, Dioscorides and Plinius. (Asparagopsis.)

A. racemosus, Willdenow, spec. plant. II, 152 (1799) ... ... — — — — — Q. N.A. B.fl.VII,17 M.fr.VII,73.

EUSTREPHUS, R. Brown, prodr. 281 (1810).

E. Brownii, F. v. M., fragm. VII, 73 (1870) ... ... — — — V. N.S.W. Q. — B.fl.VII,18 M.fr.VII,73.

GEITONOPLESIUM, Cunningham in Bot. Magaz. t. 3131 (1832). (Luzuriago partly.)

G. cymosum, Cunningham in Bot. Mag. t. 3131 (1832) ... ... — — — V. N.S.W. Q. — B.fl.VII,19 M.fr.VII,74.

DRACAENA, Vandelli, dissertatio de Dracaena t. 1 (1762). [190.

D. angustifolia, Roxburgh, fl. Indic. II, 155 (1832) ... ... — — — — — Q. N.A. B.fl.VII,20 M.fr.V,194;VI,120;IX,

CORDYLINE, Commerçon in R. Hedwig, gen. plant. 243 (1806). (Charlwoodia.)

C. terminalis, Kunth in act. acad. Berol. 30 (1820) ... ... — — — — N.S.W. Q. — B.fl.VII,21 M.fr.V,196.

C. Murchisoniae, F. v. M., fragm. V, 195 (1866) ... ... — — — — — Q. — B.fl.VII,22 M.fr.V,195.

C. stricta, Endlicher in Ann. Wien. Mus. I, 162 (1838) ... ... — — — — — N.S.W. Q. — B.fl.VII,22 M.fr.V,195;VI,121.

C. Baueri, J. Hooker in Gardeners' Chronicle 792 (1860) ... — — — — N.S.W. — — —

BLANDFORDIA, Smith, Exot. Bot. I, 5, t. 4 (1804).

B. marginata, Herbert in Bot. Regist. Misc. 84 (1842) ... ... — — T. — — — — B.fl.VII,23 M.fr.VII,71.

B. grandiflora, R. Brown, prodr. 296 (1810) ... ... — — — — N.S.W. — — B.fl.VII,23

B. nobilis, Smith, Exot. Bot. I, 5, t. 4 (1804) ... ... — — — — N.S.W. — — B.fl.VII,24 M.fr.VII,70.

B. flammea, Hooker, Bot. Mag. t. 4819 (1854) ... ... — — — — N.S.W. Q. — B.fl.VII,24

HEWARDIA, Hooker, icon. plant. 858 (1852).

H. Tasmanica, Hooker, icon. plant. t. 858 (1852)... ... ... — — T. — — — — B.fl.VII,25 M.fr.VII,72.

ASTELIA, Banks & Solander in R. Brown, prodr. 291 (1810). (Hamelinia.)

A. alpina, R. Brown, prodr. 291 (1810) ... ... — — T. V. N.S.W. — — B.fl.VII,11

MILLIGANIA, J. Hooker in Kew misc. V, 296, t. IX (1853).

M. densiflora, J. Hooker, Fl. Tasman. II, 62 (1860) ... ... — — T. — — — — B.fl.VII,26

M. longifolia, J. Hooker in Kew misc. V, 296, t. IX (1853) ... — — T. — — — — B.fl.VII,26

M. Johnstoni, F. v. M. in Benth. Fl. Austr. VII, 26 (1878) ... — — T. — — — — B.fl.VII,26

M. stylosa, F. v. M. in papers of the R. S. of Tasm. 11 (1876) — — T. — — — — B.fl.VII,27

WURMBEA, Thunberg, nov. gen. I, 18 (1781). (Anguillaria, Melanthium partly, Pleea partly.)

W. tubulosa, Bentham, Fl. Austr. VII, 28 (1878)... ... ... W.A. — — — — — — B.fl.VII,28 M.fr.VII,75.

W. dioica, F. v. M., fragm. X, 119 (1877) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,30 M.fr.VII,74.

IPHIGENIA, Kunth, enum. plant. omn. hucusq. cognit. IV, 212 (1843).

I. Indica, Kunth, enum. plant. IV, 213 (1843) ... ... — — — — — Q. N.A. B.fl.VII,31 M.fr.VII,74;IX,193.

SCHELHAMMERA, R. Brown, prodr. 273 (1810).

S. undulata, R. Brown, prodr. 274 (1810) ... ... — — — V. N.S.W. — — B.fl.VII,31 M.fr.VII,71.

S. multiflora, R. Brown, prodr. 274 (1810)... ... ... — — — — — Q. — B.fl.VII,32 M.fr.VII,72.

KREYSSIGIA, Reichenbach, iconogr. bot. exotic. III, 11, t. 229 (1829). (Kreysigia, Tripladenia.)

K. Cunninghami, F. v. M., fragm. VII, 71 (1870) ... ... — — — — N.S.W. Q. — B.fl.VII,32 M.fr.VII,71.

BURCHARDIA, R. Brown, prodr. 272 (1810).

B. umbellata, R. Brown, prodr. 272 (1810)... ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,33 M.fr.VII,77.

BULBINE, Linné, hort. Cliffort. 122 (1737). (Anthericum partly, Chrysobactron.)

| | | |

B. gracilis, F v. M., fragm. VII, 88 (1870)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,65	M.fr.I,159.
B. ramosa, F. v. M., fragm. VII, 88 (1870)	W.A.	—	—	—	—	—	—	B.fl.VII,66	M.fr.VI,255.
B. sessiliflora, F. v. M. in pap. R. S. of Tasm. 116 (1877) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,66	
B. brachyphylla; Laxm., F. v. M. in Benth. Fl. VII, 67 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,66	M.fr.I,158.
B. sessilis, F.v.M.; Laxm.,Lindl.Bot.Reg. XXV,App.LVI (1839)	W.A.	—	—	—	—	—	—	B.fl.VII,67	
STAWELLIA, F. v. M., fragm. phytogr. Austr. VII, 85 (1870).									
S. dimorphantha, F. v. M., fragm. VII, 85 (1870)	W.A.	—	—	—	—	—	—	B.fl.VII,67	M.fr.VII,85.
JOHNSONIA, R. Brown, prodr. fl. Nov. Holl. 287 (1810).									
J. lupulina, R. Brown, prodr. 287 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,68	M.fr.VII,86.
J. pubescens, Lindley, Bot. Regist. XXV, App. LVII (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VII,68	M.fr.VI,255.
ARNOCRINUM, Endlicher & Lehmann in pl. Preiss. II, 41 (1846).									
A. Drummondii, Endlicher in Lehm. pl. Preiss. II, 41 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,69	M.fr.VII,73.
A. Preissii, Lehmann, pl. Preiss. II, 42 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,70	
BORYA, Labillardiere, Nov. Holl. pl. specim. 81, t. 107 (1804).									
B. nitida, Labillardière, Nov. Holl. pl. spec. 81, t. 107 (1804) ...	W.A.	—	—	—	—	—	—	B.fl.VII,71	M.fr.VII,87.
B. septentrionalis, F. v. M., fragm. 41 (1856)	—	—	—	—	—	Q.	—	B.fl.VII,71	M.fr.V,41.
CALECTASIA, R. Brown, prodr. fl. Nov. Holl. 263 (1810).									
C. cyanea, R. Brown, prodr. 264 (1810)	W.A.	S.A.	—	V.	—	—	—	B.fl.VII,121	
BAXTERIA, R. Brown in Hook. Lond. Journ. of Bot. II, 492, t. 13-15 (1843).									
B. australis, R. Brown in Hook. Lond. Journ. II, 492 (1843) ...	W.A.	—	—	—	—	—	—	B.fl.VII,120	
ACANTHOCARPUS, Lehmann, pl. Preiss. II, 274 (1847). (Chamaexeros).									
A. Preissii, Lehmann, pl. Preiss. II, 274 (1847)	W.A.	—	—	—	—	—	—	B.fl.VII,111	
A. Serra, F. v. M.; Xerotes, Endlicher in pl. Preiss. II, 49 (1847)	W.A.	—	—	—	—	—	—	B.fl.VII,111	M.fr.VIII,211.
A. fimbriatus; Xerotes, F. v. M., fragm. VIII, 211 (1874) ...	W.A.	—	—	—	—	—	—	B.fl.VII,111	M.fr.VIII,211.
XEROTES, R. Brown, prodr. 257 (1810). (Lomandra, 1804.)									
X. Banksii, R. Brown, prodr. 263 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,96	M.fr.VIII,213.
X. dura, F. v. M. in Transact. Vict. Inst. 42 (1854)	—	S.A.	—	—	—	—	—	B.fl.VII,97	M.fr.VIII,207.
X. longifolia, R. Brown, prodr. 262 (1810)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,98	M.fr.VIII,210.
X. rigida, R. Brown, prodr. 262 (1801)	W.A.	—	—	—	—	—	—	B.fl.VII,99	M.fr.VIII,209.
X. Sonderi, F. v. M., fragm. VIII, 206 (1874)	W.A.	—	—	—	—	—	—	B.fl.VII,99	M.fr.
X. odora, Endlicher in Lehm. pl. Preiss. II, 50 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,99	
X. Brownii, F. v. M., fragm. VIII, 206 (1874)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,100	M.fr.VIII,206.
X. Ordii, F. v. M., fragm. XI, 23 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,137.
X. sororia, F. v. M., first gen. Report 15 (1854)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,100	
X. Endlicheri, F. v. M., fragm. VIII, 205 (1874)	W.A.	—	—	—	—	—	—	B.fl.VII,101	M.fr.VIII,205.
X. sericea, Endlicher in Lehm, pl. Preiss. II, 51 (1846)... ...	W.A.	—	—	—	—	—	—	B.fl.VII,101	M.fr.VIII,207.
X. purpurea, Endlicher in Lehm. pl. Preiss. II, 49 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,101	M.fr.VIII,213.
X. Preissii, Endlicher in Lehm. pl. Preiss. II, 50 (1846)... ...	W.A.	—	—	—	—	—	—	B.fl.VII,102	
X. effusa, Lindley in Mitch. Three Exped. II, 151 (1838) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,102	M.fr.VIII,209.
X. micrantha, Endl. in Lehm. pl. Preiss. II, 49 (1846)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,103	M.fr.VIII,212.
X. Thunbergii, F. v. M., fragm. VIII, 208 (1874)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,103	M.fr.VIII,208.
X. caespitosa, Benth. Fl. Austr. VII, 104 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,104	
X. pauciflora, R. Brown, prodr. 261 (1810)...	W.A.	—	—	—	—	—	—	B.fl.VII,104	M.fr.212.
X. flexifolia, R. Brown, prodr. 260 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VII,105	M.fr.VIII,207.
X. glauca, R. Brown, prodr. 260 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,105	M.fr.VIII,208.
X. elongata, Bentham, Fl. Austr. VII, 106 (1878)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VII,106	
X. rupestris, Endlicher in Lehm. pl. Preiss. II, 50 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,106	
X. collina, R. Brown, prodr. 260 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,107	
X. suaveolens, Endlicher in Lehm. pl. Preiss. II, 50 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,107	M.fr.VIII,207.
X. turbinata, Endlicher in Lehm. pl. Preiss. II, 51 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,107	M.fr.VIII,211.
X. spartea, Endlicher in Lehm. pl. Preiss. II, 51 (1846)... ...	W.A.	—	—	—	—	—	—	B.fl.VII,108	M.fr.VIII,211.
X. juncea, F. v. M. in Transact. Vict. Inst. 135 (1855)	—	S.A.	—	—	—	—	—	B.fl.VII,108	M.fr.VIII,210.
X. leucocephala, R. Brown, prodr. 260 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,109	M.fr.VIII,210.
X. hastilis, R. Brown, prodr. 263 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,110	M.frVIII,210;XI,54.
DASYPOGON, R. Brown, prodr. fl. Nov. Holl. 263 (1810).									
D. bromelifolius, R. Brown, prodr. 263 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,118	
D. Hookeri, Drummond in Hook. Lond. Journ. II, 168 (1843) ...	W.A.	—	—	—	—	—	—	B.fl.VII,119	M.fr.II,112;IV,173.
XANTHORRHOEA, Smith in Transact. Linn. Soc. IV, 219 (1798).									
X. gracilis, Endlicher in Lehm. pl. Preiss. II, 39 (1846)... ...	W.A.	—	—	—	—	—	—	B.fl.VII,114	M.fr.IV,112.
X. pumilio, R. Brown, prodr. 288 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,114	
X. macronema, F. v. M., fragm. IV, 112 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,113	
X. hastilis, R. Brown, prodr. 288 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,115	M.fr.IV,113.
X. arborea, R. Brown, prodr. 288 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,115	M.fr.IV,113.
X. semiplana, F. v. M., fragm. IV, 111 (1864)	—	S.A.	—	—	—	—	—	B.fl.VII,115	M.fr.IV,111.
X. bracteata, R. Brown, prodr. 288 (1810)...	—	—	—	—	N.S.W.	—	—	B.fl.VII,116	
X. minor, R. Brown, prodr. 288 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,114	M.fr.IV,112.
X. australis, R. Brown, prodr. 288 (1810)	—	—	T.	V.	—	—	—	B.fl.VII,116	M.fr.IV,110.
X. quadrangulata, F. v. M., fragm. IV, 111 (1864)	—	S.A.	—	—	—	—	—	B.fl.VII,117	M.fr.IV,111.
X. Preissii, Endlicher in Lehm. pl. Preiss. II, 39 (1846)... ...	W.A.	—	—	—	—	—	—	B.fl.VII,117	M.fr.IV,110.
KINGIA, R. Brown in King's Narrat. of survey of Austr. coasts II, 529 (1827).									
K. australis, R. Brown in King's Narr. of surv. II, 535 (1827)...	W.A.	—	—	—	—	—	—	B.fl.VII,119	

PALMAE.

Ray, method. pl. emend. 135 (1703).

CALAMUS, Linné, spec. plant. 325 (1753).									
C. australis, Martius, hist. nat. palm. III, 213 (1839)	—	—	—	—	—	Q.	—	B.fl.VII,134	M.fr.V,48.
C. Muelleri, Wendland & Drude in Linnaea XXXIX, 193 (1871)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,135	
C. caryotoides, A. Cunn. in Martius, hist. nat. palm. III, 212 (1839)	—	—	—	—	—	Q.	—	B.fl.VII,135	

BACULARIA, F. v. M., fragm. VII, 103 (1870). (Linospadix).
B. monostachya, F. v. M., fragm. VII, 103 (1870) ... ... — — — — N.S.W. Q. — B.fl.VII,136 M.fr.VII,82,103;XI,58.

KENTIA, Blume in Bull. Neerl. 64 (1838). (Grisebachia, Hydriastele, Hedyscepe, Kentiopsis, Veitchia partly).
K. minor, F. v. M., fragm. VIII, 235 (1874) ... ... — — — — — Q. — B.fl.VII,137 M.fr.IX,195;XI,58,137.
K. Wendlandiana, F. v. M., fragm. VII, 102 (1870) ... ... — — — — — Q. N.A. B.fl.VII,138 M.fr.VII,102.
K. Belmoreana, C. Moore & F. v. M., fragm. VII, 99 (1870) ... — — — — N.S.W. — — B.fl.VII,137 M.fr.VIII,234.
K. Forsteriana, C. Moore & F. v. M., fragm. VII, 100 (1870) ... — — — — N.S.W. — — —
K. Baueri, Seemann, fl. Vitiens. 269 (1869) ... ... — — — — N.S.W. — — — M.fr.VII,102;VIII,233.
K. Canterburyana, C. Moore & F. v. M., fragm. VII, 101 (1870) — — — — N.S.W. — — B.fl.VII,138 M.fr.VIII,234;X,119.
K. acuminata, Wendland & Drude in Linnea XXXIX, 207 (1871) — — — — — — N.A. B.fl.VII,138

CLINOSTIGMA, Herm. Wendland in Seemann's Bonplandia X, 196 (1862). (Cyphokentia). [120.
C. Mooreanum, F. v. M., fragm. VIII, 235 (1874) ... ... — — — — N.S.W. — — B.fl.VII,139 M.fr.VII,101;IX,195;X,

PTYCHOSPERMA, Labillardière in Mém. de l'Inst., ann. 1808, I, 251, pl. X (1809). (Seaforthia, Laccospadix, Archontophoenix).
P. Laccospadix, Bentham, fl. Austr. VII, 140 (1878) ... ... — — — — — Q. — B.fl.VII,140
P. Beatricae, F. v. M. in Melb. Chemist. Febr. (1882) ... ... — — — — — Q. — —
P. Alexandrae, F. v. M., fragm. V, 47, t. 43 (1865) ... ... — — — — — Q. — B.fl.VII,140 M.fr.V,47,190,213.
P. Cunninghamii, Herm. Wendland in Bot. Zeit. 346 (1858) ... — — — — N.S.W. Q. — B.fl.VII,141 M.fr.VIII,222.
P. elegans, Blume, Rumphia II, 118 (1836) ... ... — — — — — Q. — B.fl.VII,141 M.fr.VIII,222.
P. Normanbyi, F. v. M., fragm. XI, 56 (1878) ... ... — — — — — Q. — B.fl.VII,142 M.fr.VIII,235.

ARECA, Linné, spec. plant. 1189 (1753) from Ray (1688).
A. Alicae, F. v. M. in Regel's Gartenflora, 199-201 (1879) ... — — — — — Q. — — M.fr.XI,88.

COCOS, Linné, spec. plant. 1188 (1753).
C. nucifera, Linné, spec. plant. 1188 (1753) ... ... — — — — — Q. — B.fl.VII,143

CARYOTA, Linné, gen. plant. 355 (1737).
C. Rumphiana, Martius, hist. nat. palm. III, 195 (1839) ... — — — — — Q. — B.fl.VII,144

LICUALA, Rumphius, herb. Amboin. I, 44, t. 9 (1741).
L. Muelleri, Wendland & Drude in Linnaea XXXIX, 223 (1871) — — — — — Q. — B.fl.VII,145 M.fr.VIII,221;X,120.

LIVISTONA, R. Brown, prodr. fl. Nov. Holl. 267 (1810). (Corypha partly).
L. humilis, R. Brown, prodr. 268 (1810) ... ... — — — — — — N.A. B.fl.VII,146 M.fr.VIII,221.
L. inermis, R. Brown, prodr. 268 (1810) ... ... — — — — — — N.A. B.fl.VII,146
L. Mariae, F.v.M. in Giles's Geog. Trav. in Cntr. Austr.222(1875) — S.A. — — — — N.A.
L. australis, Martius, hist. nat. palm. III, 241 (1839) ... ... — — — — — — N.A. — M.fr.XI,54.

PANDANEAE.

R. Brown, prodr. Nov. Holl. (1810)

PANDANUS, Rumphius, herb. Amboin. IV, 139, t. 74 (1750) (Athrodactylis, Fiquetia).
P. odoratissimus, Linné, fil. suppl. 424 (1781) ... ... — — — — — — N.A. B.fl.VII,148 M.fr.XI,55.
P. pedunculatus, R. Brown, prodr. 341 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,149 M.fr.VIII,220;IX,193.
P. Forsteri, Moore & F. v. M., fragm. VIII, 220 (1874) ... — — — — N.S.W. — — B.fl.VII,149 M.fr.VIII,220.
P. aquaticus, F. v. M., fragm. V, 40 (1865) ... ... — — — — — Q. N.A. B.fl.VII,148 M.fr.VIII,220. [193
P. monticola, F. v. M., fragm. V, 40 (1865) ... ... — — — — — Q. — B.fl.VII,150 M.fr.VII,63;VIII,220;IX,

FREYCINETIA, Gaudichaud in Annal. des scienc. natur. III, 509 (1824).
F. Gaudichaudii, Brown & Bennett, pl. Jav. rar. 31, t. 9 (1838) — — — — — Q. — B.fl.VII,151
F. excelsa, F. v. M., fragm. V, 39 (1865) ... ... — — — — — Q. — B.fl.VII,151 M.fr.V,39.
F. Baueriana, Endlicher, prodr. fl. ins. Norf. 25 (1833) ... ... — — — — N.S.W. — — —

NIPA, Wurmb in Verhandl. Batav. Genootsch. I, 349 (1779), from Camellus, Ray & Rumphius.
N. fruticans, Wurmb in Verhandl. Batav. Genootsch. I, 349 (1779) — — — — — Q. — — M.fr.XI,128.

AROIDEAE.

A. L. de Jussieu, gen. plant. 23 (1789), from B. de Jussieu (1759).

TYPHONIUM, Schott in Wien. Zeitschr. III, 72 (1829). (Arum partly.)
T. liliifolium, F. v. M. in Schott, prodr. syst. aroid. 107 (1860) — — — — — — N.A. B.fl.VII,153 M.fr.VIII,187.
T. alismifolium, F. v. M., fragm. VIII, 186 (1874) ... ... — — — — — Q. — B.fl.VII,153 M.fr.VIII,186.
T. Brownii, Schott, icon. aroid. II, t. 15 (1857) ... . — — — — N.S.W. Q. — B.fl.VII,154 M.fr.VIII,187.
T. angustilobum, F. v. M., fragm. X, 66 (1876) ... ... — — — — — Q. — B.fl.VII,154 M.fr.X,66.

AMORPHOPHALLUS, Blume in Nouv. Ann. du Mus. III, 366 (1834). (Brachyspatha).
A. variabilis, Blume, Rumphia I, 146, t. 35 (1835) ... ... — — — — — — N.A. B.fl.VII,155 M.fr.VIII,87.

COLOCASIA, Ray, Method. 157 (1682). (Caladium partly.)
C. antiquorum, Schott, Meletemat. botan. 18 (1832) ... ... — — — — — Q. — B.fl.VII,155 M.fr.VIII,87.
C. macrorrhiza, Schott, Meletemat. bot. 18 (1832) ... ... — — — — N.S.W. Q. — B.fl.VII,155 M.fr.VIII,87.

RHAPHIDOPHORA, Hasskarl in Fl. Regensb. Beibl. II, 11 (1842). (Epipremnum partly.)
R. pertusa, Schott in Seem. Bonplandia V, 45 (1857) ... ... — — — — — Q. — B.fl.VII,156 M.fr.VIII,87.

GYMNOSTACHYS, R. Brown, prodr. fl. Nov. Holl. 337 (1810).
G. anceps, R. Brown, prodr. 337 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,157 M.fr.VIII,87.

POTHOS, Linné, amœn. acad. I, 410 (1747).
P. Loureiri, Hooker & Arnott, Bot. of Beech. Voy. 220 (1841)... — — — — N.S.W. Q. — B.fl.VII,158 M.fr.I,62;VIII,87.

TYPHACEAE.

A. L. de Jussieu, gener. plant. 25 (1789).

TYPHA, Tournefort, inst. 530, t. 301 (1700), from Theophrastos, Dioscorides and Plinius.
T. angustifolia, Linné, spec. plant. 971 (1753) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,159 M.fr.VII,116.

SPARGANIUM, Tournefort, inst. 530, t. 302 (1700).
S. angustifolium, R. Brown, prodr. 338 (1810) ... ... — — — V. N.S.W. Q. — B.fl.VII,160 M.fr.VII,117.

LEMNACEAE.

J. E. Gray, Nat. arrang. of Brit. pl. II, 729 (1821).

LEMNA, Linné, syst. nat. 9 (1735); Linné, Fl. Lappon. 351 (1737). (Telmatophace, Spirodela.)

L. trisulca, Linné, spec. plant. 970 (1753)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,162	M.fr.VIII,188.
L. minor, Linné, spec. plant. 970 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,163	M.fr.VIII,188.
L. oligorrhiza, Kurz in Journ. Linn. Soc. IX, 267 (1867) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,163	M.fr.VIII,188.
L. polyrrhiza, Linné, spec. plant. 970 (1753)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,164	M.fr.VIII,188.
L. gibba, Linné, spec. plant. 970 (1753)	W.A.	—	—	—	—	—	—	B.fl.VII,163	M.fr.VIII,188.

WOLFFIA, Horkel & Schleiden in Schlecht. Linnaea XIII, 389 (1839). (Wolfia.)

W. Michelii, Schleiden, Beitr. zur Botan. 233 (1844)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,162	M.fr.VIII,187.

FLUVIALES.

Ventenat, Tabl. du règn. véget. II, 80, (1799).

TRIGLOCHIN, Rivinus in Ruppius, fl. Jenens. 54 et 366 (1718), from Dalechamps (1586). (Cycnogeton.)

T. centrocarpa, Hooker, icon. plant. t. 728 (1848)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,167	M.fr.VI,82.
T. striata, Ruiz & Pavon, Fl. Per. et Chil. III, 72 (1802) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,166	M.fr.VI,83.
T. mucronata, R. Brown, prodr. 343 (1810)	W.A.	S.A.	—	V.	—	—	—	B.fl.VII,168	M.fr.VI,81.
T. procera, R. Brown, prodr. 343 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,168	M.fr.VI,83.
T. Maundii, F. v. M., fragm. VI, 83 (1867)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,169	

APONOGETON, Linné fil., suppl. plant. 32 (1781).

A. monostachyus, Linné fil., suppl. 214 (1781)	—	—	—	—	—	Q.	N.A.	B.fl.VII,188	M.fr.VIII,215.
A. elongatus, F. v. M. in Benth. Fl. Austr. VII, 188 (1878) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,188	M.fr.VIII,216.

POTAMOGETON, Fuchs, histor. stirp. comment. 651 (1542).

P. natans, Linné, spec. plant. 126 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,170	M.fr.VIII,217.
P. Javanicus, Hasskarl in Act. Soc. Ind. Neerl. I, 26 (1857) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,171	M.fr.VIII,217.
P. Drummondii, Bentham, Fl. Austr. VII, 171 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,171	
P. perfoliatus, Linné, spec. plant. 126 (1753)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,172	M.fr.VIII,217.
P. crispus, Linné, spec. plant. 126 (1753)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,172	M.fr.VIII,217.
P. obtusifolius, Mertens & Koch, Deutschl. Fl. I, 855 (1823) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,172	M.fr.VIII,216.
P. acutifolius, Link in Roem. & Schult. syst. veg. III, 513 (1818)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,173	M.fr.VIII,216.
P. pectinatus, Linné, spec. plant. 127 (1753)	—	S.A.	T.	V.	—	—	—	B.fl.VII,173	M.fr.VIII,217.

RUPPIA, Linné, syst. nat. 9 (1735); Linné, gen. plant. 277 (1737).

R. maritima, Linné, spec. plant. 127 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,174	M.fr.VIII,217

POSIDONIA, Koenig, Annals of Botany II, 95, t. 6 (1806). (Caulinia partly, Kernera partly.)

P. australis, J. Hooker, Fl. Tasman. II, 43 (1860)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VII,175	M.fr.VIII,218.

ZOSTERA, Linné, Waesgoeta-Resa, 166-168 et fig. (1747).

Z. nana, F. K. Mertens in Roth, enum. pl. phaner. Germ. 8 (1827)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,176	M.fr.VIII,218.
Z. Tasmanica, G. v. Mertens in Garcke, Linnaea XXXV, 68 (1867)	—	S.A.	T.	V.	—	—	—	B.fl.VII,176	M.fr.VIII,218.

CYMODOCEA, C. Koenig, Annals of Botany II, 96, t. 7 (1806). (Amphibolis, Graumuellera, Kernera partly.)

C. zosterifolia, F.v.M.; Amphibolis, Agardh., spec. algar. II, 477 (1822)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VII,177	M.fr.VIII,218.
C. ciliata, Ehrenberg in Garcke, Linnaea XXXV, 162 (1867) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,178	M.fr.VIII,218.
C. rotundata, Ascherson & Schweinfurth in Neumayer, Anleit. zu wissensch. Beob. 361 (1875)	—	—	—	—	—	—	N.A.	—	M.fr.X,162.
C. serrulata, Ascherson & Magnus in Neumayer, Anleit. zu wiss. Beob. 362 (1875)	—	—	—	—	—	Q.	—	B.fl.VII,178	M.fr.VIII,218.
C. isoetifolia, Ascherson in Garcke, Linnaea XXXV, 163 (1867)	W.A.	—	—	—	—	—	N.A.	B.fl.VII,178	

HALODULE, Endlicher, gen. plant. 1368 (1840). (Diplanthera, 1806.)

H. tridentata, Endlicher, gen. plant. 1368, implied (1840) ...	—	—	—	—	—	—	N.A.	—	M.fr.VIII,218.

LEPILAENA, Drummond & Harvey in Hook. Kew misc. VII, 57 (1855). (Zannichellia partly, Hexatheca, Altheniae subgenus.)

L. australis, J. Drummond in Hook. Kew misc. VII, 58 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.VII,179	M.fr.VIII,217.
L. Preissii, F. v. M., fragm. VIII, 217 (1874)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VII,180	M.fr.VIII,217.

NAJAS, Linné, syst. nat. 9 (1735); Linné, gen. plant. 278 (1737). (Caulinia partly.)

N. major, Allioni, Fl. Pedem. II, 221 (1785)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.VII,181	M.fr.VIII,218.
N. tenuifolia, R. Brown, prodr. 345 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,181	M.fr.VIII,219.

ALISMACEAE.

Ventenat, Tabl. du règn. véget. II, 157 (1799).

ALISMA, Rivinus in Ruppius, fl. Jen. 54 (1718), from Dioscorides and Plinius. (Caldesia; Echinodorus partly).

A. Plantago, Linné, spec. plant. 842 (1753)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,185	M.fr.VIII,215.
A. acanthocarpum, F. v. M., fragm. I, 23 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.VII,185	M.fr.VIII,214.
A. oligococcum, F. v. M., fragm. I, 23 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.VII,185	M.fr.VI,169; VIII,214.
A. parnassifolium, Bassi in act. Bonon. (1768)	—	—	—	—	—	Q.	—	B.fl.VII,186	M.fr.VIII,214.

DAMASONIUM, Tournefort, inst. 256 (1700). (Actinocarpus.)

D. australe, Salisbury in Transact. hort. Soc. Lond. I, 268 (1812)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,186	M.fr.VIII,215.

TENAGOCHARIS, Hochstetter in Regensb. bot. Zeit. June (1841). (Butomopsis.)

T. Cordofana, Hochstetter in Regensb. bot. Zeit. June (1841) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,187	M.fr.X,103.

PONTEDERIACEAE.

Humboldt, Bonpland & Kunth, nov. gen. et spec. plant. I, 265 (1815).

MONOCHORIA, Presl, reliq. Haenk. I, 127 (1827). (Limnostachys.)

M. cyanea, F. v. M., fragm. VIII, 44 (1872)	—	—	—	—	—	Q.	N.A.	B.fl.VII,74	M.fr.VIII,44.

PHILYDREAE.

R. Brown in Flinders' voy. II, 57 (1814).

PHILYDRUM, Banks in Gaertner, de fruct. I, 62 t. 16 (1788). (Philhydrum.)

P. lanuginosum, Banks in Gaertn. de fruct. I. 62 t. 16 (1788) ...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,73	M.fr.V,203.

PRITZELIA, F. v. M., Papuan plants 13 (1875). (Hetaeria, Philydrella.)

P. pygmaea, F. v. M., Papuan plants 13 (1875)	W.A.	—	—	—	—	—	—	B.fl.VII,74	M.fr.V,203.

HELMHOLTZIA, F. v. M., fragm. V, 202 (1866).

H. acorifolia, F. v. M., fragm. V, 203 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,75	M.fr.V,203.

COMMELINEAE.

R. Brown, prodr. fl. Nov. Holl. 268 (1810).

ZYGOMENES, Salisbury in Transact. hort. Soc. I, 271 (1812). (Cyanotis).

Z. axillaris, Salisbury in Transact. hort. Soc. I, 271 (1812) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,82	M.fr.VIII,62.

COMMELINA, Plumier, nov. pl. Amer. gen. 48, t. 38 (1703). (Commelynia).

C. ensifolia, R. Brown, prodr. 269 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,83	M.fr.VIII,60;IX,191.
C. cyanea, R. Brown, prodr. 269 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,84	M.fr.VIII,59;IX,191.
C. lanceolata, R. Brown, prodr. 269 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,84	M.fr.VIII,59.
C. agrostophylla, F.v. M. in J. Hook., Fl. Tasm. p. XLVII (1860)	—	—	—	—	-	Q.	N.A.	—	M.fr.VIII,59.

ANEILEMA, R. Brown, prodr. fl. Nov. Holl. 270 (1810).

A. acuminatum, R. Brown, prodr. 270 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,85	M.fr.VIII,61.
A. biflorum, R. Brown, prodr. 270 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,86	
A. sclerocarpum, F. v. M., fragm. VIII, 61 (1873)	—	—	-	—	—	Q.	—	B.fl.VII,86	M.fr.VIII,61.
A. siliculosum, R. Brown, prodr. 270 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,87	M.fr.VIII,61;IX,191.
A. gramineum, R. Brown, prodr. 270 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,87	M.fr.VIII,62.
A. giganteum, R. Brown, prodr. 271 (1810)	—	--	—	—	—	Q.	N.A.	B.fl.VII,88	M.fr.VIII,62.

FLORISCOPA, Loureiro, Fl. Cochinch. I, 192 (1790). (Floscopa, Dithyrocarpus.)

F. scandens, Loureiro, fl. Cochinch. I, 193 (1790)	—	—	—	—	—	Q.	—	B.fl.VII,89	M.fr.VIII,63.

POLLIA, Thunberg, nov. gen. plant. I, 11 (1781).

P. cyanococca, F. v. M., fragm. V, 40 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,90	M.fr.VIII,63.

CARTONEMA, R. Brown, prodr. fl. Nov. Holl. 271 (1810).

C. philydroides, F. v. M., fragm. I, 62 (1858)	W.A.	—	—	—	-	—	—	B.fl.VII,91	M.fr.VIII,64.
C. spicatum, R. Brown, prodr. 271 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,91	M.fr.I,62;VIII,64.
C. parviflorum, Hasskarl in Flora Regensb. 365 (1869) ...	—	—	—	-	—	—	N.A.	B.fl.VII,91	
C. trigonospermum, Clarke in De Cand., mon. phaner. III, 24 (1881)	—	—	—	—	—	—	N.A.	—	M.fr.XI,139.
C. brachyantherum, Bentham, Fl. Austr. VII, 92 (1878) ...	--	—	—	—	—	Q.	—	B.fl.VII,92	
C. tenue, Bentham in De Cand., monogr. phaner. III, 264 (1881)	—	—	—	—	—	—	N.A.	—	M.fr.XI,139.

XYRIDEAE.

Salisbury in Transact. of the Hortic. Soc. I, 326 (1812).

XYRIS, Gronovius in Linné, genera plantar. 11 (1737).

X. complanata, R. Brown, prodr. 256 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,77	M.fr.VIII,205.
X. pauciflora, Willdenow, phytogr. 2, t. 1 (1794)...	—	—	—	—	—	Q.	N.A.	B.fl.VII,77	M.fr.VIII,205.
X. lacera, R. Brown, prodr. 257 (1810)	W.A.	—	—	-	—	—	—	B.fl.VII,78	M.fr.VIII,204.
X. flexifolia, R. Brown, prodr. 256 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,78	M.fr.VIII,204.
X. gracilis, R. Brown, prodr. 256 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,79	M.fr.VIII,204.
X. operculata, Labillardière, Nov. Holl. pl. spec. I, 14 t. 10 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,79	M.fr.VIII,204.
X. lanata, R. Brown, prodr. 257 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,80	M.fr.VIII,204.
X. laxiflora, F. v. M., fragm. VIII, 203 (1874)	W.A.	—	-	-	—	—	-	B.fl.VII,80	M.fr.VIII,203.
X. gracillima, F. v. M., fragm. VIII, 203 (1874)	W.A.	—	—	—	—	-	—	B.fl.VII,80	M.fr.VIII,203.

JUNCEAE.

R. Brown, prodr. fl. Nov. Holl. 257 (1810).

LUZULA, De Candolle, Fl. franc. III, 158 (1805).

L. campestris, De Candolle, Fl. franc. III, 158 (1805)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,123	
L. longiflora, Bentham, Fl. Austr. VII, 123 (1878)	—	—	—	—	N.S.W.	—	—	B.fl.VII,123	

JUNCUS, Tournefort, inst. 246, t. 127 (1700), from Camerarius, J. & C. Bauhin, Ray & Morison.

J. gracilis, R. Brown, prodr. 259 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,125	
J. planifolius, R. Brown, prodr. 259 (1810)	W.A.	S.A.	T.	V.	N.S.W.	-	—	B.fl.VII,125	
J. caespititius, E. Meyer in Lehm. pl. Preiss. II, 47 (1846) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,126	
J. falcatus, E. Meyer, synops. luzular. 34 (1823)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,126	
J. bufonius, Linné, spec. plant. 328 (1753)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,127	
J. homalocaulis, F. v. M., first gen. Rep. 19 (1853)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,128	
J. Brownii, F. v. M., first gen. Rep. 19 (1853)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,128	M.fr.IX,78.
J. communis, E. Meyer, synops. Juncor. 12 (1822)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,128	
J. vaginatus, R. Brown, prodr. 258 (1810)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,129	
J. pauciflorus, R. Brown, prodr. 259 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,129	
J. pallidus, R. Brown, prodr. 258 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,130	
J. maritimus, Lamarck, Encycl. méth. III, 264 (1789)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,130	
J. prismatocarpus, R. Brown, prodr. 259 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	--	B.fl.VII,131	
J. pusillus, Buchenau in Abhandl. naturw. Vereins zu Bremen VI, 395 (1879)	—	—	T.	V.	N.S.W.	-	—	B.fl.VII,132	

ERIOCAULEAE.

Humboldt, Bonpland & Kunth, nov. gen. et spec. plant. I, 251 (1815).

ERIOCAULON, Linné, gen. plant. ed. sec. 35 (1742). (Electrosperma.)

E. setaceum, Linné, spec. plant. 87 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VII,191							
E. australe, R. Brown, prodr. 254 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,192							
E. quinquangulare, Linné, spec. plant. 87 (1753)	—	—	—	—	—	—	N.A.	B.fl.VII,192							
E. Smithii, R. Brown, prodr. 254 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,192							
E. nanum, R. Brown, prodr. 254 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,193							
E. cinereum, R. Brown, prodr. 254 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,193	M.fr.I,95.						
E. pusillum, R. Brown, prodr. 254 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,194							
E. pallidum, R. Brown, prodr. 254 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,194							
E. nigricans, R. Brown, prodr. 254 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,194							
E. electrospermum; Electrosperma Australasicum, F. v. M. in Trans. phil. Soc. Vict. I, 24 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,195							
E. lividum, F. v. M., fragm. I, 92 (1858)	—	—	—	—	—	—	N.A.	B.fl.VII,195	M.fr.I,92.						
E. concretum, F. v. M., fragm. I, 92 (1858)	—	—	—	—	—	—	N.A.	B.fl.VII,195	M.fr.I,92.						
E. Schultzii, Bentham, Fl. Austr. VII, 195 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,195							
E. tortuosum, F. v. M., fragm. I, 94 (1858)	—	—	—	—	—	—	N.A.	B.fl.VII,196	M.fr.I,91.						
E. monoscapum, F. v. M., fragm. I, 94 (1858)	—	—	—	—	—	—	N.A.	B.fl.VII,196	M.fr.I,94.						
E. spectabile, F. v. M., fragm. I, 95 (1858)	—	—	—	—	—	—	N.A.	B.fl.VII,196	M.fr.I,95.						
E. scariosum, R. Brown, prodr. 255 (1810)...	—	—	—	—	—	Q.	N.A.	B.fl.VII,197							
E. depressum, R. Brown, prodr. 255 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,197	M.fr.I,92.						

RESTIACEAE.

R. Brown, prodr. flor. Nov. Holl. 243 (1810).

TRITHURIA, J. Hooker, Fl. Tasman. II, 79, t. 138 (1860). (Juncella, 1854.)

T. submersa, J. Hooker, Fl. Tasman. II, 79, t. 138 (1860) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,199	M.fr.VIII,237.						

APHELIA, R. Brown, prodr. fl. Nov. Holl. 251 (1810). (Brizula.)

A. cyperoides, R. Brown, prodr. 252 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,200	M.fr.VIII,236.						
A. nutans, J. Hooker in Benth. Fl. Austr. VII, 200 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,200							
A. gracilis, Sonder in Schlecht. Linnaea XXVIII, 227 (1855) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,201	M.fr.VIII,237.						
A. Drummondii, Bentham, Fl. Austr. VII, 201 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,201							
A. Pumilio, F. v. M., Schlecht. in Linnaea XXVIII, 226 (1855)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,201	M.fr.VIII,237.						
A. Brizula, F. v. M., fragm. V, 203 (1866)...	W.A.	—	—	—	—	—	—	B.fl.VII,202	M.fr.VIII,237.						

CENTROLEPIS, Labillardière, Nov. Holl. pl. specim. I, 7 (1804). (Desvauxia, Alepyrum.)

C. humillima, F. v. M. in Benth. Fl. Austr. VII, 203 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,203							
C. polygyna, Hieronymus in Abh. naturf. Ges. Halle XII, 96 (1873)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,203	M.fr.VIII,237.						
C. alepyroides, Hieronymus in Abh. nat. Ges. Halle XII, 96 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,204	M.fr.VIII,237.						
C. mutica, Hieronymus in Abh. nat. Ges. Halle XII, 97 (1873)...	W.A.	—	—	—	—	—	—	B.fl.VII,204							
C. glabra, F. v. M. in Abh. nat. Ges. Halle XII, 95 (1873) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,204	M.fr.VIII,237.						
C. muscoides, Hieronymus in Abh. nat. Ges. Halle XII, 95 (1873)	—	—	T.	—	—	—	—	B.fl.VII,205							
C. monogyna, Bentham, Fl. Austr. VII, 205 (1878)	—	—	T.	—	—	—	—	B.fl.VII,205							
C. pulvinata, Desvaux in Ann. des sc. nat. XIII, 42, t. 2 (1828)	—	—	T.	—	—	—	—	B.fl.VII,205							
C. pusilla, Roemer & Schultes, syst. veg. I, 44 (1817)	—	—	—	—	—	Q.	—	B.fl.VII,205							
C. aristata, Roemer & Schultes, syst. veg. I, 44 (1817)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,206	M.fr.VIII,237.						
C. Drummondii, Hieronymus in Abh. nat. Ges. Halle XII, 98 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,206	M.fr.VIII,237.						
C. Banksii, Roemer & Schultes, syst. veg. I, 44 (1817)	—	—	—	—	—	Q.	N.A.	B.fl.VII,207	M.fr.VIII,237.						
C. fascicularis, Labillardière, Nov. Holl. pl. spec. I, 7, t. 1 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,207							
C. pilosa, Hieronymus in Abh. nat. Ges. Halle 102 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VII,207							
C. strigosa, Roemer & Schultes, syst. veg. I, 43 (1817)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,207							
C. exserta, Roemer & Schultes, syst. veg. I, 44 (1817)	—	—	—	—	—	Q.	N.A.	B.fl.VII,208	M.fr.VIII,237.						

LYGINIA, R. Brown, prodr. fl. Nov. Holl. 248 (1810). (Schoenodum partly.)

L. barbata, R. Brown, prodr. 248 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,210	M.fr.VIII,79.						

ECDEIOCOLEA, F. v. M., fragm. phytogr. Austr. VIII, 236 (1874).

E. monostachya, F. v. M., fragm. VIII, 236 (1874)	W.A.	—	—	—	—	—	-	B.fl.VII,211	M.fr.VIII,236.						

ANARTHRIA, R. Brown, prodr. fl. Nov. Holl. 248 (1810).

A. scabra, R. Brown, prodr. 249 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,212	M.fr.VIII,80.						
A. laevis, R. Brown, prodr. 249 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,212	M.fr.VIII,81.						
A. gracilis, R. Brown, prodr. 249 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,213	M.fr.VIII,72,81.						
A. prolifera, R. Brown, prodr. 249 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,213	M.fr.VIII,82.						
A. polyphylla, Nees in Lehm. pl. Preiss. II, 63 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,214	M.fr.VIII,83.						

LEPYRODIA, R. Brown, prodr. fl. Nov. Holl. 247 (1810).

L. scariosa, R. Brown, prodr. 248 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VII,215	M.fr.VIII,72.						
L. Muelleri, Bentham, Fl. Austr. VII, 215 (1878)...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,215							
L. anarthria, F. v. M., fragm. VIII, 73 (1873)	—	—	—	—	N.S.W.	—	—	B.fl.VII,216							
L. gracilis, R. Brown, prodr. 247 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VII,216	M.fr.VIII,73.						
L. Tasmanica, J. Hooker, Fl. Tasman. II, 72, t. 135 (1860) ...	—	—	T.	V.	—	—	—	B.fl.VII,216	M.fr.VIII,73,75;IX,194.						
L. interrupta, F. v. M., fragm. VIII, 74 (1873)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,217	M.fr.VIII,74.						
L. hermaphrodita, R. Brown, prodr. 248 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,217	M.fr.VIII,75.						
L. monoica, F. v. M., fragm. VIII, 76 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,217	M.fr.VIII,76.						
L. Muirii, F. v. M., fragm. VIII, 78 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,218	M.fr.VIII,76.						
L. stricta, R. Brown, prodr. 248 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,218	M.fr.VIII,74.						
L. macra, Nees in Lehm. pl. Preiss. II, 60 (1846)...	W.A.	—	—	—	—	—	—	B.fl.VII,218							
L. Drummondiana, Steudel, syn. pl. glum. II, 248 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.VII,219	M.fr.VIII,75.						
L. glauca, F. v. M., fragm. VIII, 77 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,219	M.fr.VIII,77.						
L. anaectocolea, F. v. M., fragm. VIII, 78 (1873)...	W.A.	—	—	—	—	—	—	B.fl.VII,220	M.fr.VIII,78.						

RESTIO, Linné, syst. nat. edit. duo dec. II, 735 (1767). (Megalotheca.)

R. fastigiatus, R. Brown, prodr. 246 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VII,222	M.fr.VIII,67.
R. Megalotheca, F. v. M., fragm. VIII, 99 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,222	
R. tropicus, R. Brown, prodr. 246 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,223	
R. applanatus, Sprengel, syst. I, 185 (1825)	W.A.	—	—	—	—	—	—	B.fl.VII,223	M.fr.VIII,69.
R. monocephalus, R. Brown, prodr. 245 (1810)	—	—	T.	—	—	—	—	B.fl.VII,224	M.fr.VIII,68.
R. dimorphus, R. Brown, prodr. 246 (1810)	—	—	—	—	N.S.W.	Q.	-	B.fl.VII,224	M.fr.VIII,68.
R. conferto-spicatus, Steudel, syn. glum. II, 256 (1855)	W.A.	—	—	—	—	—	—	B.fl.VII,224	
R. sphacelatus, R. Brown, prodr. 246 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,225	
R. deformis, R. Brown, prodr. 245 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,225	M.fr.
R. crispatus, R. Brown, prodr. 246 (1810)	W.A.	—	—	—	—	—	-	B.fl.VII,225	
R. nitens, Nees in Lehm. pl. Preiss. II, 59 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,226	M.fr.VIII,70.
R. gracilior, F. v. M. in Benth. Fl. Austr. VII, 226 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,226	
R. chaunocoleus, F. v. M., fragm. VIII, 64 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,226	M.fr.
R. australis, R. Brown, prodr. 245 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,227	M.fr.VIII,69.
R. gracilis, R. Brown, prodr. 245 (1810)	—	—	T.	V.	N.S.W.	Q.	-	B.fl.VII,227	M.fr.VIII,69.
R. complanatus, R. Brown, prodr. 245 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,228	M.fr.VIII,67.
R. tetraphyllus, Labillardière, Nov. Holl. pl. spec. II, 77 (1806)	—	S.A.	T.	V.	N.S.W.	Q.		B.fl.VII,228	M.fr.VIII,66.
R. laxus, R. Brown, prodr. 245 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,228	M.fr.VIII,71.
R. ornatus, Steudel, syn. pl. glum. II, 256 (1855)	W.A.	—	—	—	—	—	-	B.fl.VII,229	M.fr.VIII,70.
R. leptocarpoides, Bentham, Fl. Austr. VII, 229 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,229	M.fr.VIII,65.
R. amblycoleus, F. v. M., fragm. VIII, 65 (1873)	W.A.	—	—	—	—	—	-	B.fl.VII,230	M.fr.VIII,65.
R. tremulus, R. Brown, prodr. 230 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,230	M.fr.VIII,70.

LOXOCARYA, R. Brown, prodr. fl. Nov. Holl. 249 (1810). (Desmocladus.)

L. densa, Bentham, Fl. Austr. VII, 241 (1878)	W.A.	—	—	—	—	—	-	B.fl.VII,241	
L. Benthami, F. v. M.; Hypolaena, Masters in Cand. mon. phan. I, 367 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,242	
L. pubescens, Bentham, Fl. Austr. VII, 242 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,242	M.fr.VIII,100.
L. fasciculata, Bentham, Fl. Austr. VII, 242 (1878)	W.A.		—	—	—	—	—	B.fl.VII,242	M.fr.VIII,98.
L. flexuosa, Bentham, Fl. Austr. VII, 243 (1878)	W.A.	—	—	—	—	—	-	B.fl.VII,243	M.fr.VIII,100.
L. cinerea, R. Brown, prodr. 249 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,243	

CALOSTROPHUS, Labillardière, Nov. Holl. pl. spec. II, 78 (1806). (Calorophus, Hypolaena.)

C. elongatus, F. v. M., fragm. VIII, 86 (1873)	—	—	T.	—	—	—	—	B.fl.VII,238	M.fr.VIII,86;IX,94;X,120
C. lateriflorus, F. v. M., fragm. VIII, 87 (1873)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,238	M.fr.VIII,87.
C. gracillimus, F. v. M., fragm. VIII, 88 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,239	M.fr.VIII,88.
C. fastigiatus, F. v. M. in pap. R. Soc. Tasm. 117 (1878)	—	S.A.	T.	V.	N.S.W.	—	-	B.fl.VII,239	M.fr.VIII,84.
C. exsulcus, F. v. M.; Hypolaena, R. Brown, prodr. 251 (1810)	W.A.	—	—	—	—	-	—	B.fl.VII,240	M.fr.VIII,86.

LEPTOCARPUS, R. Brown, prodr. 250 (1810). (Schoenodum partly.)

L. scariosus, R. Brown, prodr. 250 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,232	M.fr.VIII,94.
L. tenax, R. Brown, prodr. 250 (1810)	W.A.	S.A.	T.	V.	N.S.W.	-	-	B.fl.VII,232	M.fr.VIII,96.
L. Brownii, J. Hooker, Fl. Tasman. II, 73, t. 136 (1860)	—	S.A.	T.	V.	N.S.W.		-	B.fl.VII,233	M.fr.VIII,91.
L. canus, Nees in Ann. of nat. hist. VI, 50 (1843)	W.A.	—	—	—	—		—	B.fl.VII,234	M.fr.VIII,95.
L. coangustatus, Nees in Lehm. pl. Preiss. II, 65 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,234	M.fr.VIII,94.
L. aristatus, R. Brown, prodr. 250 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,235	M.fr.VIII,90.
L. erianthus, Bentham, Fl. Austr. VII, 235 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,235	
L. ramosus, R. Brown, prodr. 250 (1810)	—	—	—	—	-	Q.	—	B.fl.VII,236	M.fr.VIII,92.
L. elatior, R. Brown, prodr. 250 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,236	
L. spathaceus, R. Brown, prodr. 250 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,236	M.fr.VIII,93.
L. Schultzii, Bentham, Fl. Austr. VII, 237 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,237	M.fr.VIII,93.

LEPIDOBOLUS, Nees in Lehm. pl. Preiss. II, 66 (1846).

L. drapetocoleus, F. v. M., fragm. VIII, 84 (1873)	—	S.A.	—	V.	—	—	—	B.fl.VII,244	M.fr.VIII,84.
L. Preissianus, Nees in Lehm. pl. Preiss. II, 66 (1846)	W.A.	—	—	—	—	—	-	B.fl.VII,245	M.fr.VIII,83.
L. chaetocephalus, F. v. M., fragm. VIII, 84 (1873)	W.A.	—	—	—	-	—	—	B.fl.VII,245	

CHAETANTHUS, R. Brown, prodr. 251 (1810). (Prionosepalum.)

C. leptocarpoides, R. Brown, prodr. 251 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,246	M.fr.VIII,97.

ONYCHOSEPALUM, Steudel, syn. pl. glum. II, 249 (1855).

O. laxiflorum, Steudel, syn. pl. glum. II, 249 (1855)	W.A.	—	—	—	—	—	—	B.fl.VII,246	M.fr.IX,51.

ACALYCEAE HYPOGYNEAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 48 (1880).

CYPERACEAE.

Haller, enum. stirp. Helvet. I, 234 (1742).

KYLLINGIA, Rottboell, descr. et icon. rar. et nov. plant. 13, t. 4 (1773). (Killingia.)

K. intermedia, R. Brown, prodr. 219 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,251	M.fr.VIII,27.
K. monocephala, Rottboell, descr. et icon. pl. 13, t. 4 (1773)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VII,251	M.fr.VIII,271;IX,54.
K. cylindrica, Nees in Wight's Contr. to Bot. of Ind. 91 (1834)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,252	M.fr.VIII,271.
K. triceps, Rottboell, descr. et icon. pl. 14, t. 4 (1773)	—	—	—	—	—	Q.	—	B.fl.VII,252	

CYPERUS, Tournefort, inst. 527, t. 299 (1700), from Hippocr., Theophr. and Plinius. (Pycreus, Mariscus, Anosporum, Sorostachys.)

C. pumilus, Linné, amoen. acad. IV, 302 (1759)	—	—	—	—	—	Q.	—	B.fl.VII,258	M.fr.VIII,267.
C. Eragrostis, Vahl, enum. plant. II, 322 (1806)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,258	M.fr.VIII,260;IX,52.
C. flavescens, Linné, spec. plant. 46 (1753)	—	—	—	—	N.S.W.	Q.	-	B.fl.VII,259	
C. globosus, Allioni, auctuar. ad fl. Pedem. 49 (1789)	—	—	—	V.	—	Q.	-	B.fl.VII,260	M.fr.VIII,260;IX,52.
C. unioloides, R. Brown, prodr. 216 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,260	M.fr.VIII,259;IX,52.

C. polystachyus, Rottboell, descr. et icon. pl. 39, t. 11 (1773) ... — — — — N.S.W. Q. N.A. B.fl.VII,261 M.fr.VIII,265,270;IX,53.
C. flavicomus, Cl. Richard in Michaux, fl. bor. Amer. I, 27 (1803) — — — — — — N.A. B.fl.VII,261 M.fr.VIII,265;IX,53.
C. pygmaeus, Rottboell, descr. et icon. pl. 20, t. 14 (1773) ... — — — V. N.S.W. Q. N.A. B.fl.VII,262 M.fr.VIII,268;IX,53.
C. cephalotes, Vahl, enum. II, 311 (1806) ... ... — — — — — Q. — B.fl.VII,263 M.fr.VIII,272;IX,54.
C. laevigatus, Linné, mantissa alt. 179 (1771) ... ... W.A. — — — — — — B.fl.VII,263 M.fr.VIII,266;IX,53.
C. platystylis, R. Brown, prodr. 241 (1810) ... ... — — — — N.S.W. — — B.fl.VII,264
C. alopecuroides, Rottboell, descr. et icon. pl. 38, t. 8 (1773) .. — — — — — Q. — B.fl.VII,264 M.fr.VIII,263;IX,52.
C. pulchellus, R. Brown, prodr. 213 (1810) ... ... — — — — — Q. N.A. B.fl.VII,265 M.fr.VIII,271.
C. tenellus, Linné, fil. suppl. 103 (1781) ... ... W.A. S.A. — — N.S.W. — — B.fl.VII,265 M.fr.VIII,261;IX,53.
C. gracilis, R. Brown, prodr. 213 (1810) ... ... — S.A. — V. N.S.W. Q. — B.fl.VII,265 M.fr.VIII,264;IX,53.
C. enervis, R. Brown, prodr. 213 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,266
C. debilis, R. Brown, prodr. 213 (1810) ... ... — — — — N.S.W. — — B.fl.VII,266
C. castaneus, Willdenow, spec. plant, I, 278 (1797) ... ... — — — — — Q. — B.fl.VII,267
C. cuspidatus, Humboldt, Bonpland & Kunth, nov. gen. et sp. pl. I, 204 (1815) ... ... — — — — — Q. — B.fl.VII,267
C. squarrosus, Linné, amoen. acad. IV, 303 (1759) ... ... — S.A. — — — Q. N.A. B.fl.VII,268 M.fr.VIII,262;IX,53.
C. difformis, Linné, amoen. acad. IV, 302 (1759)... ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,268 M.fr.VIII,262;IX,52.
C. tetraphyllus, R. Brown, prodr. 214 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,269 M.fr.VIII,264.
C. trinervis, R. Brown, prodr. 213 (1810) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,269 M.fr.VIII,267;IX,53.
C. Haspan, Linné, spec. plant. 45 (1753) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,270 M.fr.VIII,260;IX,52.
C. concinnus, R. Brown, prodr. 214 (1810)... ... . — — — V. N.S.W. Q. — B.fl.VII,271 M.fr.VIII,261.
C. filipes, Bentham, Fl. Austr. VII, 271 (1878) .. .. — — — — N.S.W. Q. — B.fl.VII,271
C. pedunculosus, F. v. M., fragm. VIII, 266 (1874) ... ... — — — — — Q. — B.fl.VII,272 M.fr.
C. vaginatus, R. Brown, prodr. 213 (1810)... ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,272 M.fr.VIII,261;IX,53.
C. Holoschoenus, R. Brown, prodr. 215 (1810) ... ... — S.A. — — — Q. N.A. B.fl.VII,273 M.fr.I,200; VIII,262.
C. dactylotes, Bentham, Fl. Austr. VII, 273 (1878) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,273
C. Gilesii, Bentham, Fl. Austr. VII, 274 (1878) ... ... — S.A. — — N.S.W. Q. — B.fl.VII,274
C. fulvus, R. Brown, prodr. 215 (1810) ... ... — S.A. — — N.S.W. Q. — B.fl.VII,274 M.fr.VIII,268.
C. carinatus, R. Brown, prodr. 216 (1810)... ... ... — — — — N.S.W. Q. N.A. B.fl.VII,274
C. alterniflorus, R. Brown, prodr. 216 (1810) ... ... W.A. S.A. — — — Q. — B.fl.VII,275
C. pilosus, Vahl, enum. II, 354 (1806) ... ... — — — — N.S.W. Q. — B.fl.VII,275 M.fr.VIII,260;IX,52.
C. ornatus, R. Brown, prodr. 217 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,276
C. Iria, Linné, spec. plant. 45 (1753) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,276 M.fr.VIII,266;IX,53.
C. eleusinoides, Wallich in Kunth, enum. II, 39 (1837) ... ... — — — — — Q. N.A. B.fl.VII,277 M.fr.VIII,264;IX,53.
C. distans, Linné, fil. suppl. 103 (1781) ... ... — — — — — Q. — B.fl.VII,277 M.fr.VIII,266;IX,53.
C. tegetiformis, Roxburgh, hort. Benghal. (1814)... ... ... — — — — — — N.A. B.fl.VII,278 M.fr.VIII,260;IX,52.
C. articulatus, Linné, spec. plant. 44 (1753) ... ... — — — — — — N.A. B.fl.VII,278
C. diphyllus, Retzius, observ. bot. V, 11 (1789) ... ... — S.A. — — — — N.A. B.fl.VII,279
C. rotundus, Linné, spec. plant. 45 (1753)... ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,279 M.fr.VIII.269;IX,53.
C. stenostachyus, Bentham, Fl. Austr. VII, 280 (1878) ... ... W.A. — — — — — — B.fl.VII,280
C. congestus, Vahl, enum. II, 350 (1806) ... ... W.A. — — — N.S.W. — — B.fl.VII,280 M.fr.VIII,269;IX,53.
C. subulatus, R. Brown, prodr. 217 (1810)... ... ... — S.A. — — N.S.W. — — B.fl.VII,281
C. sporobolus, R. Brown, prodr. 215 (1810) ... ... — — — — — — N.A. B.fl.VII,281
C. angustatus, R. Brown, prodr. 214 (1810) ... ... — — — — — Q. N.A. B.fl.VII,282 M.fr.IX,54.
C. inornatus, Boeckeler in Fl. Regensb. 86 (1875) ... ... — — — — — Q. — B.fl.VII,282 M.fr.IX,54.
C. lucidus, R. Brown, prodr. 218 (1810) ... ... — S.A. T. V. N.S.W. Q. N.A. B.fl.VII,283 M.fr.VIII,270;IX,53.
C. pennatus, Lamarck, illustr. des genr. I, 144 (1791) ... ... — — — — — Q. N.A. B.fl.VII,284 M.fr.VIII,263;IX,53.
C. exaltatus, Retzius, observ. bot. V, 11 (1789) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,285 M.fr.VIII,263;IX,53.
C. haematodes, Endlicher, prodr. fl. Norf. 22 (1839) ... ... — — — — N.S.W. — — B.fl.VII,285 M.fr.IX,54.
C. auricomus, Sieber in Sprengel, syst. I, 230 (1825) ... ... — — — — — Q. — B.fl.VII,286 M.fr.VIII,263.
C. ferax, Cl. Richard in act. soc. hist. nat. Par. I, 106 (1792) ... — — — — — Q. — B.fl.VII,286
C. Bowmanni, F. v. M. in Benth. Fl. Austr. VII, 287 (1878) ... — — — — N.S.W. Q. — B.fl.VII,287
C. trichostachys, Bentham, Fl. Austr. VII, 287 (1878) ... ... — — — — — Q. — B.fl.VII,287
C. leiocaulis, Bentham, Fl. Austr. VII, 287 (1878) ... ... — — — — N.S.W. Q. — B.fl.VII,287
C. scaber, Bentham, Fl. Austr. VII, 288 (1878) ... ... — — — — — Q. N.A. B.fl.VII,288
C. decompositus, F. v. M., fragm. VIII, 267 (1874) ... ... — — — — — Q. N.A. B.fl.VII,288 M.fr.
C. Armstrongii, Bentham, Fl. Austr. VII, 289 (1878) ... ... — — — — — Q. N.A. B.fl.VII,289
C. umbellatus, Bentham, Fl. Hongk. 386 (1861) ... ... — — — — — Q. — B.fl.VII,289 M.fr.VIII,267;IX,53.
C. conicus, Boeckeler in Schlecht. Linnaea XXXVIII, 371 (1874) W.A. — — — — Q. N.A. B.fl.VII,290 M.fr.VIII,268;IX,53.

HELEOCHARIS, R. Brown, prodr. 224 (1810). (Eleocharis, Eleochiton, Scirpidium.)

H. sphacelata, R. Brown, prodr. 224 (1810) ... ... — S.A. T. V. N.S.W. Q. N.A. B.fl.VII,292 M.fr.VI,94;VIII,239.
H. spiralis, R. Brown, prodr. 224 (1810) ... ... — — — — — Q. N.A. B.fl.VII,292
H. compacta, R. Brown, prodr. 224 (1810)... ... ... — — — — N.S.W. Q. N.A. B.fl.VII,293 M.fr.VIII,239.
H. fistulosa, J. A. & J. H. Schultes, mantiss. II, 89 (1824) ... — — — — — Q. — B.fl.VII,293 M.fr.VI,93;VIII,239.
H. tetraquetra, Nees in Wight's Contrib. Bot. of Ind. 113 (1834) — — — — N.S.W. Q. — B.fl.VII,294 M.fr.VIII,239.
H. cylindrostachys, Boeckeler in Regensb. Flora, 108 (1875) ... — — — — N.S.W. Q. — B.fl.VII,294 M.fr.VIII,240.
H. acuta, R. Brown, prodr. 224 (1810) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,294 M.fr.VIII,240.
H. multicaulis, Smith, Engl. Flor. II, 64 (1824) ... ... W.A. S.A. — V. N.S.W. — — B.fl.VII,296
H. atricha, R. Brown, prodr. 225 (1810) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,295 M.fr.VII,252;IX,100.
H. capitata, R. Brown, prodr. 225 (1810) ... ... W.A. — — — — Q. N.A. B.fl.VII,296 M.fr.VI,94;VIII,240.
H. atropurpurea, Kunth, enum. II, 151 (1837) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,296 M.fr.VIII,240.
H. acicularis, R. Brown, prodr. 224 (1810)... ... ... — S.A. T. V. N.S.W. — — B.fl.VII,297 M.fr.VIII,240.

FIMBRISTYLIS, Vahl, enum. plant. II, 285 (1806). (Abildgaardia, Trichelostylis, Oncostylis)

F. acicularis, R. Brown, prodr. 226 (1810)... ... ... — — — — — Q. N.A. B.fl.VII,301 M.fr.IX,54.
F. acuminata, Vahl, enum. II, 285 (1806) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,301 M.fr.VIII,274;IX,54.
F. punctata, R. Brown, prodr. 226 (1810) ... ... — — — — — Q. — B.fl.VII,302
F. rhyticarya, F. v. M., fragm. I, 215 (1859) ... ... — — — — — Q. N.A. B.fl.VII,302 M.fr.I,215.
F. leucostachya, Boeckeler in Garcke, Linnaea XXXVIII, 385 (1874) — — — — — — N.A. B.fl.VII,302 M.fr.IX,54.
F. nutans, Vahl, enum. II, 285 (1806) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,303 M.fr.VIII,274;IX,54.
F. pauciflora, R. Brown, prodr. 225 (1810)... ... ... — — — — — Q. N.A. B.fl.VII,303

F. cardiocarpa, F. v. M., fragm. I, 194 (1859) ... ... — — — — — — N.A. B.fl.VII,303 M.fr.I,194.
F. leucocolea, Bentham, Fl. Austr. VII, 304 (1878) ... ... — — — — — — N.A. B.fl.VII,304
F. polytrichoides, R. Brown, prodr. 226 (1810) ... ... — — — — — Q. N.A. B.fl.VII,304
F. androgyna, R. Brown, prodr. 226 (1810) ... ... — — — — — — N.A. B.fl.VII,304
F. subbulbosa, Bentham, Fl. Austr. VII, 305 (1878) ... ... — — — — — Q. — B.fl.VII,305
F. tetragona, R. Brown, prodr. 226 (1810)... ... ... — — — — — — N.A. B.fl.VII,305 M.fr.I,194;VIII,274.
F. trigastrocarya, F. v. M., fragm. I, 194 (1859) ... ... — — — — — — N.A. B.fl.VII,305 M.fr.I,194.
F. monandra, F. v. M., fragm. I, 195 (1859) ... ... — — — — — — N.A. B.fl.VII,306 M.fr.I,195.
F. pterygosperma, R. Brown, prodr. 226 (1810) ... ... — — — — — — N.A. B.fl.VII,306 M.fr.I,193;IX,55.
F. sphaerocephala, Bentham, Fl. Austr. VII, 306 (1878)... ... — — — — — Q. N.A. B.fl.VII,306
F. xyridis, R. Brown, prodr. 226 (1810) ... ... — — — — — Q. N.A. B.fl.VII,307 M.fr.VIII,274;IX,54.
F. oxystachya, F. v. M., fragm. I, 195 (1859) ... ... — — — — — — N.A. B.fl.VII,307 M.fr.VIII,272.
F. macrantha, Boeckeler in Garcke in Linnaea XXXVIII, 388 (1874) — — — — — — N.A. B.fl.VII,307 M.fr.IX,55.
F. squarrulosa, F. v. M., fragm. I, 216 (1859) ... ... — — — — — — N.A. B.fl.VII,308 M.fr.I,216.
F. monostachya, Hasskarl, pl. Jav. rar. 61-64 (1848) ... ... — — — N.S.W. Q. N.A. B.fl.VII,308 M.fr.VIII,272;IX,54.
F. Brownii, Bentham, Fl. Austr. VII, 308 (1878) ... ... — — — — — — N.A. B.fl.VII,308 M.fr.VIII,273.
F. Dallachyi, F. v. M. in Benth. Fl. Austr. VII, 309 (1878) ... — — — — — Q. — B.fl.VII,309 M.fr.VIII,273.
F. velata, R. Brown, prodr. 227 (1810) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,309 M.fr.IX,11,54.
F. aestivalis, Vahl, enum. II, 288 (1806) ... ... — — — V. N.S.W. Q. N.A. B.fl.VII,310 M.fr.IX,11,54.
F. dichotoma, Vahl, enum. II, 287 (1806) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,310 M.fr.IX,10,54.
F. depauperata, R. Brown, prodr. 227 (1810) ... ... — — — — — — N.A. B.fl.VII,311
F. spirostachya, F. v. M. in Benth. Fl. Austr. VII, 311 (1878)... — — — — — — N.A. B.fl.VII,311
F. communis, Kunth, enum. II, 234 (1837) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,311 M.fr.I,196;IX,10,54.
F. ferruginea, Vahl, enum. 291 (1806) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.VII,312 M.fr.I,197;IX,10,54.
F. denudata, R. Brown, prodr. 227 (1810)... ... ... — — — — — Q. N.A. B.fl.VII,313 M.fr.IX,9.
F. elata, R. Brown, prodr. 227 (1810) ... ... — — — — — — N.A. B.fl.VII,313
F. caespitosa, R. Brown, prodr. 228 (1810) ... ... — — — — — Q. N.A. B.fl.VII,313 M.fr.I,199.
F. spiralis, R. Brown, prodr. 226 (1810) ... ... — — — — — — N.A. B.fl.VII,314
F. subaristata, Bentham, Fl. Austr. VII, 314 (1878) ... ... — — — — — — N.A. B.fl.VII,314
F. leptoclada, Bentham, Fl. Austr. VII, 314 (1878) ... ... — — — — — Q. — B.fl.VII,314
F. debilis, F. v. M., fragm. I, 198 (1859) ... ... — — — — — — N.A. B.fl.VII,315 M.fr.I,198.
F. corynocarya, F. v. M., fragm. I, 197 (1859) ... ... — — — — — — N.A. B.fl.VII,315 M.fr.I,187.
F. solidifolia, F. v. M., fragm. I, 198 (1859) ... ... — — — — — — N.A. B.fl.VII,315 M.fr.I,198.
F. obtusangula, F. v. M., fragm. I, 198 (1859) ... ... — — — — — Q. N.A. B.fl.VII,315 M.fr.I,198.
F. miliacea, Vahl, enum. II, 287 (1806) ... ... — — — — — Q. N.A. B.fl.VII,316 M.fr.I,199;IX,12,54.
F. rara, R. Brown, prodr. 227 (1810) ... ... — — — — — — N.A. B.fl.VII,316
F. microcarya, F. v. M., fragm. I, 200 (1859) ... ... — — — — — Q. N.A. B.fl.VII,316 M.fr.I,200.
F. quinquangularis, Kunth, enum. II, 229 (1837) ... ... — — — — — — N.A. B.fl.VII,317
F. cyperoides, R. Brown, prodr. 228 (1810) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,317 M.fr.VIII,273;IX,11,54.
F. furva, R. Brown, prodr. 228 (1810) ... ... — — — — — Q. — B.fl.VII,318
F. cymosa, R. Brown, prodr. 228 (1810) ... ... — — — — — — N.A. B.fl.VII,318
F. multifolia, Boeckeler in Garcke, Linnaea XXXVIII, 397 (1874) — — — — — — N.A. B.fl.VII,319 M.fr.IX,55.
F. sericea, R. Brown, prodr. 228 (1810) ... ... — — — — — — N.A. B.fl.VII,319
F. macrostachya, Boeckeler in Garcke, Linn. XXXVIII, 386 (1874) — — — — — — N.A. B.fl.VII,319 M.fr.IX,54.
F. Neilsoni, F. v. M., fragm. IX, 79 (1875) ... ... — S.A. — — N.S.W. — — B.fl.VII,320 M.fr.IX,79.
F. capitata, R. Brown, prodr. 228 (1810) ... ... — — — — — Q. N.A. B.fl.VII,320 M.fr.I,196.
F. Schultzii, Boeckeler in Garcke, Linnaea XXXVIII, 391 (1874) — — — — — — N.A. B.fl.VII,321 M.fr.IX,55.
F. barbata, Bentham, Fl. Austr. VII, 321 (1878)... ... ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,321 M.fr.IX,55.
F. capillaris, A. Gray, Man. of Bot. N.U.S. fifth ed. 567 (1867) W.A. — — — — Q. N.A. B.fl.VII,322 M.fr.IX,55.

SCIRPUS, Tournefort, inst. 528, t. 300 (1700), from Terentius. (Isolepis, Malacochaete.)

S. humillimus, Bentham, Fl. Austr. VII, 324 (1878) ... ... — — — — — — N.A. B.fl.VII,324 M.fr.IX,7.
S. fluitans, Linné, spec. plant. 48 (1753) ... ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,325 M.fr.IX,55.
S. arenarius, Bentham, Fl. Austr. VII, 325 (1878) ... ... W.A. — — V. — — — B.fl.VII,325
S. lenticularis, Sprengel, syst. I, 208 (1825) ... ... — — — — N.S.W. — — B.fl.VII,326 M.fr.IX,55.
S. crassiusculus, J. Hooker in Benth. Fl. Austr. VII, 326 (1878) — — T. V. N.S.W. — — B.fl.VII,326
S. cyperoides, Sprengel, syst. I, 208 (1825) ... ... W.A. — — — — — — B.fl.VII,326
S. setaceus, Linné, spec. plant. 49 (1753) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,327 M.fr.IX,55.
S. riparius, Sprengel, syst. I, 208 (1825) ... ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,327
S. cartilagineus, Sprengel, syst. I, 208 (1825) ... ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,328 M.fr.IX,55.
S. squarrosus, Linné, mant. plant. alt. 181 (1771)... ... ... — — — — — Q. N.A. B.fl.VII,329
S. inundatus, Sprengel, syst. I, 207 (1825)... ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,329 M.fr.IX,55.
S. prolifer, Rottboell, descr. et icon. pl. 55 t. 17 (1773) ... — — — N.S.W. — — B.fl.VII,330 M.fr.IX,55.
S. articulatus, Linné, spec. plant. 47 (1753) ... ... — — — — — Q. N.A. B.fl.VII,331 M.fr.IX,55.
S. nodosus, Rottboell, descr. et icon. pl. 52, t. 8 (1773) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,331 M.fr.IX,55.
S. supinus, Linné, spec. plant. 49 (1753) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,330 M.fr.IX,55.
S. debilis, Pursch, flor. Amer. Sept. I, 55 (1814) ... ... — — — — Q. N.A. B.fl.VII,332
S. mucronatus, Linné, spec. plant. 50 (1753) ... ... — — — — N.S.W. Q. — B.fl.VII,332 M.fr.IX,8,56.
S. pungens, Vahl, enum. II, 255 (1806) ... ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,333 M.fr.IX,8,56.
S. lacustris, Linné, spec. plant. 48 (1753) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,333 M.fr.IX,7,56.
S. litoralis, Schrader, Fl. Germ. I, 142, t. 5 (1806) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,334 M.fr.IX,7,56.
S. maritimus, Linné, spec. plant. 51 (1753) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,335 M.fr.IX,8,56.
S. grossus, Linné, fil. suppl. plant. 104 (1781) ... ... — — — — — Q. — —
S. polystachyus, F. v. M. in Transact. phil. Soc. Vict. I, 108 (1854) — — V. N.S.W. — — B.fl.VII,335 M.fr.IX,9,56.

LIPOCARPHA, R. Brown in Tuckey's narrat. of an Exped. to Congo, 459 (1818). (Hypaelyptum.)

L. argentea, R. Brown in Tuck. Congo, 459 (1818) ... ... — — — — — Q. — B.fl.VII,336 M.fr.VIII,238;IX,57.
L. microcephala, R. Brown in Tuck. Congo, 459 (1818) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,337 M.fr.VIII,238;IX,57.

FUIRENA, Rottboell, descr. et icon. rar. pl. illustr. 70 (1773).

F. umbellata, Rottboell, descr. et icon. pl. 70, t. 19 (1773) ... — — — — — Q. N.A. B.fl.VII,337 M.fr.VIII,238;IX,57.
F. glomerata, Lamarck, Illustr. des genr. I, 150 (1791) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,338 M.fr.VIII,238;IX,57.

HYPELYTRUM, L. C. Richard in Persoon, synops. I, 70 (1805). (Hypolytrum.)

H. latifolium, L. C. Richard in Pers. syn. I, 70 (1805) ... ... — — — — Q. — B.fl.VII,339 M.fr.VIII,238;IX,57.

EXOCARYA, Bentham in J. Hooker, icon. plant. t. 1206 (1877).

E. scleroides, Bentham in Hook. icon. pl. t. 1206 (1877)... ... — — — — N.S.W. Q. — B.fl.VII,339 M.fr.IX,12.

MAPANIA, Aublet, Hist. pl. Guian. 47, t. 17 (1775). (Pandanophyllum.)

M. hypelytroides, F. v. M., fragm. IX, 16 (1875)... ... ... — — — — — Q. — B.fl.VII,341 M.fr.IX,16.

SCIRPODENDRON, Zippelius in Journ. Asiat. Soc. Beng. XXXVIII, 85 (1869).

S. costatum, Kurz in Journ. Asiat. Soc. Beng. XXXVIII, 85 (1869) — — — — — Q. — B.fl.VII,341 M.fr.X,104.

LEPIRONIA, L. C. Richard in Persoon, synops. I, 70 (1805). (Chondrachne).

L. mucronata, L. C. Richard in Pers. syn. I, 70 (1805) ... ... — — — — N.S.W. Q. — B.fl.VII,342 M.fr.IX,17,58.

CHORIZANDRA, R. Brown, prodr. 221 (1810). (Chorisandra.)

C. spaerocephala, R. Brown, prodr. 221 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,344 M.fr.IX,18,58.
C. enodis, Nees in Lehm. pl. Preiss. II, 73 (1846)... ... ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,344 M fr.IX,18.
C. multiarticulata, Nees in Ann. and Mag. of nat. Hist. VI, 48 (1842) W.A. — — — — — B.fl.VII,345 M.fr.IX,18.
C. cymbaria, R. Brown, prodr. 221 (1810) ... ... W.A. — — V. N.S.W. Q. — B.fl.VII,345 M.fr.IX,18,58

OREOBOLUS, R. Brown, prodr. fl. Nov. Holl. 235 (1810).

O. Pumilio, R. Brown, prodr. 226 (1810) ... ... — — T. V. N.S.W. — — B.fl.VII,346 M.fr.IX,20,58.

REMIREA, Aublet, Hist. des pl. de la Guian. franc. I, 44, t. 16 (1775).

R. maritima, Aublet, Hist. pl. Guian. I, 45, t. 16 (1775) ... — — — — — Q. — B.fl.VII,347 M.fr.V,92;IX,20,58.

RHYNCHOSPORA, Vahl, enum. plant. II, 229 (1806). (Morisia.)

R. aurea, Vahl, enum. II, 229 (1806) ... ... — — — — — Q. — B.fl.VII,348 M.fr.IX,17,57.
R. glauca, Vahl, enum. II, 229 (1806) ... ... — — — — N.S.W. Q. — B.fl.VII,349 M.fr.IX,17,57.
R. Wallichiana, Kunth, enum. II, 289 (1837) ... ... — — — — — Q. N.A. B.fl.VII,349 M.fr.IX,17,52.
R. longisetis, R. Brown, prodr. 230 (1810)... ... ... — — — — — Q. N.A. B.fl.VII,350 M.fr.IX,17,58.
R. tenuifolia, Bentham, Fl. Austr. VII, 350 (1878) ... ... — — — — — Q. N.A. B.fl.VII,350

CYATHOCHAETE, Nees in Lehm. pl. Preiss. II, 86 (1846). (Tetralepsis.)

C. clandestina, Bentham, Fl. Austr. VII, 351 (1878) ... ... W.A. — — — — — — B.fl.VII,351 M.fr.IX,40.
C. avenacea, Bentham, Fl. Austr. VII, 351 (1878) ... ... W.A. — — — — — — B.fl.VII,351 M.fr.IX,40.
C. diandra, Nees in Lehm. pl. Preiss. II, 86 (1846) .. ... — — — V. N.S.W. — — B.fl.VII,352 M.fr.IX,39.

CARPHA, Banks & Solander in R. Brown, prodr. 230 (1810).

C. alpina, R. Brown, prodr. 280 (1310) ... ... — — T. V. N.S.W. — — B.fl.VII,381 M.fr.IX,39.

SCHOENUS, Linné, coroll. gen. 2 (1737). (Chaetospora, Elynanthus, Tricostularia, Helothrix, Isoschoenus, Gymnochaete, Gymnoschoenus, Mesomelaena, Discopodium.)

S. cruentus, F. v. M., fragm. IX, 36 (1875) ... . W.A. — — — — — — B.fl.VII,357 M.fr.IX,37.
S. Benthami, F.v.M.; S.compressus, Benth. Fl.Austr. VII,357 (1878) W.A. — — — — — — B.fl.VII,357 M.fr.IX,37.
S. lanatus, Labillard., Nov. Holl. pl. spec. I, 19, t. 20 (1804) ... W.A. — — — — — — B.fl.VII,357 M.fr.IX,37.
S. curvifolius, Poiret, Encycl. méth. suppl. II, 251 (1811) ... W.A. — — — — — — B.fl.VII,358 M.fr.IX,36.
S. capitatus, F. v. M., fragm. IX, 58 (1875) ... ... W.A. — — — — — — B.fl.VII,358 M.fr.IX,37.
S. setifolius, Bentham, Fl. Austr. VII, 359 (1878) ... .. W.A. — — — — — — B.fl.VII,359
S. Drummondii, Bentham, fl. Austr. VII, 359, non Steud. (1870) W.A. — — — — — — B.fl.VII,359 M.fr.IX,37.
S. turbinatus, Poiret, Encycl. méth. suppl. II, 251 (1811) ... — — — — N.S.W. — — B.fl.VII,359 M.fr.IX,33.
S. barbatus, Boeckeler in Garcke, Linnaea XXXVIII, 277 (1874) W.A. — — — — — — B.fl.VII,360 M.fr.IX,30.
S. flavus, Boeckeler in Garcke, Linnaea XXXVIII, 278 (1874) W.A. — — — — — — B.fl.VII,360 M.fr.IX,30.
S. brevisetis, Poiret, Encycl. méth. suppl. II, 251 (1811) ... W.A. — — — — — — B.fl.VII,360 M.fr.IX,30.
S. Armeria, Boeckeler in Garcke, Linnaea XXXVIII, 279 (1874) W.A. — — — — — — B.fl.VII,361 M.fr.IX,30
S. aphyllus, Boeckeler in Garcke, Linnaea XXXVIII, 280 (1874) — S.A. — V. N.S.W. — — B.fl.VII,361 M.fr.IX,28.
S. imberbis, R. Brown, prodr. 231 (1810) ... ... — — — V. N.S.W. — — B.fl.VII,361 M.fr.IX,28.
S. ericetorum, R. Brown, prodr. 231 (1810) ... . — — — — N.S.W. Q. — B.fl.VII,362 M.fr.IX,58
S. nitens, Poiret, Encycl. méth. suppl. II, 251 (1811) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,362 M.fr.IX,35.
S. cygneus, Nees in Lehmann, pl. Preiss. II, 81 (1846) ... ... W.A. — — — — — — B.fl.VII,363 M.fr.IX,80.
S. minutulus, F. v. M., fragm. IX, 32 (1875) ... ... W.A. — — — — — — B.fl.VII,363 M.fr.IX,32.
S. trachycarpus, F. v. M., fragm. IX, 33 (1875) ... ... W.A. — — — — — — B.fl.VII,363 M.fr.IX,33.
S. Tepperi, F. v. M., fragm. XI, 106 (1880) ... ... — S.A. — — — — — — — M.fr.XI,106.
S. nanus, F. v. M., fragm. IX, 36 (1875) ... ... W.A. — — — — — — B.fl.VII,364 M.fr.IX,28.
S. pleiostemoneus, F. v. M., fragm. IX, 52 (1875) ... ... W.A. — — — — — — B.fl.VII,364 M.fr.IX,52.
S. breviculmis, Bentham, Fl. Austr. VII, 364 (1878) ... ... W.A. — — — — — — B.fl.VII,364
S. deformis, Poiret, Encycl. méth. suppl. II, 251 (1811) ... — S.A. — — — — — B.fl.VII,364 M.fr.IX,39.
S. unispiculatus, F. v. M. in Benth. Fl. Austr. VII, 365 (1878) W.A. — — — — — — B.fl.VII,365
S. obtusifolius, Boeckeler in Garcke, Linnaea XXXVIII, 281 (1874) W.A. — — — — — — B.fl.VII,366 M.fr.IX,31.
S. grammatophyllus, F. v. M., fragm. IX, 31 (1875) ... ... W.A. — — — — — — B.fl.VII,366 M.fr.IX,31.
S. asperocarpus, F. v. M., fragm. IX, 29 (1875) ... ... W.A. — — — — — — B.fl.VII,366 M.fr.IX,29.
S. Moorei, Bentham, Fl. Austr. VII, 367 (1878) ... ... — — — — N.S.W. — — B.fl.VII,367
S. villosus, R. Brown, prodr. 231 (1810) ... ... — — — — N.S.W. — — B.fl.VII,367 M.fr.IX,28.
S. grandiflorus, F. v. M., fragm. IX, 30 (1875) ... ... W.A. — — — — — — B.fl.VII,367 M.fr.IX,30.
S. distans; Chaetospora, F. v. M., fragm. IX, 35 (1875) ... W.A. — — — — — — — M.fr.IX,35.
S. calostachyus, Poiret, Encycl. méth. suppl. II, 251 (1811) ... — — — — N.S.W. Q. — B.fl.VII,368 M.fr.IX,35.
S. scabripes, Bentham, Fl. Austr. VII, 368 (1878) ... ... — — — — — Q. — B.fl.VII,368
S. multiglumis, Bentham, Fl. Austr. VII, 368 (1878) ... ... W.A. — — — — — — B.fl.VII,368
S. efoliatus, F. v. M., fragm. IX, 32 (1875) ... ... W.A. — — — — — — B.fl.VII,369 M.fr.IX,32.
S. acuminatus, R. Brown, prodr. 231 (1810) ... ... W.A. — — — — — — B.fl.VII,369 M.fr.IX,80.
S. pedicellatus, Poiret, Encycl. méth. suppl. II, 251 (1811) ... W.A. — — — — — — B.fl.VII,369
S. fascicularis, Nees in Ann. and Mag. of nat. Hist. VI, 48 (1842) W.A. — — — — — — B.fl.VII,370
S. brevifolius, R. Brown, prodr. 231 (1810) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,370 M.fr.IX,29,80.
S. melanostachys, R. Brown, prodr. 231 (1810) ... ... — — — V. N.S.W. Q. — B.fl.VII,370 M.fr.IX,29.
S. spartens, R. Brown, prodr. 231 (1810) ... ... — — — — — Q. N.A. B.fl.VII,371
S. vaginatus, F. v. M. in Benth. Fl. Austr. VII, 371 (1878) ... — — — — N.S.W. Q. — B.fl.VII,371

S. falcatus, R. Brown, prodr. 232 (1810) ... ... — — — — — Q. N.A. B.fl.VII,372 M.fr.IX,29.
S. punctatus, R. Brown, prodr. 232 (1810)... ... ... — — — — — — N.A. B.fl.VII,372
S. indutus, F. v. M. in Benth. Fl. Austr. VII, 372 (1878) ... W.A. — — — — — — B.fl.VII,372
S. bifidus, Boeckeler in Garcke, Linnaea XXXVIII, 282 (1874) W.A. — — — — — — B.fl.VII,373 M.fr.IX,38,80.
S. apogon, Roemer et Schultes, syst. veg. II, 77 (1817) ... — S.A. T. V. N.S.W. Q. — B.fl.VII,373 M.fr.IX,56;X,120.
S. odontocarpus, F. v. M., fragm. IX, 32 (1875) ... ... W.A. — — — — — — B.fl.VII,374 M.fr.IX,32.
S. humilis, Bentham, Fl. Austr. VII, 374 (1878) ... ... W.A. — — — — — — B.fl.VII,374
S. sculptus, Boeckeler in Garcke, Linnaea XXXVIII, 286 (1874) W.A. — — — — — — B.fl.VII,374 M.fr.IX,30.
S. axillaris, Poiret, encycl. méth. suppl. II, 251 (1811) ... ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,375 M.fr.IX,34.
S. tenellus, Bentham, Fl. Austr. VII, 375 (1878)... ... ... W.A. — — — — — — B.fl.VII,375
S. natans, F. v. M., fragm. IX, 36 (1875) ... ... W.A. — — — — — — B.fl.VII,375 M.fr.IX,38.
S. fluitans, J. Hooker, Fl. Tasman. II, 81, t. 141 (1860)... ... — — T. — — — — B.fl.VII,376 M.fr.IX,28.
S. octandrus, F. v. M., fragm. IX, 31 (1875) ... ... W.A. — — — — — — B.fl.VII,377 M.fr.IX,31.
S. capillaris, Chaetospora, F. v. M., fragm. IX, 377 (1875) ... W.A. S.A. T. V. — — — B.fl.VII,377 M.fr.IX,34,80.
S. stygius, Poiret, encycl. méth. suppl. II, 251 (1811) ... ... W.A. — — — — — — B.fl.VII,378 M.fr.IX,36.
S. deustus, F. v. M.; Carpha deusta, R. Brown, prodr. 230 (1810) — — — — N.S.W. Q. — B.fl.VII,379 M.fr.IX,39.
S. tetragonus, Poiret, Encycl. méth. suppl. II, 251 (1811) ... W.A. — — — — — — B.fl.VII,380 M.fr.IX,36.
S. sphaerocephalus, Poiret, Encycl. méth. suppl. II, 251 (1811)... — — T. V. N.S.W. — — B.fl.VII,380 M.fr.IX,33.
S. anceps, Poiret, Encycl. méth. suppl. II, 251 (1811) ... ... W.A. — — — — — — B.fl.VII,380 M.fr.IX,30.
S. paludosus, Poiret, Encycl. méth. suppl. II, 251 (1811) ... — — — — N.S.W. Q. — B.fl.VII,382 M.fr.IX,35.
S. pauciflorus; Lepidosperma, F. v. M., fragm. IX, 23 (1875) ... — — — V. — — — B.fl.VII,383 M.fr.IX,23.
S. Tricostularia, F. v. M.; Tricostularia compressa, Nees in pl. Preiss. II, 83 (1846) ... ... W.A. — — — — — — B.fl.VII,383 M.fr.IX,37.
S. Neesii, F.v. M.; Tricostularia, Lehmann, pl. Preiss. II, 83 (1846) W.A. — — — — — — B.fl.VII,383 M.fr.IX,37.
S. fimbristyloides; Chaetospora, F. v. M., fragm. IX, 34 (1875) — — — — — — N.A. B.fl.VII,384 M.fr.IX,34.

LEPIDOSPORA, F. v. M., fragm. phytograph. Austral. IX, 34 (1875).
L. tenuissima, F. v. M., fragm. IX, 34 (1875) ... ... — S.A. T. V. N.S.W. — — B.fl.VII,365 M.fr.IX,34.

LEPIDOSPERMA, Labillardière, Nov. Holl. pl. spec. I, 14 (1804).
L. gladiatum, Labillardière, Nov. Holl. pl. spec. I, 15, t. 12 (1804) W.A. S.A. T. V. N.S.W. — — B.fl.VII,387
L. effusum, Bentham, Fl. Austr. VII, 387 (1878)... ... ... W.A. — — — — — — B.fl.VII,387
L. rupestre, Bentham, Fl. Austr. VII, 388 (1878)... ... ... W.A. — — — — — — B.fl.VII,388
L. elatius, Labillardière, Nov. Holl. pl. spec. I, 15, t. 11 (1804) — S.A. T. V. — — — B.fl.VII,388 M.fr.IX,25.
L. tetraquetrum, Nees in Lehm. pl. Preiss. II, 90 (1846) ... W.A. — — — — — — B.fl.VII,388 M.fr.IX,24.
L. Oldfieldii, J. Hooker, Fl. Tasman. II, 91, t. 146 (1860) ... — — T. — — — — B.fl.VII,389
L. exaltatum, R. Brown, prodr. 234 (1810)... ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,389
L. longitudinale, Labillardière, Nov. Holl. pl. spec. I, 16, t. 13 (1804) W.A. S.A. T. V. N.S.W. — — B.fl.VII,389 M.fr.IX,25.
L. concavum, R. Brown, prodr. 234 (1810)... ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,390 M.fr.IX,24.
L. angustatum, R. Brown, prodr. 235 (1810) ... ... W.A. — — — — — — B.fl.VII,391 M.fr.IX,26,80.
L. Drummondii, Bentham, Fl. Austr. VII, 391 (1878) ... ... W.A. — — — — — — B.fl.VII,391
L. Brunonianum, Nees in Lehm. pl. Preiss. II, 92 (1846) ... W.A. — — — — — — B.fl.VII,392
L. tuberculatum, Nees in Lehm. pl. Preiss. II, 90 (1846) ... W.A. — — — — — — B.fl.VII,392 M.fr.IX,26.
L. resinosum, F. v. M. in Benth. Fl. Austr. VII, 392 (1878) ... W.A. — — — — — — B.fl.VII,392
L. viscidum, R. Brown, prodr. 234 (1810) ... ... — S.A. — V. N.S.W. — — B.fl.VII,393
L. costale, Nees in Lehm. pl. Preiss. II, 92 (1846) ... ... W.A. — — — — — — B.fl.VII,393
L. laterale, R. Brown, prodr. 234 (1810) ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,393
L. congestum, R. Brown, prodr. 234 (1810) ... ... — S.A. — — — — B.fl.VII,394
L. globosum, Labillardière, Nov. Holl. pl. spec. I, 16, t. 14 (1804) — S.A. T. V. — — — B.fl.VII,394
L. lineare, R. Brown, prodr. 235 (1810) ... ... — S.A. T. V. N.S.W. — — B.fl.VII,395 M.fr.IX,26.
L. aphyllum, R. Brown, prodr. 235 (1810)... ... ... W.A. — — — — — — B.fl.VII,395
L. gracile, R. Brown, prodr. 235 (1810) ... ... W.A. — — — — — — B.fl.VII,395
L. semiteres, F. v. M. in Garcke, Linnaea XXXVIII, 327 (1874) — S.A. — V. — — — B.fl.VII,395
L. canescens, Boeckeler in Garcke, Linnaea XXXVIII, 330 (1874) — S.A. — V. N.S.W. — — B.fl.VII,396 M.fr.IX,24.
L. pubisquameum, Steudel, syn. pl. glum. II, 158 (1855) ... W.A. — — — — — — B.fl.VII,396 M.fr.IX,27.
L. scabrum, Nees in Lehm. pl. Preiss. II, 92 (1846) ... ... W.A. — — — — — — B.fl.VII,397 M.fr.IX,27.
L. tenue, Bentham, Fl. Austr. VII, 397 (1878) ... ... W.A. — — — — — — B.fl.VII,397
L. leptostachyum, Bentham, Fl. Austr. VII, 397 (1878)... ... W.A. — — — — — — B.fl.VII,397
L. leptophyllum, Bentham, Fl. Austr. VII, 398 (1878) ... ... W.A. — — — — — — B.fl.VII,398
L. tortuosum, F. v. M., general Report 9 (1855) ... ... — — — V. — — — B.fl.VII,398 M.fr.IX,23.
L. flexuosum, R. Brown, prodr. 235 (1810) ... ... — — — — N.S.W. — — B.fl.VII,398
L. filiforme, Labillardière, Nov. Holl. pl. spec. I, 17, t. 15 (1804) — S.A. T. V. N.S.W. — — B.fl.VII,398 M.fr.IX,27.
L. striatum, R. Brown, prodr. 235 (1810) ... ... W.A. — — — — — — B.fl.VII,399 M.fr.IX,26,27.
L. Neesii, Kunth, enum. II, 319 (1837) ... ... — — — V. N.S.W. — — B.fl.VII,399 M.fr.IX,24,27.
L. carphoides, F. v. M. in Benth. Fl. Austr. VII, 400 (1878) ... W.A. S.A. — V. — — — B.fl.VII,400

CLADIUM, P. Browne, Civ. and Nat. Hist. of Jamaic. 114 (1756). (Gahnia, Lampocarya, Melachne, Epiandra, Didymonema, Baumea, Chapelliera, Morelotia.)
C. Mariscus, R. Brown, prodr. 236 (1810) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,402 M.fr.IX,14,57.
C. insulare, Bentham, Fl. Austr. 403 (1878) ... ... — — — — N.S.W. — — B.fl.VII,403
C. articulatum, R. Brown, prodr. 237 (1810) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,403 M.fr.IX,14,57.
C. arthrophyllum, F. v. M., fragm. IX, 14 (1875) ... ... W.A. — — — — — — B.fl.VII,403 M.fr.IX,14.
C. glomeratum, R. Brown, prodr. 237 (1810) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,404 M.fr.IX,15,57.
C. Preissii, F. v. M. in Benth. Fl. Austr. VII, 405 (1878) ... W.A. — — — — — — B.fl.VII,405 M.fr.IX,13,15,56.
C. laxum, Bentham, Fl. Austr. VII, 405 (1878) ... ... W.A. — — — — — — B.fl.VII,405
C. riparium, Bentham, Fl. Austr. VII, 405 (1878) ... ... W.A. — — — — — — B.fl.VII,405
C. teretifolium, R. Brown, prodr. 237 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,406 M.fr.IX,15,56.
C. tetraquetrum, J. Hooker, Fl. Tasman. II, 95, t. 149 (1860) ... — S.A. T. V. N.S.W. — — B.fl.VII,406
C. schoenoides, R. Brown, prodr. 237 (1810) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,407
C. Gunnii, J. Hooker, Fl. Tasman. II, 95, t. 148 (1860)... ... — S.A. T. V. N.S.W. — — B.fl.VII,407 M.fr.IX,15,57.
C. junceum, R. Brown, prodr. 237 (1810) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,408 M.fr.IX,16,57.
C. vaginale, Bentham, Fl. Austr. VII, 409 (1878)... ... ... W.A. — — — — — — B.fl.VII,408
C. elynanthoides, F. v. M., fragm. IX, 31 (1875)... .. ... W.A. — — — — — — B.fl.VII,409 M.fr.IX,31.

C. Filum, R. Brown, prodr. 237 (1810) ... ... — S.A. T. V. — — — B.fl.VII,409 M.fr.IX,14,57.
C. trifidum, F. v. M. in Pap. R. S. Tasm. 117 (1878) ... ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,413
C. decompositum, R. Brown, prodr. 237 (1810) ... ... W.A. — — — — — — B.fl.VII,417 M.fr.IX,18.
C. microstachyum, F.v.M.; Gahnia, Benth. Fl.Austr.VII,414(1878) — — — V. N.S.W. — — B.fl.VII,414
C. polyphyllum, F. v. M.; Gahnia, Benth. Fl. Austr. VII, 415 (1878) W.A. — — — — — — B.fl.VII,415
C. ancistrophyllum, F. v. M. in Benth. Fl. Austr. VII, 415 (1878) W.A. — — — — — — B.fl.VII,415
C. lanigerum, R. Brown, prodr. 237 (1810)... ... ... W.A. S.A. — — — — — B.fl.VII,415 M.fr.IX,14,57.
C. aristatum, F. v. M. in Benth. Fl. Austr. 416 (1878) ... ... W.A. — — — — — — B.fl.VII,416
C. deustum, R. Brown, prodr. 237 (1810) ... ... W.A. S.A. — — — — — B.fl.VII,416 M.fr.IX,14.
C. Radula, R. Brown, prodr. 237 (1810) ... ... — — T. V. — — — B.fl.VII,417 M.fr.IX,13,56.
C. Sieberi, F. v. M., fragm. IX, 14 (1875)... ... ... — — — — N.S.W. — — B.fl.VII,414 M.fr.IX,14,57.
C. tetragonocarpum, F. v. M.; Gahnia, Boeckeler in Linnaea XXXVIII, 347 (1874) ... ... — — — V. — — — B.fl.VII,418
C. melanocarpum, F. v. M., fragm. IX, 13 (1875) ... ... — — T. V. N.S.W. — — B.fl.VII,414 M.fr.IX,13,56.
C. psittacorum, F. v. M., fragm. IX, 13 (1875) ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,418 M.fr.IX,13,56.
C. asperum, F. v. M., fragm. IX, 12 (1875) ... ... — — — — N.S.W. Q. — B.fl.VII,412 M.fr.IX,12,56.
C. xanthocarpum, F. v. M., fragm. IX, 13 (1875)... ... ... — — — — N.S.W. — — B.fl.VII,418 M.fr.IX,13,56.

CAUSTIS, R. Brown, prodr. 239 (1810). (Eurostorrhiza.)

C. pentandra, R. Brown, prodr. 240 (1810)... ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,420 M.fr.IX,19,58.
C. flexuosa, R. Brown, prodr. 239 (1810) ... ... — — — V. N.S.W. Q. N.A. B.fl.VII,421 M.fr.IX,58.
C. recurvata, Sprengel, syst. cur. post. 26 (1827)... ... ... — — — — N.S.W. — — B.fl.VII,421 M.fr.IX,19.
C. restiacea, F. v. M. in Benth. Fl. Austr. VII, 421 (1878) ... — — — V. N.S.W. — — B.fl.VII,421 M.fr.IX,19.
C. dioica, R. Brown, prodr. 239 (1810) ... ... W.A. — — — — — — B.fl.VII,422 M.fr.IX,19,58.

ARTHROSTYLIS, R. Brown, prodr. 229 (1810). (Arthrostyles.)

A. aphylla, R. Brown, prodr. 229 (1810) ... ... — — — — — Q. N.A. B.fl.VII,423 M.fr.IX,9,55.

REEDIA, F. v. M., fragm. I, 240, t. 10 (1859).

R. spathacea, F. v. M., fragm. I, 240, t. 10 (1859) ... ... W.A. — — — — — — B.fl.VII,423 M.fr.IX,15.

EUANDRA, R. Brown, prodr. 239 (1810). (Evandra.)

E. aristata, R. Brown, prodr. 239 (1810) ... ... W.A. — — — — — — B.fl.VII,424 M.fr.IX,18.
E. pauciflora, R. Brown, prodr. 239 (1810) ... ... W.A. — — — — — — B.fl.VII,425 M.fr.IX,18.

SCLERIA, Bergius in Svensk. Vet. Acad. Handl. 149, t. 4 (1765). (Diplacrum, Hypoporum, Sphaeropus.)

S. caricina, Bentham, Fl. Austr. VII (1878) ... ... — — — — — Q. — B.fl.VII,426 M.fr.IX,58.
S. pygmaea, R. Brown, prodr. 240 (1810) ... ... — — — — — Q. N.A. B.fl.VII,427 M.fr.IX,22,58.
S. rugosa, R. Brown, prodr. 240 (1810) ... ... — — — — — Q. N.A. B.fl.VII,428 M.fr.IX,22,58.
S. laxa, R. Brown, prodr. 240 (1810) ... ... — — — — — Q. N.A. B.fl.VII,428 M.fr.IX,21.
S. Brownii, Kunth, enum. II, 349 (1837) ... ... — — — — — Q. N.A. B.fl.VII,429 M.fr.IX,21,58.
S. lithosperma, Willdenow, spec. plant. IV, 316 (1805)... ... — — — — — Q. N.A. B.fl.VII,429 M.fr.IX,21,58.
S. tessellata, Willdenow, spec. plant. IV, 315 (1805) ... ... — — — — — Q. — B.fl.VII,430
S. margaritifera, Willdenow, spec. plant. IV, 312 (1805) — — — — — Q. — B.fl.VII,430 M.fr.IX,21,58.
S. Graeffeana, Boeckeler in Regensb. Flora, 121 (1875) ... ... — — — — — Q. N.A. B.fl.VII,431
S. hebecarpa, Nees in Wight, Contrib. to Bot. of Ind. 117 (1834) — — — — N.S.W. Q. N.A. B.fl.VII,431 M.fr.IX,21,58.
S. Chinensis, Kunth, enum. II, 357 (1837)... ... ... — — — — — Q. — B.fl.VII,431 M.fr.IX,20,58.
S. oryzoides, Presl, reliq. Haenk. I, 201 (1830) ... ... — — — — — — N.A. B.fl.VII,432
S. sphacelata, F. v. M., fragm. IX, 20 (1875) ... ... — — — — N.S.W. Q. — B.fl.VII,432 M.fr.IX,20.

UNCINIA, Persoon, synops. plant. I, 534 (1807).

U. tenella, R. Brown, prodr. 241 (1810) ... ... — — T. V. — — — B.fl.VII,433 M.fr.VIII,151.
U. compacta, R. Brown, prodr. 241 (1810)... ... ... — — T. V. — — — B.fl.VII,434 M.fr.VIII,152.
U. riparia, R. Brown, prodr. 241 (1810) ... ... — — T. V. N.S.W. — — B.fl.VII,434 M.fr.VIII,152.
U. debilior, F. v. M., fragm. VIII, 151 (1874) ... ... — — — — N.S.W. — — B.fl.VII,435 M.fr.VIII,151.

CAREX, Ruppius, Fl. Jenens. 306 (1718).

C. cephalotes, F. v. M. in Transact. phil. Soc. Vict. I, 110 (1854) — — — V. N.S.W. — — B.fl.VII,437 M.fr.VIII,251.
C. acicularis, Boott in J. Hook. Fl. N. Zel. I, 280, t. 63 (1853) — — T. V. N.S.W. — — B.fl.VII,437 M.fr.VIII,251.
C. capillacea, Boott, Illustr. of the gen. Carex I, 44, t. 110 (1858) — — — — N.S.W. — — B.fl.VII,438 M.fr.IX,191.
C. inversa, R. Brown, prodr. 242 (1810) ... . W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,438 M.fr.VIII,252.
C. canescens, Linné, spec. plant. 974 (1753) ... ... — — — V. N.S.W. — — B.fl.VII,439 M.fr.VIII,253.
C. echinata, Murray, prodr. design. stirp. Goett. 76 (1770) ... — — — V. N.S.W. — — B.fl.VII,439 M.fr.VIII,252.
C. hypandra, F. v. M., fragm. VIII, 259 (1874) ... ... — — — V. N.S.W. — — B.fl.VII,439 M.fr.VIII,259.
C. chlorantha, R. Brown, prodr. 242 (1810) ... ... — S.A. T. V. N.S.W. — — B.fl.VII,440 M.fr.VIII,256.
C. paniculata, Linné, amoen. acad. IV, 294 (1759) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,440 M.fr.VIII,256.
C. declinata, Boott, Ill. of the gen. Carex IV, 17, t. 580 (1867) — — — V. N.S.W. Q. — B.fl.VII,441 M.fr.VIII,257.
C. tereticaulis, F. v. M., fragm. VIII, 256 (1874) ... ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,441 M.fr.VIII,256.
C. fissilis, Boott, Ill. of the gen. Carex II, 86, t. 245 (1860) ... — — — — — Q. — B.fl.VII,441 M.fr.VIII,249;IX,80.
C. gracilis, R. Brown, prodr. 242 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,442 M.fr.VIII,250.
C. contracta, F. v. M., fragm. VIII, 258 (1874) ... ... — — — — N.S.W. — — B.fl.VII,442 M.fr.VIII,258.
C. Gaudichaudiana, Kunth, enum. plant. II, 417 (1837)... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,442 M.fr.VIII,257.
C. acuta, Linné, spec. plant. 978 (1753) ... ... — — — V. N.S.W. Q. — B.fl.VII,443 M.fr.VIII,259.
C. lobolepis, F. v. M., fragm. VIII, 258 (1874) ... ... — — — — N.S.W. — — B.fl.VII,443 M.fr.VIII,258.
C. flava, Linné, spec. plant. 975 (1753) ... ... — — T. — — — — B.fl.VII,444 M.fr.VIII,250.
C. Buxbaumii, Wahlenberg in Act. Holmiens. 139 (1803) ... — — — V. N.S.W. — — B.fl.VII,444 M.fr.VIII,252.
C. pumila, Thunberg, Fl. Japon. 38 (1784)... .. .. — S.A. T. V. N.S.W. Q. — B.fl.VII,445 M.fr.VIII,251.
C. breviculmis, R. Brown, prodr. 242 (1810) ... ... — S.A. T. V. N.S.W. — — B.fl.VII,445 M.fr.VIII,255.
C. Preissii, Nees in Lehm. pl. Preiss. II, 94 (1846) ... ... W.A. — — — — — — B.fl.VII,446 M.fr.VIII,251.
C. Gunniana, Boott in Transact. Linn. Soc. XX, 143 (1851) ... — S.A. T. V. N.S.W. — — B.fl.VII,446 M.fr.VIII,251.
C. Bichenoviana, Boott in J. Hook. Fl. Tasm. II, 101 (1860) ... — — T. — — — — B.fl.VII,446
C. maculata, Boott in Transact. Linn. Soc. XX, 128 (1851) ... — — — — N.S.W. Q. — B.fl.VII,447 M.fr.VIII,258.
C. Brownii, Tuckerman, enum. meth. Caric. 21 (1849) ... ... — — — V. N.S.W. — — B.fl.VII,447 M.fr.IX,250.
C. alsophila, F. v. M., fragm. VIII, 257 (1874) ... ... — — — V. — — — B.fl.VII,447 M.fr.VIII,257.
C. longifolia, R. Brown, prodr. 242 (1810)... ... ... — — T. V. N.S.W. Q. — B.fl.VII,448 M.fr.VIII,250.
C. Pseudo-Cyperus, Linné, spec. plant. 978 (1753) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,448 M.fr.VIII,249.

GRAMINEAE.

Haller, enum. stirp. Helv. I, 203 (1742).

ERIOCHLOA, Humboldt, Bonpland & Kunth, nov. gen. et sp. pl. I, 94, t. 30 (1815). (Helopus.)

E. punctata, Hamilton, prodr. pl. Ind. occ. 5 (1825) ... ... — — — V. N.S.W. Q. N.A. B.fl.VII,462 M.fr.VI,85;VIII,126
E. annulata, Kunth, Graminées I, 30 (1832) ... ... — S.A. — — N.S.W. Q. — B.fl.VII,463 M.fr.VI,85.

PASPALUM, Linné, syst. nat. ed. decim. 855 (1759).

P. scrobiculatum, Linné, mantiss. 29 (1767) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,360 M.fr.VIII,156.
P. distichum, Linné, syst. nat. ed. decim. 855 (1759) ... ... W.A. — — — N.S.W. Q. — B.fl.VII,460 M.fr.VIII,156.
P. brevifolium, Fluegge, gram. monogr. 150 (1810) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,461 M.fr.VIII,195.
P. minutiflorum, Steudel, sym. glum. I, 17 (1855) ... ... — — — — — Q. — B.fl.VII,461

PANICUM, Tournefort, inst. 515, t. 298 (1700), from Plinius. (Isachne, Digitaria, Echinochloa, Coridochloa, Chamaeraphis.)

P. coenicolum, F. v. M. in Transact. Vict. Inst. 45 (1854) ... W.A. S.A. — V. N.S.W. — — B.fl.VII,467
P. divaricatissimum, R. Brown, prodr. 192 (1810) ... — S.A. — V. N.S.W. Q. — B.fl.VII,467 M.fr.VIII,154;X,76.
P. macractinum, Bentham, Fl. Austr. VII, 468 (1878) ... ... — — — — N.S.W. Q. — B.fl.VII,468
P. papposum, R. Brown, prodr. 192 (1810) ... ... — — — — — — N.A. B.fl.VII,468 M.fr.VIII,155,196.
P. sanguinale, Linné, spec. plant. 57 (1753) ... ... W.A. — — V. N.S.W. Q. N.A. B.fl.VII,469 M.fr.VIII,153.
P. ctenanthum, F. v. M., fragm. VIII, 153 (1874) ... ... — — — — — — N.A. B.fl.VII,469 M.fr.VIII,153.
P. stenostachyum, Bentham, Fl. Austr. VII, 470 (1878) ... — — — — — N.A. B.fl.VII,470
P. tenuissimum, Bentham, Fl. Austr. VII, 470 (1878) ... — — — — N.S.W. Q. — B.fl.VII,470
P. parviflorum, R. Brown, prodr. 192 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,470 M.fr.VIII,196;XI,129.
P. Baileyi, Bentham, Fl. Austr. VII, 470 (1878) ... ... — — — — N.S.W. Q. — B.fl.VII,471
P. gibbosum, R. Brown, prodr. 193 (1810)... ... ... — — — — — Q. N.A. B.fl.VII,471 M.fr.VIII,155;XI,129.
P. leucophaeum, Humb., Bonpl. & Kunth, nov. gen. et spec. I, 97 (1815) — S.A. — V. N.S.W. Q. — B.fl.VII,472 M.fr.VIII,155.
P. semialatum, R. Brown, prodr. 192 (1810) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,472 M.fr.VIII,196.
P. rarum, R. Brown, prodr. 189 (1810) ... ... — — — — — — N.A. B.fl.VII,473
P. argenteum, R. Brown, prodr. 190 (1810) ... ... — — — — — — N.A. B.fl.VII,473
P. holosericeum, R. Brown, prodr. 190 (1810) ... ... — — — — — Q. N.A. B.fl.VII,473 M.fr.VIII,155.
P. flavidum, Retzius, observ. IV, 15 (1787) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,474 M.fr.VIII,189.
P. gracile, R. Brown, prodr. 190 (1810) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,475 M.fr.VIII,189.
P. prostratum, Lamarck, illustr. I, 171 (1791) ... ... — — — — — Q. N.A. B.fl.VII,476 M.fr.VIII,191.
P. helopus, Trinius in Spreng. Neue Entd. II, 84 (1821) ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,476
P. Gilesii, Bentham, Fl. Austr. VII, 477 (1878) ... ... — S.A. — — — Q. N.A. B.fl.VII,477 M.fr.XI,129.
P. piligerum, F. v. M. in Benth. Fl. Austr. VII, 477 (1878) ... — — — — — — N.A. B.fl.VII,477
P. polyphyllum, R. Brown, prodr. 190 (1810) ... ... — — — — — — N.A. B.fl.VII,477 M.fr.VIII,194.
P. distachyon, Linné, mantiss. alt. pl. 138 (1771) ... ... — S.A. — — N.S.W. Q. — B.fl.VII,478 M.fr.VIII,194.
P. reversum, F. v. M., fragm. VIII, 152 (1874) ... ... W.A. S.A. — — N.S.W. Q. — B.fl.VII,478 M.fr.X,76.
P. colonum, Linné, syst. nat. ed. decim. 5 (1758)... ... ... — — — — — — N.A. B.fl.VII,478
P. Crus Galli, Linné, spec. plant. 56 (1753) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,479 M.fr.VIII,198.
P. myosuroides, R. Brown, prodr. 189 (1810) ... ... — — — — — Q. N.A. B.fl.VII,480
P. Indicum, Linné, mantiss. pl. alt. 184 (1771) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,480 M.fr.VIII,197;XI,129.
P. Myurus, Lamarck, illustr. I, 172 (1791) ... ... — — — — — Q. — B.fl.VII,480 M.X,114.
P. foliosum, R. Brown, prodr. 191 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,481 M.fr.VIII,194.
P. adspersum, Trinius, spec. gram. t. 169 (1830) ... . — S.A. — — N.S.W. Q. — B.fl.VII,481
P. inaequale, F. v. M., fragm. VIII, 189 (1874) ... ... — — — — — Q. — B.fl.VII,482 M.fr.XI,129. [129.
P. uncinulatum, R. Brown, prodr. 191 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,482 M.fr.VIII,195;X,76;XI,
P. majusculum, F. v. M. in Benth. Fl. Austr. VII, 482 (1878)... — — — — — — N.A. B.fl.VII,482
P. pauciflorum, R. Brown, prodr. 183 (1810) ... ... — S.A. — — — Q. N.A. B.fl.VII,483 M.fr.VIII,193.
P. semitonsum, F. v. M. in Benth. Fl. Austr. VII, 483 (1878) . — — — — — — N.A. B.fl.VII,483
P. antidotale, Retzius, observ. bot. IV, 17 (1878)... ... ... — — — — — — N.A. B.fl.VII,483
P. repens, Linné, spec. pl. ed. sec. 87 (1762) ... ... — — V. N.S.W. Q N.A. B.fl.VII,484
P. Buncei, F. v. M. in Benth. Fl. Austr. VII, 487 (1878) ... — — — — Q. — B.fl.VII,487
P. capillipes, Bentham, Fl. Austr. VII, 484 (1878) ... ... — — — — — — N.A. B.fl.VII,484 M.fr.VIII,193.
P. pygmaeum, R. Brown, prodr. 191 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,484 M.fr.VIII,193.
P. brevifolium, Linné, spec. plant. 59 (1753) ... ... — — — — — Q N.A. B.fl.VII,485 M.fr.VIII,193;XI,129.
P. hermaphroditum, Steudel, syn. glum. I, 67 (1855) ... ... — — — — — Q. — B.fl.VII,485
P. marginatum, R. Brown, prodr. 190 (1810) .. .. — — — V. N.S.W. Q. — B.fl.VII,485 M.fr.VIII,190.
P. lachnophyllum, Bentham, Fl. Austr. VII, 486 (1878) ... — — — — — Q. — B.fl.VII,486
P. obseptum, Trinius, gram. panic. diss. alt. 149 (1826)... ... — — — N.S.W. — — B.fl.VII,486
P. bicolor, R. Brown, prodr. 191 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,487
P. melananthum, F. v. M. in Transact. Vict. Inst. 47 (1854) ... — — — V. N.S.W. Q. — B.fl.VII,488 M.fr.VIII,192.
P. effusum, R. Brown, prodr. 191 (1810) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,488 M.fr.VIII,191.
P. Mitchelli, Bentham, Fl. Austr. VII, 489 (1878) ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,489 M.fr.X,76.
P. decompositum, R. Brown, prodr. 191 (1810) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,489 M.fr.VI,86;VIII,191.
P. trachyrhachis, Bentham, Fl. Austr. VII, 490 (1878) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,490 M.fr.VIII,192;XI,129.
P. prolutum, F. v. M. in Transact. Vict. Inst. 46 (1854) ... — S.A. — V. N.S.W. Q. — B.fl.VII,490 M.fr.VIII,192.
P. spinescens, R. Brown, prodr. 193 (1810) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,499 M.fr.VIII,197;XI,129.
P. paradoxum, R. Brown, prodr. 193 (1810) ... ... — — V. N.S.W. — — B.fl.VII,499 M.fr.VIII,197.
P. Chamaeraphis, Trinius in Mem. acad. Petersb. ser. 6, III, 217 (1835) — — — — — — N.A. B.fl.VII,500
P. atro-virens, Trinius in Spreng. neue Entdeckung. II, 89 (1821) — — — V. N.S.W. Q. — B.fl.VII,625 M.fr.VIII,193;XI,129.
P. Myosotis, Steudel, syn. glum. I, 96 (1855) ... ... — — — — — Q. — B.fl.VII,625 M.fr.VIII,193.

OPLISMENUS, Palisot, Flore d'Owarc et de Benin. II, 14 (1807). (Orthopogon.)

O. compositus, Palisot, agrostogr. 54 (1812) ... ... — — — V. N.S.W. Q. — B.fl.VII,491 M.fr.VIII,199.

SETARIA, Palisot, Essai d'une nouv. Agrostogr. 51, t. 13 (1812).

S. glauca, Palisot, Agrostogr. 51, t. 13 (1812) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,492 M.fr.VIII,110.
S. macrostachya, Humb., Bonpl. & Kunth, nov. gen. et spec. I, 110 (1815) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,493 M.fr.VIII,110;XI,129.
S. viridis, Palisot, Agrostogr. 51, (1812) ... ... W.A. S.A. — — — — N.A. B.fl.VII,494
S. verticillata, Palisot, Agrostogr. 51 (1812) ... ... W.A. — — — — — N.A. B.fl.VII,494 M.fr.VIII,110.

PENNISETUM, L. C. Richard in Persoon, synops. plant. I. 72 (1805). (Gymnothrix, Plagiosetum.)

P. compressum, R. Brown, prodr. 195 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,495 M.fr.VIII,109.
P. Arnhemicum, F. v. M., fragm. VIII, 109 (1874) ... ... — — — — — — N.A. B.fl.VII,496 M.fr.VIII,109.
P. refractum, F. v. M., fragm. VIII, 109 (1874) ... ... — S.A. — — — Q. N.A. B.fl.VII,495 M.fr.III,147;IX,194.

CENCHRUS, Linné, coroll. gen. 20 (1737).

C. australis, R. Brown, prodr. 196 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,497 M.fr.VIII,107;XI,129.
C. inflexus, R. Brown, prodr. 195 (1810) ... ... — — — — — — N.A. B.fl.VII,497 M.fr.VIII,107.
C. elymoides, F. v. M., fragm. VIII, 107 (1874) ... ... — — — — — Q. N.A. B.fl.VII,497 M.fr.VIII,107.

XEROCHLOA, R. Brown, prodr. fl. Nov. Holl. 196 (1810).

X. imberbis, R. Brown, prodr. 197 (1810) ... ... — — — — — — N.A. B.fl.VII,501 M.fr.VIII,117.
X. barbata, R. Brown, prodr. 197 (1810) ... ... — — — — — — N.A. B.fl.VII,501 M.fr.VIII,117.
X. laniflora, Bentham, Fl. Austr. VII, 502 (1878) ... ... — — — — — — N.A. B.fl.VII,502

THUAREA, Persoon, synops. plant. I, 110 (1805). (Thouarsea.)

T. sarmentosa, Persoon, synops. I, 110 (1805) ... ... — — — — — Q. N.A. B.fl.VII,502 M.fr.VI,86.

SPINIFEX, Linné, syst. nat. ed. XII, 757 (1767).

S. hirsutus, Labillard., Nov. Holl. pl. spec. II, 81, t. 230 (1806) W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,503 M.fr.VIII,138.
S. longifolius, R. Brown, prodr. 198 (1810) ... ... W.A. — — — — Q. N.A. B.fl.VII,504 M.fr.VIII,139. [IX,195.
S. paradoxus, Bentham in Hook. icon. plant. t. 1243 (1877) ... — S.A. — V. N.S.W. Q. — B.fl.VII,504 M.fr.VI,86;VIII,117,199;

LEPTASPIS, R. Brown, prodr. fl. Nov. Holl. 211 (1810).

L. Banksii, R. Brown, prodr. 211 (1810) ... ... — — — — — Q. — B.fl.VII,548 M.fr.VIII,116.

PEROTIS, Aiton, hort. Kewens. I, 85 (1789).

P. rara, R. Brown, prodr. 172 (1810) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,509 M.fr.VIII,115.

POLYPOGON, Desfontaines, flor. Atlantic. I, 66 (1798).

P. fugax, Nees in Steudel, syn. glum. I, 184 (1855) ... ... W.A. — — — — — — B.fl.VII,547
P. tenellus, R. Brown, prodr. 173 (1810) ... ... W.A. — — — — — — B.fl.VII,547 M.fr.VIII,114.

ARUNDINELLA, Raddi, agrostograph. Brasil. 37 (1823).

A. Nepalensis, Trinius, spec. gram. t. 268 (1833)... ... ... — — — — N.S.W. Q. N.A. B.fl.VII,545 M.fr.VIII,139.
A. Schultzii, Bentham, Fl. Austr. VII, 545 (1878) ... ... — — — — — — N.A. B.fl.VII,545

TRAGUS, Haller, hist. stirp. Helvet. n. 1413 (1768). (Lappago.)

T. racemosus, Haller, stirp. Helv. n. 1413 (1768)... ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,507 M.fr.VIII,107.

NEURACHNE, R. Brown, prodr. fl. Nov. Holl. 196 (1810). [129.

N. alopecuroides, R. Brown, prodr. 196 (1810) ... ... W.A. S.A. — V. N.S.W. — — B.fl.VII,507 M.fr.VI,86;VIII,200; XI,
N. Mitchelliana, Nees in Hook. Lond. Journ. of Bot. II, 410 (1843) — S.A. — V. N.S.W. — — B.fl.VII,508 M.fr.VI,86;VIII,200.
N. Munroi, F. v. M., fragm. VIII, 200 (1874) ... ... — S.A. — V. N.S.W. — — B.fl.VII,508 M.fr.V,204;XI,129.

ZOYSIA, Willdenow, Neue Schrift. nat. Freunde zu Berl. III, 440 (1801).

Z. pungens, Willdenow, Neue Schrift. nat. Fr. Berl. III, 440 (1801) — — T. V. N.S.W. Q. — B.fl.VII,506 M.fr.VIII,116.

IMPERATA, Cyrillo, plant. rar. Neap. II, t. 11 (1792).

I. arundinacea, Cyrillo, plant. rar. Neap. II, t. 11 (1792) ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,536 M.fr.VIII,126.

ERIANTHUS, L. C. Richard in Michaux, fl. bor. Amer. I, 54 (1803). (Pollinia, Eulalia, Pogonatherum, Leptatherum, Saccharum partly.)

E. articulatus, F. v. M., fragm. VIII, 118 (1874) ... ... — — — — — Q. N.A. B.fl.VII,525 M.fr.X,76.
E. irritans, Kunth, Graminées I, 160 (1832) ... ... — — — — — Q. N.A. B.fl.VII,526 M.fr.VIII,118.
E. fulvus, Kunth, Graminées I, 160 (1832) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.VII,526 M.fr.VIII,118.
E. Roxburghii, F. v. M., fragm. VIII, 117 (1874) ... ... — — — — — Q. — B.fl.VII,527 M.fr.VIII,117.
E. Mackinlayi; Pollinia, F. v. M. in Benth. fl. Austr. VII, 527 (1878) — — — — — — N.A. B.fl.VII,527 M.fr.VIII,118.

DIMERIA, R. Brown, prodr. fl. Nov. Holl. 204 (1810).

D. acinaciformis, R. Brown, prodr. 204 (1810) ... ... — — — — — Q. — B.fl.VII,523
D. tenera, Trinius in Mém. de l'Acad. de Petersb. six. sér. II, 225 (1833) ... ... — — — — — Q. N.A. B.fl.VII,523 M.fr.VII,104;XI,129.

ARTHRAXON, Palisot, Agrostograph. 111, t. XI, f. 6 (1812). (Batratherum.)

A. ciliare, Palisot, Agrostogr. 111, t. 11 (1812) ... ... — — — — N.S.W. Q. — B.fl.VII,524 M.fr.VIII,119;X,76.

ELIONURUS, Willdenow, spec. plant. IV, 941 (1805).

E. citreus, Munro in Benth. Fl. Austral. VII, 510 (1878) ... — — — — — Q. — B.fl.VII,510 M.fr.XI,129.

LEPTURUS, R. Brown, prodr. fl. Nov. Holl. 207 (1810).

L. incurvatus, Trinius, fundam. agrostogr. 123 (1820) ... ... — S.A. — V. N.S.W. — — B.fl.VII,668 M.fr.VIII,117.
L. cylindricus, Trinius, fundam. agrostogr. 123 (1820) ... ... W.A. S.A. — V. N.S.W. — — B.fl.VII,668
L. repens, R. Brown, prodr. 207 (1810) ... ... — — — — — Q. N.A. B.fl.VII,668 M.fr.VI,86.

ROTTBOELLIA, Linné, fil. gram. gen. 22 (1779). (Coelorrhachis.)

R. formosa, R. Brown, prodr. 206 (1810) ... ... — — — — — Q. N.A. B.fl.VII,513 M.fr.VI,85;XI,129.
R. exaltata, Linné, fil. suppl. 114 (1781) ... ... — — — — — Q. N.A. B.fl.VII,513
R. muricata, Retzius, observ. bot. III, 12 (1786)... ... ... — — — — — Q. N.A. B.fl.VII,514
R. ophiuroides, Bentham, Fl. Austr. VII, 514 (1878) ... .. — — — — — Q. N.A. B.fl.VII,514 M.fr.VIII,123.

OPHIURUS, K. F. Gaertner, de fruct. III, 3, t. 181 (1805). (Ophiurus.)

O. corymbosus, K. F. Gaertner, de fruct. III, 3, t. 181 (1805)... — — — — — Q. N.A. B.fl.VII,512 M.fr.VI,85.

MANISURIS, Linné, syst. nat. ed. duodec. 762 (1767).

M. granularis, Swartz, Fl. Ind. occid. I, 186 (1797) ... ... — — — — — Q. N.A. B.fl.VII,514 M.fr.VII,116;XI,129.

HEMARTHRIA, R. Brown, prodr. fl. Nov. Holl. 207 (1810).

H. compressa, R. Brown, prodr. 207 (1810) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,511 M.fr.XI,129.

ISCHAEMUM, Linné, syst. nat. 8 (1735); Linné, gen. pl. ed. sec. 525 (1742). (Hologamium, Spodiopogon partly.)

I. truncatiglume, F. v. M. in Benth. Fl. Austr. VII, 518 (1878) — — — — — — N.A. B.fl.VII,518
I. arundinaceum, F. v. M. in Benth. Fl. Austr. VII, 519 (1878) — — — — — — N.A. B.fl.VII,519
I. triticeum, R. Brown, prodr. 205 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,519

I. australe, R. Brown, prodr. 205 (1810) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,519
I. muticum, Linné, spec. plant. 1049 (1753) ... ... — — — — — Q. — B.fl.VII,520 M.fr.VIII,120.
I. ciliare, Retzius, observ. bot. VI, 36 (1791) ... ... — — — — N.S.W. — — B.fl.VII,520 M.fr.VIII,119.
I. decumbens, Bentham, Fl. Austr. VII, 521 (1878) ... ... — — — — — — N.A. B.fl.VII,521
I. pectinatum, Trinius in Mem. de l'Acad. de Petersb. six. sèr. II, 296 (1833)... ... ... — — — — N.S.W. Q. — B.fl.VII,521 M.fr.VIII,118.
I. fragile, R. Brown, prodr. 205 (1810) ... ... — — — — — Q. — B.fl.VII,522 M.fr.VIII,123;XI,130.
I. laxum, R. Brown, prodr. 205 (1810) ... ... — — — — — Q. N.A. B.fl.VII,522 M.fr.VIII,119.

ANDROPOGON, Royen, prodr. exhib. pl. hort. acad. Lugd. 52 (1740). (Sorghum, Chrysopogon, Heteropogon, Schizachyrium, Trachypogon, Holcus partly, Spodiopogon partly.)

A. erianthoides, F. v. M., fragm. X, 75 (1876) ... ... — — — — N.S.W. Q. — B.fl.VII,529 M.fr. X, 75.
A. sericeus, R. Brown, prodr. 201 (1810) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,529
A. affinis, R. Brown, prodr. 201 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,530
A. pertusus, Willdenow, spec. plant. IV, 922 (1805) ... ... — S.A. — V. N.S.W. Q. — B.fl.VII,530 M.fr.VIII,122;X,76.
A. annulatus, Forskael, Fl. Aegypt. Arab. 173 (1775) ... ... — S.A. — — — Q. N.A. B.fl.VII,531 M.fr.VIII,123.
A. Ischaemum, Linné, spec. plant. 1047 (1753) ... ... W.A. — — — — — B.fl.VII,531 M.fr.VIII,122.
A. intermedius, R. Brown, prodr. 202 (1810) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,531
A. procerus, R. Brown, prodr. 202 (1810) ... ... — — — — — — N.A. B.fl.VII,532 M.fr.VIII,124.
A. exaltatus, R. Brown, prodr. 202 (1810) ... ... W.A. S.A. — — — — N.A. B.fl.VII,532 M.fr.VIII,124.
A. bombycinus, R. Brown, prodr. 202 (1810) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,533 M.fr.XI,20.
A. Schoenanthus, Linné, spec. plant. 1046 (1753) .. .. — — — — — Q. — B.fl.VII,534 M.fr.VIII,124.
A. refractus, R. Brown, prodr. 202 (1810) ... ... — — — V. N.S.W. Q. N.A. B.fl.VII,534 M.fr.VIII,124;X,76.
A. lachnatherus, Bentham, Fl. Austr. VII, 534 (1874) ... ... — — — — N.S.W. Q. — B.fl.VII,534
A. fragilis, R. Brown, prodr. 202 (1810) ... ... — — — — — Q. N.A. B.fl.VII,535
A. contortus, Linné, spec. plant. 1045 (1753) ... ... W.A. — — — N.S.W. Q. N.A. B.fl.VII,517 M.fr.VIII,120.
A. triticeus, R. Brown, prodr. 201 (1810) ... ... — — — — — Q. N.A. B.fl.VII,517 M.fr.VIII,120.
A. Gryllus, Linné, amoen. acad. IV, 332 (1759) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,537 M.fr.VIII,121;XI,130.
A. montanus, Roxburgh, Flor. Indic. edit. Carey. I, 271 (1820) — — — V. N.S.W. Q. N.A. B.fl.VII,538 M.fr.VIII,122;X,76;XI,
A. acicularis, Retzius, observ. bot. V, 22 (1789) ... ... — — — — — Q. N.A. B.fl.VII,538 M.fr.VIII,122. [130.
A. elongatus, Sprengel, syst. I, 289 (1825)... ... ... — — — — — Q. N.A. B.fl.VII,539 M.fr.VIII,121.
A. Halepensis, Sibthorp & Smith, Fl. Graeca, I, 52, t. 68 (1806) W.A. — — — N.S.W. Q. — B.fl.VII,540 M.fr.VIII,119.
A. australis, Sprengel, syst. I, 287 (1825) ... ... — — — V. N.S.W. Q. N.A. B.fl.VII,540
A. tropicus, Sprengel, syst. I, 287 (1825) ... ... — — — — — Q. N.A. B.fl.VII,541
A. intrans; Sorghum, F. v. M. in Benth. Fl. Austr. VII, 541 (1878) — — — — — — N.A. B.fl.VII,541

ANTHISTIRIA, Linné, fil. gram. gen. 35 (1779). (Iseilema.)

A. ciliata, Linné fil., gram. gen. 35 (1779)... ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,542 M.fr.V,207;VIII,139.
A. frondosa, R. Brown, prodr. 200 (1810) ... ... — — — — — Q. N.A. B.fl.VII,542 M.fr.VIII,139;XI,130.
A. avenacea, F. v. M., fragm. V, 206 (1866) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,543 M.fr.VI,253;VIII,139.
A. membranacea, Lindley in Mitch. Trop. Austr. 88 (1848) ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,543 M.fr.V,207;VIII,139.

APLUDA, Linné, spec. plant. 82 (1753).

A. mutica, Linné, spec. plant. 82 (1753) ... ... — — — — N.S.W. Q. — B.fl.VII,544 M.fr.VIII,140.

CHIONACHNE, R. Brown in Horsfield's pl. Jav. rar. 15 (1838).

C. barbata, R. Brown in Bennett, pl. Javan. rar. 18 (1838) ... — — — — — Q. — B.fl.VII,515
C. cyathopoda, F. v. M. in Benth. Fl. Austr. VII, 516 (1878) ... — — — — — — N.A. B.fl.VII,515 M.fr.VIII,116.

ALOPECURUS, Linné, syst. nat. 8 (1735); Linné, gen. plant. 18 (1737). (Perhaps immigrated.)

A. geniculatus, Linné, spec. plant. 60 (1753) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,555 M.fr.VIII,138.

LEERSIA, Solander in Swartz, nov. gen. et spec. plant. 1 et 21 (1788). (Asprella, Asperella.)

L. hexandra, Swartz, nov. gen. et spec. pl. 21 (1788) ... ... — — — — N.S.W. Q. — B.fl.VII,549 M.fr.VIII,115.

ORYZA, Tournefort, inst. 513, t. 296 (1700), from Theophrastos & Dioscorides.

O. sativa, Linné, spec. plant. 333 (1753) ... ... — — — — — — N.A. B.fl.VII,550 M.fr.VIII,115;XI,130.

POTAMOPHILA, R. Brown, prodr. fl. Nov. Holl. 211 (1810).

P. parviflora, R. Brown, prodr. 211 (1810)... ... ... — — — — N.S.W. — — B.fl.VII,550 M.fr.VIII,126.

EHRHARTA, Thunberg in Svensk. Ventsk. Acad. Handl. 216, t. 8 (1779). (Tetrarrhena, Michrolaena, Diplax.)

E. distichophylla, Labill., Nov. Holl. pl. spec. I, 90, t. 117 (1804) — — T. V. — — — B.fl.VII,554 M.fr.VII,90.
E. juncea, Sprengel, syst. II, 114 (1825) ... ... — — T. V. N.S.W. — — B.fl.VII,554 M.fr.VII,90.
E. laevis, Sprengel, syst. II, 115 (1825) ... ... W.A. — — — — — B.fl.VII,554 M.fr.VII,90.
E. acuminata, Sprengel, syst. II, 114 (1825) ... ... — — T. V. — — — B.fl.VII,555 M.fr.VII,90;XI,130.
E. stipoides, Labillard., Nov. Holl. pl. spec. I, 91, t. 118 (1804) W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,552 M.fr.VII,90.
E. diarrhena, F. v. M., fragm. VII, 89 (1870) ... . — — T. — — — B.fl.VII,553 M.fr.VII,89.

HIEROCHLOE, J. G. Gmelin, Fl. Sibiric. I, 100 (1747). (Hierochloa, Disarrhenum.)

H. redolens, R. Brown, prodr. 209 (1810) ... ... — — T. V. N.S.W. — — B.fl.VII,558 M.fr.VIII,137.
H. rariflora, J. Hooker, Fl. Austr. I, 93 (1844) ... ... — — T. V. N.S.W. — — B.fl.VII,559 M.fr.VIII,138.

ARISTIDA, Linné, spec. plant. 82 (1753). (Chaetaria, Arthratherum.)

A. hygrometrica, R. Brown, prodr. 174 (1810) ... ... — — — — — — N.A. B.fl.VII,561
A. stipoides, R. Brown, prodr. 174 (1810) .. .. W.A. S.A. — — N.S.W. Q. N.A. B.fl.VII,561 M.fr.VIII,111.
A. arenaria, Gaudichaud in Freyc. voy. Bot. 407 (1826)... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,561
A. Behriana, F. v. M. in Transact. Vict. Inst. 44 (1854) ... — S.A. — V. N.S.W. Q. — B.fl.VII,562 M.fr.XI,130.
A. leptopoda, Bentham, Fl. Austr. VII, 562 (1878) ... ... — S.A. — V. N.S.W. Q. — B.fl.VII,562 M.fr.XI,130.
A. vagans, Cavanilles, icon. et descr. pl. V, 45, t. 471 (1799) ... — — — — N.S.W. Q. — B.fl.VII,562 M.fr.VIII,111.
A. ramosa, R. Brown, prodr. 173 (1810) ... ... — S.A. — — N.S.W. Q. — B.fl.VII,563
A. calycina, R. Brown, prodr. 173 (1810) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,563 M.fr.XI,130.
A. depressa, Retzius, observ. bot. IV, 22 (1787) ... ... — — — — N.S.W. Q. — B.fl.VII,563

STIPA, Linné, spec. plant. 78 (1737). (Streptachne, Urachne partly.)

S. elegantissima, Labillardière, Nov. Holl. pl. spec. I, 23, t. 29 (1804) W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,565 M.fr.VIII,103.
S. Tuckeri, F. v. M., fragm. XI, 128 (1881) ... ... — — — V. N.S.W. — — — M.fr.XI,129.
S. verticillata, Nees in Spreng. syst. veg. cur. post. 30 (1827) ... — — — — N.S.W. Q. — B.fl.VII,566 M.fr.VIII,105.

S. flavescens, Labillardière, Nov. Holl. pl. spec. I, 24, t. 30 (1804) W.A. S.A. T. V. N.S.W. — — B.fl.VII,566 M.fr.VIII,104.
S. teretifolia, Steudel, syn. glum. I, 128 (1855) ... ... W.A. — T. V. — — — B.fl.VII,567 M.fr.VIII,104.
S. compressa, R. Brown, prodr. 175 (1810)... ... ... W.A. — — — — — — B.fl.VII,567
S. Drummondii, Steudel, syn. glum. I, 128 (1855) ... ... W.A. — — — — — — B.fl.VII,567
S. pycnostachya, Bentham, Fl. Austr. VII, 568 (1878) ... ... W.A. — — — — — — B.fl.VII,568
S. setacea, R. Brown, prodr. 174 (1810) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,568
S. semibarbata, R. Brown, prodr. 174 (1810) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,568 M.fr.VIII,104;XI,129.
S. hemipogon, Bentham, Fl. Austr. VII, 569 (1878) ... ... W.A. — — — — — B.fl.VII,569 M.fr.VIII,104.
S. pubescens, R. Brown, prodr. 174 (1810)... ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,569 M.fr.VIII,104.
S. aristiglumis, F. v. M. in Transact. Vict. Inst. 43 (1854) ... — S.A. — V. N.S.W. Q. — B.fl.VII,570 M.fr.VIII,103.
S. Eriopus, Bentham, Fl. Austr. VII, 570 (1878)... ... ... W.A. — — — — — — B.fl.VII,570
S. trichophylla, Bentham, Fl. Austr. VII, 570 (1878) ... ... W.A. — — — — — — B.fl.VII,570
S. scabra, Lindley in Mitch. Trop. Austr. 31 (1848) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,570
S. Streptachne, F. v. M. in journ. R.S. of N.S.W. 237 (1881) ... — — — — — Q. — B.fl.VII,572

DICHELACHNE, Endlicher, prodr. fl. Norfolk. 20 (1833). (Muehlenbergia partly.)
D. crinita, J. Hooker, Fl. N. Zel. I, 293 (1853) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,574 M.fr.VIII,105.
D. sciurea, J. Hooker, Fl. N. Zel. I, 294 (1853) ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,574 M.fr.VIII,105;XI,129.

PENTAPOGON, R. Brown, prodr. fl. Nov. Holl. 173 (1810).
P. Billardieri, R. Brown, prodr. 173 (1810)... ... ... — S.A. T. V. N.S.W. — — B.fl.VII,572 M.fr.VIII,106.

ECHINOPOGON, Palisot, Essai d'une nouv. Agrostogr. 42, t. 9, f. 5 (1812). (Cinna partly.)
E. ovatus, Palisot, Agrostogr. 42, t. 9 (1812) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,599 M.fr.VIII,106.

DIPLOPOGON, R. Brown, prodr. fl. Nov. Holl. 176 (1810). (Dipogonia.)
D. setaceus, R. Brown, prodr. 176 (1810) ... ... W.A. — — — — — — B.fl.VII,573 M.fr.VI,86;VIII,106.

AMPHIPOGON, R. Brown, prodr. 175 (1810). (Aegopogon partly, Pentacraspedion, Gamelythrum.)
A. debilis, R. Brown, prodr. 175 (1810) ... ... W.A. — — — — — — B.fl.VII,597 M.fr.VIII,201.
A. strictus, R. Brown, prodr. 175 (1810) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,597 M.fr.VIII,201.
A. laguroides, R. Brown, prodr. 175 (1810) ... ... W.A. — — — — — — B.fl.VII,598 M.fr.VI,86;VIII,201.
A. cygnorum, Nees in Lehm. pl. Preiss. II, 100 (1846) ... ... W.A. — — — — — — B.fl.VII,599
A. turbinatus, R. Brown, prodr. 175 (1810) ... ... W.A. — — — — — — B.fl.VII,599 M.fr.VI,86;VIII,201.

PAPPOPHORUM, Schreber, gen. pl. II, 787 (1791). (Euneapogon).
P. commune, F. v. M., plants of Gregory's Exp. 10 (1859) ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,600 M.fr.VIII,200.

SPOROBOLUS, R. Brown, prodr. fl. Nov. Holl. 169 (1810). (Vilfa partly.)
S. Virginicus, Humboldt & Kunth, gram. I, 67 (1832) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,621 M.fr.VI,84.
S. Indicus, R. Brown, prodr. 170 (1810) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,622 M.fr.VI,84;X,76.
S. diander, Palisot, Agrostogr. 26 (1812) ... ... — — — — N.S.W. Q. — B.fl.VII,622 M.fr.VI,84;VIII,140.
S. pulchellus, R. Brown, prodr. 170 (1810)... ... ... — — — — N.S.W. Q. N.A. B.fl.VII,623 M.fr.VIII,140.
S. Lindleyi, Bentham, Fl. Austr. VII, 623 (1878)... ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,623 M.fr.XI,130. [130.
S. actinocladus, F. v. M., fragm. VIII, 140 (1874) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,623 M.fr.VI,84;IX,194;XI,

AGROSTIS, Linné, syst. nat. 8 (1735); Linné, gen. plant. 19 (1737). (Deyeuxia, Didymochaeta, Trichodium, Bromidium, Lachnagrostis partly).
A. Muelleri, Bentham, Fl. Austr. VII, 576 (1878) ... ... — — — V. N.S.W. — — B.fl.VII,576 M.fr.VI,86.
A. scabra, Willdenow, spec. plant. I, 378 (1797) ... ... — S.A. T. V. N.S.W. — — B.fl.VII,576
A. venusta, Trinius in Mém. de l'Acad. de Petersb. VI, 340 (1850) W.A. S.A. T. V. N.S.W. — — B.fl.VII,576
A. aequata, Nees in Hook. Lond. Journ. of Bot. II, 412 (1843)... — — T. — — — — B.fl.VII,579
A. Solandri, F. v. M., Veget. of the Chath. Isl. 60 (1864) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,579 M.fr.XI,27,130.
A. montana, R. Brown, prodr. 171 (1810) ... ... — S.A. T. V. N.S.W. — — B.fl.VII,581
A. quadriseta, R. Brown, prodr. 171 (1810) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,581
A. cylindrica, R. Brown, prodr. 171 (1810)... ... ... W.A. — — — — — — B.fl.VII,582
A. densa, F. v. M.; Deyeuxia, Benth. Fl. Austr. VII, 582 (1878) — S.A. — V. — — — B.fl.VII,582
A. frigida, F. v. M. in Benth. Fl. Austr. VII, 583 (1878) ... — — T. V. N.S.W. — — B.fl.VII,583
A. scabra, R. Brown, prodr. 172 (1810) ... ... — — T. V. N.S.W. Q. — B.fl.VII,583 M.fr.VI,85.
A. nivalis, F. v. M. in Transact. Vict. Inst. 43 (1854) ... ... — — — V. N.S.W. — — B.fl.VII,584
A. Gunniana, F. v. M.; Deyeuxia, Benth. Fl. Austr. VII, 584 (1878) — — T. — — — — B.fl.VII,584
A. breviglumis, F.v.M.; Deyeuxia, Benth. Fl. Austr.VII, 584 (1878) — — — — N.S.W. — — B.fl.VII,584

COELACHNE, R. Brown, prodr. fl. Nov. Holl. 187 (1810).
C. pulchella, R. Brown, prodr. 187 (1810) ... ... — — — — — Q. — B.fl.VII,626

MICRAIRA, F. v. M., fragm. V, 208 (1866).
M. subulifolia, F. v. M., fragm. V, 208 (1866) ... ... — — — — — Q. — — M.fr.V,208.

AIRA, Linné, gen. plant. 335 (1737). (Aera, Deschampsia.)
A. caespitosa, Linné, spec. plant. 64 (1753) ... ... — S.A. T. V. N.S.W. — — B.fl.VII,587 M.fr.VIII,138.

TRISETUM, Persoon, synops. plant. I, 97 (1805).
T. subspicatum, Palisot, Agrostogr. 88 (1812) ... ... — — T. V. N.S.W. — — B.fl.VII,588 M.fr.VIII,136.

ERIACHNE, R. Brown, prodr. fl. Nov. Holl. 183 (1810).
E. stipacea, F. v. M., fragm. V, 206 (1866) ... ... — — — — — Q. N.A. B.fl.VII,626 M.fr.VIII,137;X,76.
E. Armitii, F. v. M. in Benth. Fl. Austr. VII, 627 (1878) ... — — — — — Q. N.A. B.fl.VII,627
E. squarrosa, R. Brown, prodr. 183 (1810)... ... ... — — — — — Q. N.A. B.fl.VII,628
E. glauca, R. Brown, prodr. 184 (1810) ... ... — — — — — — N.A. B.fl.VII,628
E. rara, R. Brown, prodr. 183 (1810) ... ... — — — — — Q. — B.fl.VII,628
E. agrostidea, F. v. M., fragm. VII, 82 (1870) ... ... — — — — — — N.A. B.fl.VII,628 M.fr.VIII,137.
E. ciliata, R. Brown, prodr. 184 (1810) ... ... — — — — — Q. N.A. B.fl.VII,629
E. setacea, Bentham, Fl. Austr. VII, 629 (1878) ... ... — — — — — — N.A. B.fl.VII,629
E. avenacea, R. Brown, prodr. 184 (1810) ... ... — — — — — — N.A. B.fl.VII,629
E. aristidea, F. v. M., fragm. V, 205 (1866) ... ... W.A. S.A. — — N.S.W. Q. — B.fl.VII,629 M.fr.V,205.
E. pallescens, R. Brown, prodr. 184 (1810) ... ... — — — — — Q. — B.fl.VII,630
E. festucacea, F. v. M., fragm. V, 205 (1866) ... ... — — — — — — N.A. B.fl.VII,630 M.fr.V,205.

E. ovata, Nees in Hook. Lond. Journ. II, 418 (1843) ... ... W.A. S.A. — — – — N.A. B.fl.VII,630 M.fr.VIII,137.
E. meliacea, F. v. M., fragm. V, 205 (1866) ... ... — — — — — — N.A. B.fl.VII,631 M.fr.V,205.
E. pallida, F. v. M. in Benth. Fl. Austr. VII, 631 (1878) ... — — — — — — N.A. B.fl.VII,631
E. scleranthoides, F. v. M., fragm. VIII, 233 (1874) ... ... — S.A. — — — — — B.fl.VII,631
E. mucronata, R. Brown, prodr. 184 (1810) ... ... — — — — — Q. — B.fl.VII,632
E. obtusa, R. Brown, prodr. 184 (1810) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,632 M.fr.VIII,137;XI,130.
E. capillaris, R. Brown, prodr. 184 (1810)... ... ... — — — — — — N.A. B.fl.VII,632

ANISOPOGON, R. Brown, prodr. fl. Nov. Holl. 176 (1810).
A. avenaceus, R. Brown, prodr. 176 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,590 M.fr.VIII,106.

DANTHONIA, De Candolle, Fl. franç. III, 32 (1805). (Amphibromus, Monacather, Plinthacanthus.)
D. paradoxa, R. Brown, prodr. 177 (1810) ... ... — — — — N.S.W. — — B.fl.VII,591
D. bipartita, F. v. M., fragm. I, 160 (1859) ... ... W.A. S.A. — V. N.S.W. — — B.fl.VII,592 M.fr.VIII,136.
D. carphoides, F. v. M. in Benth. Fl. Austr. VII, 592 (1878) ... — S.A. — V. N.S.W. — — B.fl.VII,592 M.fr.XI,130.
D. penicillata, F. v. M., fragm. VIII, 135 (1873)... ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,592 M.fr.VIII,135.
D. robusta, F. v. M. in Transact. Vict. Inst. 44 (1854) ... ... — — — V. N.S.W. — — B.fl.VII,593 M.fr.VIII,136.
D. pauciflora, R. Brown, prodr. 177 (1810) ... ... — — T. V. N.S.W. — — B.fl.VII,596
D. nervosa, J. Hooker, Fl. Tasman. II, 121, t. 163 (1860) ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,589 M.fr.VIII,135.

ASTREBLA, F. v. M., fragm. phytogr. Austr. X, 76 (1876).
A. pectinata, F. v. M., fragm. X, 76 (1876) ... . — S.A. — — N.S.W. Q. N.A. B.fl.VII,602 M.fr.VIII,134.
A. triticoides, F. v. M., fragm. X, 76 (1876) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,602 M.fr.VIII,134.

MICROCHLOA, R. Brown, prodr. fl. Nov. Holl. 208 (1810).
M. setacea, R. Brown, prodr. 208 (1810) .. .. — — — — — — N.A. B.fl.VII,608

CYNODON, L. C. Richard in Persoon, synops. I, 85 (1805).
C. Dactylon, L. C. Richard in Persoon, synops. I, 85 (1805) ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,609 M.fr.VIII,113.
C. tenellus, R. Brown, prodr. 187 (1810) ... ... — — — — — — N.A. B.fl.VII,609 M.fr.VIII,113.
C. convergens, F. v. M., fragm. VIII, 113 (1874)... ... ... — — — — — Q. N.A. B.fl.VII,610 M.fr.X,76.
C. ciliaris, Bentham, Fl. Austr. VII, 610 (1878) ... ... — S.A. — — — — — B.fl.VII,610

CHLORIS, Swartz, nov. gen. et spec. plant. 25 (1788).
C. unispicea, F. v. M., fragm. VII, 118 (1871) ... ... — — — — — Q. — B.fl.VII,611 M.fr.
C. pumilio, R. Brown, prodr. 186 (1810) ... ... — — — — — Q. N.A. B.fl.VII,611
C. pectinata, Bentham, Fl. Austr. VII, 612 (1878) ... ... — S.A. — — — Q. N.A. B.fl.VII,612
C. divaricata, R. Brown, prodr. 186 (1810) ... ... — — — — — Q. N.A. B.fl.VII,612 M.fr.VI,85.
C. acicularis, Lindley in Mitch. Trop. Austr. 33 (1848) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,612 M.fr.VI,85.
C. truncata, R. Brown, prodr. 186 (1810) ... ... — S.A. — V. N.S.W. Q. — B.fl.VII,612 M.fr.VI,85;XI,130.
C. ventricosa, R. Brown, prodr. 186 (1810) .. .. — — — — N.S.W. Q. — B.fl.VII,613 M.fr.VI,85.
C. barbata, Swartz, nov. gen. et spec. plant. 25 (1788) ... ... — S.A. — — — — N.A. B.fl.VII,613 M.fr.VI,85.
C. scariosa, F. v. M., fragm. VI, 85 (1868) ... ... — — — — — Q. N.A. B.fl.VII,614 M.fr.XI,130.

ELEUSINE, J. Gaertner, de fruct. I, 7 (1788). (Leptochloa, Dactyloctenium, Acrachne).
E. Aegyptiaca, Persoon, syn. I, 82 (1805) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,615 M.fr.VIII,111.
E. Indica, J. Gaertner, de fruct. I, 7 (1788) ... ... — — — — N.S.W. Q. — B.fl.VII,615 M.fr.VIII,112.
E. verticillata, Roxburgh, Fl. Ind. ed. Car. & Wall. I, 346 (1820) — — — — — Q. N.A. B.fl.VII,616 M.fr.VIII,112. [130.
E. digitata, Sprengel, syst. cur. post. 36 (1827) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.VII,617 M.fr.I,216;VIII,112;XI,
E. Chinensis, F. v. M.; Poa, Koenig in Roth, nov. pl. spec. 65 (1821) — — — — N.S.W. Q. — B.fl.VII,617 M.fr.VIII,132.
E. digitata, Sprengel, syst. veg. cur. post. 36 (1827) ... ... — — — — — Q. N.A. B.fl.VII,617 M.fr.VIII,113.

CENTOTHECA, Desvaux in Palisot, Agrost. 69, t. 14 (1812). [76.
C. lappacea, Desvaux in Palisot, Agrost. 69, t. 14 (1812) ... — — — — — Q. — B.fl.VII,640 M.fr.VI,85;VIII,116;X,

POA, Linné, gen. plant. 20 (1737). (Glyceria partly.)
P. Billardieri, Steudel, syn. glum. I, 262 (1855) ... ... W.A. S.A. T. V. — — N.A. B.fl.VII,651
P. homomalla, Nees in Lehm. pl. Preiss. II, 104 (1846) ... ... W.A. — — — — — — B.fl.VII,651
P. caespitosa, G. Forster, fl. ins. Austr. prodr. 89 (1786) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,651 M.fr.VIII,133;XI,13.
P. Maxwelli, Bentham, Fl. Austr. VII, 653 (1878) ... ... W.A. — — — — — — B.fl.VII,652
P. nodosa, Nees in Lehm. pl. Preiss. II, 105 (1846) ... ... W.A. S.A. — V. N.S.W. — — B.fl.VII,653 M.fr.VIII,132.
P. saxicola, R. Brown, prodr. 654 (1810) ... ... — — T. — — - B.fl.VII,654
P. lepida, F. v. M., fragm. VIII, 130 (1874) ... ... W.A. S.A. — V. N.S.W. - -- B.fl.VII,654 M.fr.XI,130.
P. Fordeana, F. v. M., fragm. VIII, 130 (1874) ... ... — S.A. — V. N.S.W. — — B.fl.VII,657 M.fr.VIII,130.
P. fluitans, Scopoli, flor. Carniol. 106 (1772) ... ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,657 M.fr.VIII,129.
P. latispicea; Festuca, F. v. M., VIII, 127 (1873) ... ... — — — — N.S.W. — — B.fl.VII,658 M.fr.VIII,128.
P. syrtica, F. v. M. in Transact. Vict. Inst. 45 (1854) ... ... W.A. S.A. T. V. — -- — B.fl.VII,658 M.fr.VIII,130. [130.
P. dives; Festuca, F. v. M., fragm. III, 147 (1863) ... ... — — — V. — — — B.fl.VII,659 M.fr.VI,147;VIII,129,XI,
P. ramigera, F. v. M. in Transact. Vict. Inst. 45 (1854)... ... W.A. S.A. — V. N.S.W. — — B.fl.VII,659 M.fr.VI,85;VIII,131,XI, 130.

FESTUCA, Dillenius, nov. gen. 90, t. 3 (1719). (Glyceria partly, Vulpia partly, Schedonorus, Schenodorus, Brizopyrum.)
F. duriuscula, Linné, spec. plant. 74 (1753) ... ... — S.A. T. V. N.S.W. — — B.fl.VII,663 M.fr.VI,85.
F. scirpoidea, F. v. M., fragm. VIII, 129 (1874) ... ... W.A. — — — — — — B.fl.VII,655 M.fr.XI,130.
F. litoralis, Labillardière, Nov. Holl. spec. plant. I, 22, t. 27 (1804) W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,656 M.fr.VIII,129.
F. Hookeriana, F. v. M. in J. Hooker, Fl. Tasm. II, 122, t. 115 (1860) — — T. V. N.S.W. — — B.fl.VII,656 M.fr.VIII,131.

DIPLACHNE, Palisot, Essai d'une nouv. Agrostogr. 9, t. 16 (1812). (Uralapis.)
D. loliiformis, F. v. M. in Benth. fl. Austr. VII, 618 (1878) ... — S.A. — V. N.S.W. Q. — B.fl.VII,616 M.fr.VIII,128;XI,130.
D. Muelleri, Bentham, Fl. Austr. VII, 619 (1878) ... ... — S.A. — — — — N.A. B.fl.VII,619
D. fusca, Palisot, Agrost. 163 (1812) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,619 M.fr.VIII,127.
D. parviflora, Bentham, Fl. Austr. VII, 620 (1878) ... ... — — — — — — N.A. B.fl.VII,620 M.fr.VIII,129,

TRIODIA, R. Brown, prodr. fl. Nov. Holl. 182 (1810).

T. Mitchelli, Bentham, Fl. Austr. VII, 606 (1878)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,606		
T. pungens, R. Brown, prodr. 182 (1810)	...	...	...	...	—	S.A.	—	—	—	—	N.A.	B.fl.VII,606 M.fr.VIII,129.
T. Cunninghamii, Bentham, Fl. Austr. VII, 606 (1878)	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,606		
T. irritans, R. Brown, prodr. 182 (1810)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,607 M.fr.VIII,129.
T. procera, R. Brown, prodr. 182 (1810)	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.VII,607
T. microstachya, R. Brown, prodr. 182 (1810)	...	...	...		—	—	—	—	—	—	N.A.	B.fl.VII,607
T. microdon, F. v. M.; Triraphis, Benth. Fl. Austr. VII, 605 (1878)			—	—	—	—	N.S.W.	—	—	B.fl.VII,605		

DISTICHLIS, Rafinesque in Journ. de Phys. LXXXIX, 104 (1819). (Uniola partly, Brizopyrum partly.)

D. maritima, Rafinesque in Journ. de Phys. LXXXIX, 104 (1819)			—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,637		M.fr.VIII,129.

BROMUS, Dillenius in Linné, syst. nat. 8 (1735); Linné, gen. plant. 15 (1737).

B. arenarius, Labillardière, Nov. Holl. pl. spec. I, 23, t. 28 (1804)			W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,660		M.fr.VI,126;XI,20.

ERAGROSTIS, Palisot, Essai d'une nouv. Agrostogr. 70, t. 14 (1812). (Poa partly).

E. tenella, Palisot in Roem. et Schult. syst. veg. II, 576 (1817)			—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,643		M.fr.VIII,132;XI,130
E. nigra, Nees in Steud. nomencl. bot. I, 563 (1841)	...	...	—	—	—	—	N.S.W.	—	—	B.fl.VII,643		
E. imbecilla, Bentham, Fl. Austr. VII, 643 (1878)		...	—	—	—	—	—	Q.	—	B.fl.VII,643		
E. trichophylla, Bentham, Fl. Austr. VII, 644 (1878)	...	...	—	S.A.	—	—	—	—	—	B.fl.VII,644		
E. leptocarpa, Bentham, Fl. Austr. VII, 644 (1878)	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.VII,644		
E. megalosperma, F. v. M. in Benth. Fl. Austr. VII, 644 (1878)			—	—	—	—	N.S.W.	Q.	—	B.fl.VII,644		
E. pilosa, Palisot, Agrost. 71 (1812)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,645 M.fr.VIII,133;XI,130
E. leptostachya, Steudel, syn. glum. I, 279 (1855)		...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,645	
E. Schultzii, Bentham, Fl. Austr. VII, 646 (1878)		...	...	—	—	—	—	—	—	N.A.	B.fl.VII,646	
E. diandra, Steudel, syn. glum. I, 279 (1855)	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,646	M.fr.XI,130.
E. Brownii, Nees in Steud. nomencl. bot. I, 562 (1841)		...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,646	
E. concinna, Steudel, syn. glum. I, 279 (1855)	...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.VII,647	
E. speciosa, Steudel, syn. glum. I, 279 (1855)	...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.VII,648	
E. laniflora, Bentham, Fl. Austr. VII, 648 (1878)		...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VII,648	
E. eriopoda, Bentham, Fl. Austr. VII, 648 (1878)	...	...	—	S.A.	—	—	N.S.W.	—	N.A.	B.fl.VII,648		
E. setifolia, Nees in Hook. Lond. journ. II, 419 (1843)	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,648		
E. lacunaria, F. v. M., first gen. Rep. 20 (1853)	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,649	
E. falcata, Gaudichaud in Voy. Freyc. Bot. 408, t. 25 (1826)	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,649			
E. stenostachya, Steudel, syn. glum. I, 279 (1855)	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,650		

HETERACHNE, Bentham in J. Hook. icon. plant. XIII, 39 (1877).

H. Brownii, Bentham in J. Hook. icon. XIII, 40 (1877)	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,635		M.fr.VIII,132.
H. Gulliveri, Bentham in J. Hook. icon. XIII, 39, t. 1250 (1877)			—	—	—	—	—	Q.	N.A.	B.fl.VII,635		

ECTROSIA, R. Brown, prodr. fl. Nov. Holl. 185 (1810).

E. Schultzii, Bentham, Fl. Austr. VII, 633 (1878)	...	...	—	—	—	—	—	—	N.A.	B.fl.VII,633		
E. leporina, R. Brown, prodr. 186 (1810)	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,633 M.fr.VIII,109.
E. agrostoides, Bentham, Fl. Austr. VII, 634 (1878)	...	...	—	—	—	—	—	—	N.A.	B.fl.VII,634		
E. Gulliveri, F. v. M., fragm. VIII, 201 (1874)	...	...	...	—	—	—	—	Q.	N.A.	B.fl.VII,634		M.fr.VIII,201.

ELYTHROPHORUS, Palisot, Essai d'une nouv. Agrostogr. 67, t. 14 (1812).

E. articulatus, Palisot, Agrost. 67, t. 14 (1812)	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,638	M.fr.VIII,109.

TRIRHAPHIS, R. Brown, prodr. fl. Nov. Holl. 185 (1810). (Triraphis.)

T. mollis, R. Brown, prodr. 185 (1810)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,603 M.fr.VIII,108.
T. pungens, R. Brown, prodr. 185 (1810)	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.VII,604 M.fr.VIII,108,125.
T. bromoides, F. v. M., fragm. VIII, 108 (1874)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,604	M.fr.VIII,108.
T. danthonioides, F. v. M., fragm. VIII, 125 (1874)	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,604		M.fr.VIII,108.

AGROPYRON, J. Gaertner, nov. comm. Petrop. XIV, 539 (1770). (Agropyrum, Triticum partly, Anthosachne, Vulpia partly.)

A. scabrum, Palisot, Agrost. 102 (1812)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,665 M.fr.VI,85;XI,27,130.
A. velutinum, Nees in Hook. Lond. Journ. of Bot. II, 417 (1843)			—	—	T.	V.	N.S.W.	—	—	B.fl.VII,665		
A. pectinatum, Palisot, Agrost. 102 (1812)	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,666 M.fr.VI,85.

ARUNDO, Tournefort, inst. 526 (1700), from Varro. (Phragmites.)

A. Phragmites, Dodoens, stirp. hist. pemptad. 602 (1583)	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,637			
A. Roxburghii, F. v. M.; Phragmites, Kunth in nov. act. acad. Caes. XIX, suppl. I, 152 (1843)	...	...	...	...	—	—	—	—	—	Q.	N.A.	— M.fr.XI,139.

BAMBUSA, Schreber, gener. plantar. I, 236 (1789).

Species (three, of this or allied genera, not yet found in flower) undetermined	...	...	...	...	...	...	—	—	—	—	—	Q. N.A. — M.fr.VI,85;XI,129.

ACOTYLEDONEAE.

A. L. de Jussieu, genera plantarum 1 (1789).

ACOTYLEDONEAE VASCULARES.

Meissner, plantarum vascularium genera, 429 (1843).

RHIZOSPERMAE.

G. Weber, primit. Fl. Holsat. 74 (1780).

AZOLLA, Lamarck, Encycl. méthod. I, 343 (1783).

A. pinnata, R. Brown, prodr. 167 (1810) ... ... — S.A. — V. N.S.W. Q. — B.fl.VII,679 M.fr.V,140.
A. filiculoides, Lamarck, Encycl. méth. I, 343 (1783) ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,680 M.fr.V,140;XI,127.

MARSILEA, Linné, syst. nat. 8 (1735); Linné, gen. plant. 326 (1737). (Marsiglia.)

M. quadrifolia, Linné, spec. plant. 1099 (1753) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,683 M.fr.V,140.

PILULARIA, Vaillant, Bot. Paris. 159, t. 15 (1727).

P. globulifera, Linné, spec. plant. 1100 (1753) ... ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,684 M.fr.V,140.

ISOETES, Linné, skanska resa 420 (1751).

I. Drummondii, A. Braun in Berl. Monatsb. 593 (1863) ... ... W.A. — — V. — — — B.fl.VII,672 M.fr.V,140.
I. elatior, F. v. M. & A. Braun in Linnaea XXV, 722 (1852) ... — — T. — — — — — M.fr.V,140;XI,139
I. humilior, F. v. M. & A. Braun in Linnaea XXV, 722 (1852)... — — T. — — — — — M.fr.V,140;XI,139.
I. Gunnii, A. Braun in Berl. Monatsb. 535 (1868) ... ... — — T. — — — — — M.fr.XI,139.
I. Muelleri, A. Braun in Berl. Monatsb. 535 (1868) ... ... — — — — — Q. — — M.fr.XI,139.

LYCOPODINEAE.

Swartz, synops. filic. XV (1806).

PSILOTUM, Swartz in Schrader, Journ. fuer die Bot. II, 109 (1800.)

P. triquetrum, Swartz, synops. filic. 187 (1806) ... ... — — — — N.S.W. Q. — B.fl.VII,681 M.fr.V,112;X,118.
P. complanatum, Swartz, synops. filic. 187 (1806) ... ... — — — — — Q. — B.fl.VII,682 M.fr.V,112.

TMESIPTERIS, Bernhardi in Schrader, Journ. fuer die Bot. II, 131 (1800).

T. Tannensis, Bernhardi in Schrad. Journ. II, 131, t. 2 (1800)... — — T. V. N.S.W. Q. — B.fl.VII,680 M.fr.V,112;X,118.

LYCOPODIUM, Ruppius, Fl. Jenens. 32 (1718), from Tabernaemontanus (1588).

L. Selago, Linné, spec. plant. 1102 (1753) ... ... — — T. V. N.S.W. — — B.fl.VII,674 M.fr.V,111;X,118.
L. Phlegmaria, Linné, spec. plant. 1101 (1753) ... ... — — — — — Q. — B.fl.VII,674 M.fr.V,111;X,118.
L. clavatum, Linné, spec. plant. 1101 (1753) ... ... — — T. V. N.S.W. — — B.fl.VII,675 M.fr.V,111.
L. Carolinianum, Linné, spec. plant. 1104 (1753)... ... ... W.A. — T. V. N.S.W. Q. — B.fl.VII,675 M.fr.V,111;X,118.
L. laterale, R. Brown, prodr. 165 (1810) ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,675 M.fr.V,111;VI,124;X,118
L. cernuum, Linné, spec. plant. 1103 (1753) ... ... — — — — — Q. N.A. B.fl.VII,676 M.fr.V,111;X,118.
L. densum, Labill., Nov. Holl. plant. spec. II, 104, t. 251 (1806) — S.A. T. V. N.S.W. — — B.fl.VII,676 M.fr.V,111;X,118.
L. scariosum, G. Forster, florul. ins. austr. prodr. 87 (1786) ... — — T. V. — — — B.fl.VII,676 M.fr.V,112.
L. volubile, G. Forster, florul. ins. austr. prodr. 86 (1786) ... — — — — — — N.A. B.fl.VII,677 M.fr.V,112.

SELAGINELLA, Palisot, prodr. des fam. de l'Aethéogamie 101 (1805). (Lycopodium partly.)

S. Preissiana, Spring in Mém. de l'Ac. de Brux. 61 (1842) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,677 M.fr.V,112;VIII,275.
S. uliginosa, Spring in Mém. de l'Ac. de Brux. II, 60 (1842) ... — S.A. T. V. N.S.W. Q. — B.fl.VII,678 M.fr.V,112;VIII,275.
S. flabellata, Spring in Mém. de l'Ac. de Brux. 174 (1849) ... — — — — — Q. — B.fl.VII,678 M.fr.VIII,175,275.
S. concinna, Spring in Mém. de l'Ac. de Brux. 199 (1849) ... — — — — — Q. — B.fl.VII,678 M.fr.V,112;VIII,274.
S. Belangeri, Spring in Mém. de l'Ac. de Brux. 242 (1849) ... — — — — N.S.W. Q. N.A. B.fl.VII,679

PHYLLOGLOSSUM, Kunze in Mohl & Schlechtendal, Bot. Zeitung, 721 (1843).

P. Drummondii, Kunze in der Bot. Zeit. 721 (1843) ... ... W.A. — T. V. N.S.W. — — B.fl.VII,672 M.fr.V,112;X,118.

FILICES.

Linné, gen. plant. 322 (1737).

OPHIOGLOSSUM, Tournefort, inst. rei herb. 548, t. 325 (1700), from Bock (1539). (Ophioderma.)

O. vulgatum, C. Bauhin, pinax theatr. bot. 354 (1623) ... ... — S.A. T. V. N.S.W. Q. N.A. B.fl.VII,688 M.fr.V,112;VI,124.
O. pendulum, Linné, sp. pl. edit. sec. 1518 (1763) ... ... — — — — N.S.W. Q. — B.fl.VII,689 M.fr.V,113;VI,124.

BOTRYCHIUM, Swartz in Schrader, Journ. fuer die Bot. II, 110 (1800).

B. Lunaria, Swartz in Schrad. Journ. II, 110 (1800) ... ... — — T. V. N.S.W. — — B.fl.VII,690 M.fr.V,113.
B. ternatum, Swartz, synops. filic. 172 (1806) ... ... — — T. V. N.S.W. Q. — B.fl.VII,690 M.fr.V,113.

HELMINTHOSTACHYS, Kaulfuss, enum. filic. Chamiss. 28 (1824).

H. Zeylanica, Kaulfuss, enum. fil. 28 (1824) ... ... — — — — — Q. — B.fl.VII,690 M.fr.V,113;IX,190.

LYGODIUM, Swartz in Schrader, Journ. fuer die Bot. II, 7 & 106 (1800). (Hydroglossum, Lygodictyon.)

L. scandens, Swartz in Schrad. Journ. II, 106 (1800) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,691 M.fr.V,113;VII,84,X,118.
L. reticulatum, Schkuhr. kryptog. Gewaechse 139, t. 139 (1809) — — — — — Q. — B.fl.VII,692 M.fr.VII,83;VIII,157.
L. Japonicum, Swartz, syn. filic. 154 (1806) ... ... — — — — — Q. N.A. B.fl.VII,692 M.fr.V,113;VII,84,156.

SCHIZAEA, Smith in Mem. de l'Acad. Turin. V, 149, t. 19 (1791). (Lopidium, Actinostachys.)

S. rupestris, R. Brown, prodr. 162 (1810) ... ... — — — — N.S.W. — — B.fl.VII,693 M.fr.V,113.
S. fistulosa, Labillardière, Nov. Holl. pl. spec. II, 103, t. 250 (1806) — S.A. T. V. — — — B.fl.VII,693 M.fr.V,113.
S. dichotoma, Smith in Mém. de l'Ac. Tur. V, 149 (1791) ... — S.A. T. V. N.S.W. Q. N.A. B.fl.VII,694 M.fr.V,113;X,118.
S. Forsteri, Sprengel, Anleit. zur Kenntn. der Gew. III, 157 (1804) — — — — — Q. — — M.fr.VIII,275.

ANGIOPTERIS, G. Hoffmann in Comment. Goett. XII, 29 (1796). [118.

A. evecta, G. Hoffmann in Comment. Goett. XII, 29 (1796) ... — — — — — Q. — B.fl.VII,694 M.fr.V,114;VII,120;X,

MARATTIA, Swartz, nov. gen. et spec. plant. 8 & 128 (1788).

M. fraxinea, Smith, plant. icon. hact. ined. t. 48 (1790)... ... — — — — N.S.W. Q. — B.fl.VII,695 M.fr.V,114;X,118.

TRICHOMANES, Linné, syst. nat. 8 (1835); Linné, hort. Cliffort. 476 (1737).

T. peltatum, Baker in Journ. Linn. Soc. IX, 336, t. 8 (1865) ... — — — — — Q. — B.fl.VII,701 M.fr.X,117.
T. Vitiense, Baker in Journ. Linn. Soc. IX, 338, t. 8 (1865) ... — — — — N.S.W. Q. — B.fl.VII,701 M.fr.VIII,32;IX,132.
T. Motleyi, Bosch, Hymenoph. Javan (1861) ... ... — — — — — Q. — — —
T. cuspidatum, Willdenow, spec. plant. V, 499 (1810) ... ... — — — — — Q. — — M.fr.XI,131&139.
T. parvulum, Poiret, Encycl. méth. VIII, 44 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,701 M.fr.VII,155.
T. digitatum, Swartz, synops. filic. 370 & 422 (1806) ... ... — — — — N.S.W. — — B.fl.VII,702
T. venosum, R. Brown, prodr. 159 (1810) ... ... — — T. V. N.S.W. Q. — B.fl.VII,702 M.fr.V,116;IX,191.
T. Javanicum, Blume, flor. Jav. filic. 224 (1836) ... ... — — — — — Q. — B.fl.VII,702
T. rigidum, Swartz, nov. gen. et spec. pl. 137 (1788) ... ... — — — — N.S.W. Q. — B.fl.VII,702 M.fr.V,115.
T. bipunctatum, Poiret, Encycl. méth. VIII, 44 (1810) ... ... — — — — — Q. — B.fl.VII,703 M.fr.X,117;XI,132.
T. humile, G. Forster, florul. ins. Austr. prodr. 84 (1786) ... — — — — N.S.W. — — —
T. caudatum, Brackenridge, Bot. of Wilk. U.S. Exped. fil. 256, t. 36 (1854) ... ... — — — — N.S.W. Q. — B.fl.VII,703 M.fr.V,116;VII,122.
T. Bauerianum, Endlicher, prodr. fl. Norf. 17 (1833) ... ... — — — — N.S.W. Q. — B.fl.VII,703 M.fr.VII,121;XI,132.
T. parviflorum, Poiret, Encycl. méth. VIII, 44 (1810) ... ... — — — — — Q. — B.fl.VII,704 M.fr.VII,122.

HYMENOPHYLLUM, Smith in Roemer, Archiv I, 56 (1797).

H. marginatum, Hooker & Greville, icon. filic.I, t. 34 (1829) ... — — — — N.S.W. — — B.fl.VII,705
H. nitens, R. Brown, prodr. 159 (1810) ... ... — — T. V. N.S.W. Q. — B.fl.VII,705
H. Javanicum, Sprengel, syst. veg. IV, 132 (1827) ... ... — — T. V. N.S.W. Q. — B.fl.VII,705 M.fr.V,116.
H. Tunbridgense, Smith in Roemer, Archiv. I, 56 (1797) ... — — T. V. N.S.W. Q. — B.fl.VII,706 M.fr.V,116.
H. multifidum, Swartz, synops. filic. 149 & 378 (1806) ... ... — — — — N.S.W. — — B.fl.VII,707

CYATHEA, Smith in Mém. de l'Acad. Turin. V, 416 (1791). (Hemitelia, Amphicosmia.)

C. Lindsayana, Hooker, syn. filic. 25 (1865) ... ... — — — — N.S.W. Q. — B.fl.VII,708 M.fr.VIII,177.
C. arachnoidea, Hooker, syn. filic. 24 (1866) ... ... — — — — — Q. — B.fl.VII,708 M.fr.VI,200.
C. Macarthurii, F. v. M., fragm. VII, 177 (1874) ... ... — — — — N.S.W. — — B.fl.VII,708 M.fr.VIII,176;X,120.
C. medullaris, Swartz, synops. filic. 140 (1806) ... ... — — T. V. N.S.W. — — B.fl.VII,708 M.fr.V,116;VIII,177.
C. Cunninghami, J. Hooker, icon. plant. 985 (1854) ... ... — — T. V. — — — —
C. brevipinnea, Baker in Bentham fl. Austr. VII, 709 (1878) ... — — — — N.S.W. — — B.fl.VII,709 M.fr.XI,139.
C. Moorei, F. v. M.; Hemitelia, Baker in Gardn. Chron. 252 (1872) — — — — N.S.W. — — B.fl.VII,709 M.fr.VIII,157.

CERATOPTERIS, Brogniart in Bull. de la Soc. philom. 186 (1821). (Parkeria.)

C. thalictroides, Brogniart in Bull. Soc. philom. 186 (1821) ... — — — — — Q. N.A. B.fl.VII,695 M.fr.V,122;VIII,158.

GLEICHENIA, Smith in Mém. de l'Acad. Turin. V, 418 (1791). (Mertensia, Platyzoma, Stromatopteris.) [118.

G. platyzoma, F. v. M., fragm. V, 114 (1866) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,696 M.fr.V,114;VIII,157;X,
G. circinata, Swartz, syn. filic. 165 et 394 (1806) ... ... — S.A. T. V. N.S.W. Q. N.A. B.fl.VII,697 M.fr.V,115.
G. dicarpa, R. Brown, prodr. 161 (1810) ... . — — T. V. N.S.W. Q. — B.fl.VII,698 M.fr.V,115.
G. flabellata, R. Brown, prodr. 161 (1810) ... ... — — T. V. N.S.W. Q. — B.fl.VII,698 M.fr.V,114;VIII,157.
G. Hermanni, R. Brown, prodr. 161 (1810) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,698 M.fr.V,114;X,118.

OSMUNDA, Tournefort, inst. rei herb. 547, t. 324 (1700), from l'Obel (1576). (Todea, Leptopteris.)

O. barbara, Thunberg, prodr. plant. Capens. 171 (1800) ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,699 M.fr.V,114.
O. Fraseri, F. v. M.; Todea, Hooker & Greville, icon. filic. t. 101 (1829) — — — — N.S.W. — — B.fl.VII,700 M.fr.V,114;VII,156.
O. Moorei, F. v. M.; Todea, Baker in Trimen, Journ. of Bot. 16 (1873) — — — — N.S.W. — — B.fl.VII,700 M.fr.VIII,157.

ALSOPHILA, R. Brown, prodr. 158 (1810).

A. Rebeccae, F. v. M., fragm. V, 53 (1865) ... ... — — — — — Q. — B.fl.VII,710 M.fr.V,117,215;VIII,180.
A. Loddigesii, Kunze in Schlecht. Linnaea XXIII, 221 (1850) ... — — — — N.S.W. Q. — B.fl.VII,710 [VIII,178.
A. australis, R. Brown, prodr. 158 (1810) ... ... — — T. V. N.S.W. Q. — B.fl.VII,710 M.fr.V,52,116;VII,120;
A. excelsa, R. Brown, prodr. 158 (1810) ... ... — — — — N.S.W. Q. — —
A. Leichhardtiana, F. v. M., fragm. V, 53 (1865) ... ... — — — — N.S.W. Q. — B.fl.VII,711 M.fr.V,117;VI,200;VII,
A. Robertsiana, F. v. M., fragm. V, 54 (1865) ... ... — — — — — Q. — B.fl.VII,712 M.fr.V,117,213. [120.

DICKSONIA, L'Heritier, sert. Angl. 31 (1788). (Cibotium, Patania, Dennstaedtia, Deparia, Balantium.)

D. Billardieri, F. v. M., fragm. VIII, 175 (1874) ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,712 M.fr.V,117;VI,199;VII,
D. Youngiae, C. Moore in Hook. & Bak. syn. filic. 51 (1865) ... — — — — N.S.W. Q. — B.fl.VII,713 M.fr.VI,200. [120.
D. davallioides, R. Brown, prodr. 158 (1810) ... ... — — — V. N.S.W. Q. — B.fl.VII,713 M.fr.VIII,158. [104.
D. nephrodioides; Deparia, Baker in Gardn. Chron. 253 (1872) ... — — — — N.S.W. — — B.fl.VII,714 M.fr.VIII,157;IX,78;X,

DAVALLIA, Smith in Mém. de l'Acad. de Turin. V, 414 (1791). (Microlepia, Humata.)

D. solida, Swartz, synops. filic. 132 & 345 (1806) ... ... — — — — — Q. — B.fl.VII,715
D. elegans, Swartz, synops. filic. 132 & 347 (1806) ... ... — — — — — Q. — B.fl.VII,715 M.fr.V,118;VIII,157.
D. pyxidata, Cavanilles, descripcion de las plantas n. 694 (1802) — — — V. N.S.W. Q. — B.fl.VII,716 M.fr.V,118.
D. pedata, Smith in Mém. de l'Ac. de Turin. V, 414 (1791) ... — — — — — Q. — B.fl.VII,716 M.fr.V,118.
D. dubia, R. Brown, prodr. 157 (1810) ... ... — — T. V. N.S.W. Q. — B.fl.VII,716 M.fr.V,118.
D. flaccida, R. Brown, prodr. 157 (1810) ... ... — — — — — Q. — B.fl.VII,717 M.fr.V,118;VII,156.
D. tripinnata, F. v. M. in Benth. Fl. Austr. VII, 717 (1878) ... — — — — — Q. — B.fl.VII,717

VITTARIA, Smith in Mém. de l'Acad. de Turin. V, 414 (1791).

V. elongata, Swartz, synops. filic. 109 & 302 (1806) ... ... — — — — N.S.W. Q. — B.fl.VII,718 M.fr.V,122;VII,121.

LINDSAYA, Dryander in Mém. de l'Acad. de Turin. V, 413 (1791). (Lindsaea, Isoloma, Synophlebium, Schizoloma.)

L. linearis, Swartz, synops. filic. 118 & 318, t. 3 (1806) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,719 M.fr.V,119.
L. dimorpha, Bailey, Queensland Ferns 19 (1874) ... ... — — — — — Q. — B.fl.VII,719
L. cultrata, Swartz, synops. filic. 119 (1806) ... ... — — — — — Q. — B.fl.VII,719
L. flabellulata, Dryander in Transact. Linn. Soc. III, 41, t. 8 (1797) — — — — — Q. N.A. B.fl.VII,720 M.fr.V,119.
L. lobata, Poiret, Encycl. méth. XI (1813) ... ... — — — — — Q. — B.fl.VII,720
L. trichomanoides, Dryander in Trans. Linn. Soc. III, 43, t. 11 (1797) — — T. — — — — B.fl.VII,720 M.fr.V,118.
L. microphylla, Swartz, synops. filic. 120 (1806) ... ... — — — — N.S.W. Q. — B.fl.VII,721 M.fr.V,119.
L. incisa, Prentice in Trim. Journ. of Bot. 295 (1873) ... ... — — — — — Q. — B.fl.VII,721
L. Fraseri, Hooker, spec. filic. I, 221, t. 70 (1846) ... ... — — — — — Q. — B.fl.VII,721 M.fr.V,118.

L. ensifolia, Swartz, synops. filic. 118 (1806) ... ... -- — — — — Q. N.A. B.fl.VII,721 M.fr.V,118.
L. lanuginosa, Wallich, numerical list n. 154 (1828) ... ... — — — — — Q. — B.fl.VII,722 M.fr.V,118.

ADIANTUM, Tournefort, in 543, t. 317 (1700), from Hippocrates, Theophrastos, Dioscorides and Plinius.

A. lunulatum, N. Burmann, Fl. Ind. 235 (1768) ... ... — — — — — Q. N.A. B.fl.VII,723 M.fr.V,119.
A. Capillus veneris, Linné, spec. plant. 1096 (1753) ... ... — — — — — Q. — B.fl.VII,723
A. Aethiopicum, Linné, syst. nat. edit. decim. n. 15 (1759) ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,724 M.fr.V,119.
A. formosum, R. Brown, prodr. 155 (1810) ... ... — — — V. N.S.W. Q. — B.fl.VII,724 M.fr.V,120;VIII,157.
A. affine, Willdenow, spec. plant. V, 448 (1810) ... ... - — — — N.S.W. Q. — B.fl.VII,724 M.fr.V,119.
A. diaphanum, Blume, flora Javae, filic. 215 (1836) ... ... — — — - N.S.W. Q. — B.fl.VII,725 M.fr.VII,121.
A. hispidulum, Swartz, syn. filic. 124 & 321 (1806) ... ... — — — V. N.S.W. Q. — B.fl.VII,725 M.fr.V,120.

CHEILANTHES, Swartz, syn. filic. 126, t. 3 (1806). (Notholaena, Nothochlaena.)

C. Prenticei, Luerssen in Uhlw. & Behr. Bot. Centralblatt IX, 440 (1882) ... ... — — — — — — N.A. —
C. pumilio, F. v. M.; Notholaena, R. Brown, prodr. 146 (1810) — — — — — Q. — B.fl.VII,773 M.fr.VIII,175.
C. fragillima, F. v. M., fragm. V, 123 (1866) ... ... — — — — — — N.A. B.fl.VII,774 M.fr.V,123.
C. vellea, F. v. M., fragm. V, 123 (1866) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,773 M.fr.V,123;VIII,176.
C. distans, A. Braun, index sem. hort. Berol. (1859) ... .. W.A. S.A. — V. N.S.W. Q. — B.fl.VII,774 M.fr.V,122;VIII,176.
C. tenuifolia, Swartz, syn. filic. 126 (1806) .. .. W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,726 M.fr.V,122.

PTERIS, Linné, syst. nat. 9 (1735); Linné, gen. plant. 322 (1737.) (Pellaea, Cheiloplecton, Platyloma, Litobrochia.)

P. geraniifolia, Raddi, syn. filic. Brasil. 46 (1819) ... ... — — — — N.S.W. Q. — B.fl.VII,728 M.fr.V,124.
P. paradoxa, Baker in Benth. Fl. Austr. VII, 729 (1878) ... — — — — N.S.W. Q. — B.fl.VII,729
P. falcata, R. Brown, prodr. 154 (1810) ... ... — — T. V. N.S.W. Q. — B.fl.VII,729 M.fr.V,123.
P. rotundifolia, G. Forster, florul. ins. Austr. prodr. 79 (1786) .. — — — — — Q. — B.fl.VII,730
P. longifolia, Linné, spec. plant. 1074 (1753) ... ... — — — V. N.S.W. Q. — B.fl.VII,730 M.fr.V,126.
P. ensiformis, N. Burmann, Fl. Ind. 230 (1768) ... ... -- — — — — Q. — B.fl.VII,730 M.fr.V,123.
P. umbrosa, R. Brown, prodr. 154 (1810) ... ... - — — V. N.S.W. Q. — B.fl.VII,730 M.fr.V,126.
P. quadriaurita, Retzius, observ. bot. VI, 38 (1791) ... ... — — — — — Q. — B.fl.VII,731 M.fr.V,125.
P. arguta, Aiton, hort. Kew. III, 458 (1789) ... ... — — T. V. N.S.W. Q. — B.fl.VII,731 M.fr.V,126.
P. aquilina, Linné, spec. plant. 1075 (1753) ... ... W.A S.A. T. V. N.S.W. Q. — B.fl.VII,731 M.fr.V,126.
P. incisa, Thunberg, prodr. pl. Capens. 171 (1800) ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,732 M.fr.V,124;VIII,158.
P. marginata, Bory. voy. dans les isles d'Afr. II, 192 (1804) ... — — — — — Q. — B.fl.VII,733 M.fr.V,125.
P. comans, G. Forster, florul. ins. Austr. prodr. 79 (1786) ... — — T. V. N.S.W. Q. — B.fl.VII,733 M.fr.V,125;VII,121.

LOMARIA, Willdenow in Magaz. der Ges. naturf. Freunde zu Berlin, III, 160 (1809.) (Stegania, Plagiogyria.)

L. Patersoni, Sprengel, syst. veg. IV, 62 (1827) ... ... — — T. V. N.S.W. Q. — B.fl.VII,734 M.fr.V,122;VII,158.
L. vulcanica, Blume, fl. Javae, filic. 202 (1836) .. .. — — T. — — — — B.fl.VII,735 M.fr.V,121.
L. discolor, Willdenow, spec. plant. V, 293 (1810) ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,735 M.fr.V,121.
L. lanceolata, Sprengel, syst. veg. IV, 62 (1827) ... . — S.A. T. V. N.S.W. — — B.fl.VII,735 M.fr.V,121.
L. attenuata, Willdenow, spec. plant V, 290 (1810) ... ... — — — — N.S.W. -- — B.fl.VII,736
L. alpina, Sprengel, syst. veg. IV, 62 (1827) ... ... — — T. V. N.S.W. — — B.fl.VII,736 M.fr.V,121.
L. fluviatilis, Sprengel, syst. veg. IV, 65 (1827) ... ... — — T. V. N.S.W. -- — B.fl.VII,736 M.fr.V,121.
L. Fullagari, F. v. M., fragm. VIII, 157 (1874) ... ... — — — — N.S.W. — B.fl.VII,737 M.fr.VIII,157.
L. Capensis, Willdenow, spec. plant. V, 291 (1810) ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,737 M.fr.V,121.
L. euphlebia, Kunze in Mohl & Schlecht. Bot. Zeit. VI, 521 (1849) — — — — — Q. - B.fl.VII,738 M.fr.VI,124.

BLECHNUM, Linné, spec. plant. 1077 (1753.)

B. cartilagineum, Swartz, syn. filic. 114 & 312 (1806) ... ... — — -- V. N.S.W. Q. — B.fl.VII,738 M.fr.V,120.
B. laevigatum, Cavanilles, descripcion de las plantas 650 (1802) — — — — N.S.W. — — B.fl.VII,739 M,fr.V,120.
B. serrulatum, C. Richard in Annal. du Mus. d'Hist. nat. Par. I, 114 (1802) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,739 M.fr.V,120.
B. orientale, Linné, syst. nat. ed. decim. (1759) ... ... -- — — — — Q. N.A. B.fl.VII,740

MONOGRAMMA, Commerçon in Schkuhr's Kryptog. Gew. 82, t. 87 (1809). (Monogramme, Vaginularia, Pleurogramme, Diclidopteris.)

M. Junghuhnii, Hooker, spec. filic. V, 123, t. 289 (1864) ... — — — — — Q. — B.fl.VII,740 M.fr.VII,119.

WOODWARDIA, Smith in mém. de l'Acad. de Turin. V, 411 (1791). (Doodia.)

W. aspera, Mettenius, fil. hort. bot. Lips. 65 (1856) ... — — — V. N.S.W. Q. — B.fl.VII,740 M.fr.V,130.
W. caudata, Cavanilles, descripcion de las pl. n. 653 (1802) ... — — T. V. N.S.W. Q. N.A. B.fl.VII,742 M.fr.V,129;VIII,158.

ASPLENIUM, Linné, gen. plant. 322 (1737) from C. Bauhin (1651.) (Scolopendrium, Allantodea, Diplazium, Callipteris, Anisogonium, Thamnopteris, Neottopteris, Darea, Caenopteris, Athyrium, Diplora, Asplenum.)

A. Nidus, Linné, spec. plant. 1079 (1753) ... ... — — — V. N.S.W. Q. — B.fl.VII,744 M.fr.V,130.
A. simplicifrons, F. v. M., fragm. V, 74 (1865) ... ... — — — — — Q. — B.fl.VII,744 M.fr.V,74,130;VIII,158.
A. attenuatum, R. Brown, prodr. 150 (1810) ... ... - — — — N.S.W. Q. — B.fl.VII,745 M.fr.V,130.
A. Trichomanes, Linné, spec. plant. 1080 (1753) ... ... — — T. V. N.S.W. — — B.fl.VII,745 M.fr.V,131
A. flabellifolium, Cavanilles, descripcion de las pl. n. 636 (1802) W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,745 M.fr.V,131.
A. paleaceum, R. Brown, prodr. 150 (1810) ... ... — — — — — Q. — B.fl.VII,746 M.fr.V,131;VIII,158.
A. falcatum, Lamarck, Encycl. méth. I, 303 (1783) ... ... — — — — N.S.W. Q. — B.fl.VII,746 M.fr.V,131;VII,156.
A. Hookerianum, Colenso in Journ. of Nat. Sc. Tasm. II, 169 (1842) — — — V. N.S.W. — — B.fl.VII,747
A. furcatum, Thunberg, prodr. pl. Cap. 172 (1800) ... ... W.A. S.A. — V. N.S.W. — - B.fl.VII,748 M.fr.V,131.
A. laserpitiifolium, Lamarck, Encycl. méth. I, 310 (1783) ... — — — — — Q. — B.fl.VII,749 M.fr.V,131.
A. marinum, Linné, spec. plant. 1081 (1753) ... ... — — T. V. N.S.W. — — B.fl.VII,747 M.fr.V,132.
A. bulbiferum, G. Forster, floral. ins. Austr. prodr. 433 (1786)... — S.A. T. V. N.S.W. Q. — B.fl.VII,749 M.fr.V,132.
A. pteridoides, Baker, syn. filic. sec. edit. 488 (1874) ... ... — — — — N.S.W. — — B.fl.VII,749
A. umbrosum, J. Smith in Hook. Lond. Journ. IV, 174 (1842)... — — T. V. N.S.W. — — B.fl.VII,749 M.fr.V,132.
A. pallidum, Blume, fl. Javae, filic. 776 (1836) ... ... — — — — — Q. — — M.fr.XI,131.
A. sylvaticum, Presl, reliq. Haenk. I, 42 (1830) ... ... — — — — — Q. — B.fl.VII,750 M.fr.V,133.
A. maximum, D. Don, prodr. fl. Nepal. 8 (1825) ... ... -- — — — N.S.W. Q. — B.fl.VII,751
A. polypodioides, Mettenius, fil. hort. Lips. 78 (1856) ... ... — — — — — Q. — B.fl.VII,751 M.fr.V,132.
A. melanochlamys, Hooker, spec. filic. III, 259 (1860) ... ... — — — — N.S.W. — — B.fl.VII,751
A. decussatum, Swartz, spec. fil. 76 & 260 (1806) ... ... — — — — — Q. — B.fl.VII,751

CYSTOPTERIS, Bernhardi in Schrader's neuem Journ. II, 26 (1806).

C. fragilis, Bernhardi in Schrad. Journ. I, 26 (1806)	—	—	T.	—	—	—	—	B.fl.VII,752	M.fr.V,118.			

ASPIDIUM, Swartz in Schrader's Journ. II, 429 (1000). (Nephrodium, Nephrolepis, Polystichum, Lastraea, Sagenia, Oleandra.)

A. cordifolium, Swartz, syn. filic. 45 (1806)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,754	M.fr.V,136.			
A. exaltatum, Swartz, syn. filic. 45 (1806)...	—	—	—	—	—	Q.	N.A.	B.fl.VII,754	M.fr.V,136			
A. ramosum, Palisot, Fl. d'Oware et de Benin II, 53, t. 91 (1807)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,754	M.fr.V,135.			
A. unitum, Swartz, syn. filic. 47 (1806)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,755	M.fr.V,137;	VIII,158.		
A. pteroides, Swartz, syn. filic. 49 (1806)	—	—	—	—	—	Q.	—	B.fl.VII,755				
A. molle, Swartz, syn. filic. 49 (1806)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,756	M.fr.V,135.			
A. truncatum, Gaudichaud in Voy. Freyc. Bot. 332, t. 10 (1826)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,756	M.fr.V,135.			
A. confluens, Mettenius in Schlecht. Linnaea XXXVI, 125 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.VII,757	M.fr.V,133;	VIII,158.		
A. aculeatum, Swartz in Schrad. Journ. II, 37 (1800)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,757	M.fr.V,134.			
A. aristatum, Swartz, syn. filic. 53 & 253 (1806)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,757	M.fr.V,134.			
A. Capense, Willdenow, spec. plant. V, 267 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,758	M.fr.V,134.			
A. decompositum, Sprengel, syst. IV, 109 (1827)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,758	M.fr.V,136.			
A. tenerum, Sprengel, syst. IV, 109 (1827)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,759				
A. uliginosum, Kunze in Schlecht. Linnaea XX, 6 (1847) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,759	M.fr.V,133.			
A. hispidum, Swartz, syn. filic. 56 (1806)	—	—	T.	V.	—	—	—	B.fl.VII,760	M.fr.V,133.			

POLYPODIUM, Tournefort, inst. 540, t. 316 (1700), from Theophrastos, Dioscorides & Plinius. (Niphobolus, Goniophlebium, Goniopteris, Phlebodium, Phegopteris, Phymatodes, Pleopeltis, Drynaria, Dictyopteris, Arthropteris, Xiphopteris, Meniscium partly; from Dodoens, l'Ecluse, Bauhin, Morison, Ray, Plumier and particularly Petiver as Polypodium.)

P. australe, Mettenius, filic. hort. bot. Lips. 36 (1856)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,762	M.fr.V,127.			
P. Hookeri, Brackenridge, Bot. Wilk. Exped. Filic. 4 (1854) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,763				
P. contiguum, Brackenridge, Bot. Wilk. Exp. Fil. 6 t. (1854) ...	—	—	—	—	—	Q.	—	B.fl.VII,763	M.fr.V,127.			
P. grammitidis, R. Brown, prodr. 147 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,764	M.fr.V,127.			
P. tenellum, G. Forster, florul. ins. Austr. prodr. 440 (1786) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,764	M.fr.V,126;	VII,121.		
P. proliferum, Roxburgh in Wall. num. list 11 (1828)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,765	M.fr.V,128.			
P. urophyllum, Wallich, numerical list 11 (1828)	—	—	—	—	—	Q.	—	B.fl.VII,765	M.fr.IV,165;	V,137;	VIII,	
P. Hillii, Baker in Hook., syn. filic. sec. edit. 505 (1874) ...	—	—	—	—	—	Q.	—	B.fl.VII,766				[121.
P. poecilophlebium, Hooker, spec. filic. V, 14 (1863)	—	—	—	—	—	Q.	—	B.fl.VII,766	M.fr.V,127.			
P. serpens, G. Forster, florul. insul. austr. prodr. 81 (1786) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,767	M.fr.V,129.			
P. confluens, R. Brown, prodr. 146 (1810)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,767				
P. acrostichoides, G. Forster, florul. insul. austr. prodr. 81 (1786)	—	—	—	—	—	Q.	—	B.fl.VII,767	M.fr.V,129.			
P. attenuatum, R. Brown, prodr. 146 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,768	M.fr.V,128;	V,121.		
P. simplicissimum, F. v. M. in Hook. & Bak. syn. fil. sec. edit. 513 (1874)	—	—	—	—	—	Q.	—	B.fl.VII,768	M.fr.VII,120,156.			
P. nigrescens, Blume, Fl. Javae, filic. 161, t. 70 (1836)	—	—	—	—	—	Q.	—	B.fl.VII,769				
P. phymatodes, Linné, mant. pl. alt. 306 (1771)	—	—	—	—	—	Q.	N.A.	B.fl.VII,769				
P. pustulatum, G. Forster, florul. ins. Austr. pr. 81 (1786) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,769	M.fr.V,127.			
P. scandens, G. Forster, florul. ins. Austr. pr. 81 (1786) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,770	M.fr.V,128.			
P. verrucosum, Wallich, numerical list 11 (1828)...	—	—	—	—	—	Q.	—	B.fl.VII,770				
P. subauriculatum, Blume, Fl. Javae, filic. 177, t. 83 (1836) ...	—	—	—	—	—	Q.	—	B.fl.VII,770	M.fr.V,128.			
P. rigidulum, Swartz, synops. filic. 38 & 320 (1806) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,771				
P. quercifolium, Linné, spec. plant. 1087 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VII,772	M.fr.V,127.			
P. irioides, Poiret, Encycl. méth. V, 513 (1804)	—	—	—	—	—	Q.	—	B.fl.VII,771	M.fr.V,127.			
P. punctatum, Thunberg, fl. Japonic. 337 (1784)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,764	M.fr.V,129.			

HYPOLEPIS, Bernhardi in Schrader, neu. Journ. I, 34 (1806).

H. tenuifolia, Bernhardi in Schrad. Journ. II, 24 (1800) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,726	M.fr.V,129.			

GRAMMITIS, Swartz in Schrader, Journ. II, 3 & 17 (1800). (Gymnogramme, Selliguea).

G. Reynoldsii, F. v. M. in Benth. Fl. Austr. VII, 775 (1878) ...	—	S.A.	—	—	—	—	—	B.fl.VII,775	M.fr.VIII,175.			
G. Muelleri, Hooker in F. v. M. fragm. V, 138 (1866)	—	—	—	—	—	Q.	—	B.fl.VII,775	M.fr.V,138;	VI,124;	VII,	
G. rutifolia, R. Brown, prodr. 146 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,775	M.fr.V,137.			[121.
G. leptophylla, Swartz, syn. fil. 23 et 218, t. 1 (1806)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,776	M.fr.V,137.			
G. pinnata, F. v. M., fragm. VI, 124 (1868)	—	—	—	—	—	Q.	—	B.fl.VII,776	M.fr.VI,124.			
G. ampla, F. v. M., fragm. V, 188 (1866)	—	—	—	—	—	Q.	—	B.fl.VII,777	M.fr.V,188.			

ANTROPHYUM, Kaulfuss, enum. filic. 197 (1824).

A. reticulatum, Kaulfuss, enum. filic. 197 (1824)...	—	—	—	—	—	Q.	—	B.fl.VII,777	M.fr.V,138.			

ACROSTICHUM, Linné, gen. plant. 322 (1737), indicative. (Elaphoglossum, Stenochlaena, Lomariopsis, Hymenolepis, Gymnopteris, Chrysodium.)

A. conforme, Swartz, syn. filic. 10 & 192, t. 1 (1836)	—	—	—	—	—	Q.	—	B.fl.VII,778	M.fr.V,138.			
A. scandens, J. Smith in Hook. spec. fil. V, 249 (1864) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,778	M.fr.VI,124.			
A. sorbifolium, Linné, spec. plant. 1069 (1753)	—	—	—	—	—	Q.	—	B.fl.VII,779	M.fr.VII,119.			
A. neglectum, Bailey in proc. Linn. Soc. of N.S.W., V, 32 (1880)	—	—	—	—	—	Q.	—	—	M.fr.XI,131.			
A. repandum, Blume, Fl. Javae 39, t. 14 & 15 (1836)	—	—	—	—	—	Q.	—	B.fl.VII,779	M.fr.V,138.			
A. aureum, Linné, spec. plant. 1069 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,779	M.fr.V,138;	VII,120.		
A. spicatum, Linné, fil. suppl. pl. 444 (1781)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,780	M.fr.V,138.			
A. pteroides, R. Brown, prodr. 145 (1810)...	—	—	—	—	—	Q.	N.A.	B.fl.VII,780	M.fr.X,76.			

PLATYCERIUM, Desvaux in Mém, de la Soc. Linn. de Paris VI, 213 (1827).

P. alcicorne, Desvaux in Mém. Soc. Linn. Par. VI, 213 (1827)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,780	M.fr.V,139.			
P. grande, J. Smith in Hooker, Lond. Journ. III, 402 (1841) .	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,781	M.fr.V,139.			

ADDITIONAL SPECIES.

Pachynema sphenandrum, F. v. M. & Tate in Transact. R. S. of S. Austr. V, 79 (1882)	—	—	—	—	—	—	N.A.	—		
Polyalthia Holtzeana, F. v. M. in Wing's S. Sc. Record II, 230 (1882)	—	—	—	—	—	—	N.A.	—		
Philotheca Hassellii, F. v. M., fragm. XII, inedit.	W.A.	—	—	—	—	—	—	—		
Tribulus astrocarpus, F. v. M., fragm. XII, 4 (1882)	W.A.	—	—	—	—	—	—	—		M.fr.XII,
Euphorbia obliqua, Bauer in Endl. prodr. fl. Norfolk. 85 (1833)	—	—	—	—	N.S.W.	—	—	—		
Phyllanthus Tatei, F. v. M. in Wing's S. Sc. Record II. 55 (1882)	—	S.A.	—	—	N.S.W.	—	—	—		
Ficus Pinkiana, F. v. M. in Wing's S. Sc. Record II, 273 (1882)	—	—	—	—	—	Q.	—	—		
Peperomia Baueriana, Miquel, syst. piperac. 120 (1843) ...	—	—	—	—	N.S.W.	—	—	—		
Nepenthes Bernaysii, Bailey in proceed. L. S. of N.S.W. 185 (1880)	—	—	—	—	—	Q.	—	—		
Ptilotus Polakii, F. v. M. in Wing's S. Sc. Record II, 274 (1882)	W.A.	—	—	—	—	—	—	—		
Atriplex Bunburyanum, F. v. M. in Wing's S. Sc. Record II, 274 (1882)	W.A.	—	—	—	—	—	—	—		
Kochia prosthecochaeta, F. v. M., fragm. XII, 14 (1882) ...	W.A.	—	—	—	—	—	—			M.fr.XII,14
Kochia melanocoma, F. v. M., fragm. XII, 14 (1882)	W.A.	—	—	—	—	—	—	—		M.fr.XII,14
Bassia astracantha, F. v. M., fragm. XII, 12 (1882)	W.A.	—	—	—	—	—	—	—		M.fr.XII,12
Bassia tridens, F. v. M., fragm. XII, 12 (1882)	W.A.	—	—	—	—	—	—			M.fr.XII,12
Bassia Forrestiana, F. v. M., fragm. XII, 12 (1882)	W.A.	—	—	—	—	—	—	—		M.fr.XII,12
Bossiaea Scortechinii, F. v. M. (inedit).	—	—	—	—	N.S.W.	Q.	—	—		
Bossiaea Webbii, F. v. M. in Melb. Chemist, Dec. (1882) ...	W.A.	—	—	—	—	—	—	—		
Isotropis Forrestii, F. v. M. in Wing's S. Sc. Record II, 252 (1882)	W.A.	—	—	—	—	—	—	—		
Daviesia arborea, F. v. M. & Scortechini in proc. L. S. of N.S.W. VII, 221 (1882)	—	—	—	—	N.S.W.	Q.	—	—		
Crotalaria striata, De Candolle, prodr. II, 131 (1825)	—	—	—	—	—	Q.	—	—		
Indigofera tinctoria, Linné, spec. plant. 751 (1753)	—	—	—	—	—	Q.	—	—		
Swainsona Oliverii, F. v. M. in Wing's S. Sc. Rec. II, 152 (1882)	W.A.	S.A.	—	—	—	—	—	—		
Callistemon pithyoides, Miquel in Nederl. Kruidk. Arch. IV, 142 (1859)	—	—	—	V.	N.S.W.	Q.	—	—		
Melaleuca cylindracea, R. Brown in Benth. Fl. Austral. III, 146 (1866)	—	S.A.	—	—	—	—	—	—		
Eucalyptus Foelscheana, F. v. M. in Melb. Chemist, Nov. (1882)	—	—	—	—	—	—	N.A.	—		
Cryptandra Wayii; Trymalium, F. v. M. & Tate in Transact. R. S. of S. Austr. V, 80 (1882)	—	S.A.	—	—	—	—	—	—		
Adenanthos Forrestii, F. v. M. in Wing's S. Sc. Record II, 230 (1882)	W.A.	—	—	—	—	—	—	—		
Grevillea deflexa, F. v. M. inedit.	W.A.	—	—	—	—	—	—	—		
Epaltes Tatei, F. v. M. inedit.	—	S.A.	—	—	—	—	—	—		
PODOSPERMA, Labillardière, Nov. Holl. plant. specim. II, 35, t. 177 (1806). (Podotheca.)										
Podosperma Polakii, F. v. M., fragm. XII, 21 (1882)	W.A.	—	—	—	—	—	—	—		M.fr.XII,21
Helipterum sterilescens, F. v. M. in Wing's S. Sc. Record II, 274 (1882)	W.A.	—	—	—	—	—	—	—		
Helipterum Forrestii, F. v. M. in Wing's S. Sc. Record II, 273 (1882)	W.A.	—	—	—	—	—	—	—		
Velleya macroplectra, F. v. M., fragm. XII, 22 (1882)	W.A.	—	—	—	—	—	—	—		M.fr.XII,22
Thunbergia fragrans, J. Koenig in Roxb. pl. of Coromand. I, 44, t. 67 (1795)	—	—	—	—	—	—	N.A.	—		
ISANDRA, F. v. M., fragm. phyt. Austr. XII (inedit.) Near Anthocercis.										
Isandra Bancroftii, F. v. M., fr. XII (inedit.)	W.A.	—	—	—	—	—	—	—		
DICLADANTHERA, F. v. M., fragm. phytogr. Austral. XII, 23 (1882).										
Dicladanthera Forrestii, F. v. M., fragm. XII, 23 (1882) ...	W.A.	—	—	—	—	—	—	—		M.fr.XII,23
Prostanthera Campbelli, F. v. M. in Wing's S. Sc. Record II, 252 (1882)	W.A.	—	—	—	—	—	—	—		
Chloanthes lepidota, F. v. M. inedit.	W.A.	—	—	—	—	—	—	—		
Myoporum obscurum, Endlicher, prodr. fl. Norfolk. 54 (1833)...	—	—	—	—	N.S.W.	—	—	—		
Eremophila Pantoni, F. v. M. in Wing's S. Sc. Record II, 251 (1882)	W.A.	—	—	—	—	—	—	—		
ERIA, Lindley in Edwards, Bot. Regist. t. 904 (1825).										
Eria Fitzalani, F. v. M. in Wing's S. Sc. Record II, 252 (1882)	—	—	—	—	—	Q.	—	—		
LUISIA, Gaudichaud in Freycin. voy. Bot. 426, t. 37 (1826).										
Luisia teretifolia, Gaudichaud in Freyc. voy. 426, t. 37 (1826)	—	—	—	—	—	Q.	—	—		
Prasophyllum Tepperi, F. v. M. in Tepp. pl. about Ardross., Oct. (1880)	—	S.A.	—	—	—	—	—	—		
Smilax purpurata, G. Forster, florul. ins. Austr. prodr. 70 (1786)	—	—	—	—	N.S.W.	—	—	—		
Pandanus Moorei, F. v. M. inedit.	—	—	—	—	N.S.W.	—	—	—		
Carex haematostoma, Nees in Wight, contrib. to the Bot. of Ind. 125 (1834)	—	—	T.	—	—	—	—	—		

ADDITIONAL ANNOTATIONS.

Myosurus from l'Obel (1576).
Myosurus minimus ... ... Q.
Ranunculus from Plinius.
Ranunculus aquatilis ... ... N.S.W.
Hibbertia hirsuta... ... ... S.A.
Doryphora Sassafras ... ... Q.
Endiandra Sieberi ... ... Q.
Tristichocalyx pubescens ... ... N.A.
Papaver from Plinius.
Cleome viscosa ... ... W.A.
Capparis spinosa ... ... S.A.
Capparis lasiantha ... ... W.A.
Barbarea from l'Obel (1576).
Barbarea vulgaris ... ... S.A.
Capsella pilosula ... ... W.A.
Senebiera=Carara, Medicus, Pflanzen-Gattung. 34 (1792), from Cesalpini (1583). (Coronopus partly).
Lepidium strongylophyllum ... ... W.A.
Lepidium leptopetalum ... ... W.A.
Lepidium rotundum ... ... S.A.
Cakile from l'Obel (1576).
Hybanthus enneaspermus ... ... W.A.
Hymenanthera Banksii ... ... S.A.
Pittosporum revolutum ... ... M.fr.XII,4.
Pittosporum ferrugineum ... ... M.fr.XII,4.
Pittosporum melanospermum ... ... Q.
Hymenosporum flavum ... ... M.fr.XII,4.
Marianthus microphyllus ... ... M.fr.XII,2.
Citriobatus pauciflorus ... ... N.S.W.
Billardiera variifolia ... ... M.fr.XII,2.
Cheiranthera linearis ... ... M.fr.XII,4.
Cheiranthera filifolia ... ... M.fr.XII,4.
Bergia perennis ... ... S.A.
Polygala from Dioscorides & Plinius.
Polygala persicarifolia ... ... Q.
Polygala Chinensis ... ... S.A.
Comesperma sylvestre ... ... S.A.,N.S.W.
Comesperma viscidulum... ... ... S.A.
Tribulus Hystrix ... ... W.A.
Hedraianthera porphyropetala ... ... N.S.W.
Tribulus macrocarpus ... ... W.A.
Tribulus platypterus ... . W.A.
Sida virgata ... ... W.A.
Sida petrophila ... ... W.A.
Sida rhombifolia ... . S.A.
Sida physocalyx ... ... W.A.
Sida lepida ... ... W.A.,S.A.
Abutilon tubulosum ... ... N.A.
Abutilon cryptopetalum ... ... M.fr.IX,131.
Abutilon geranioides ... ... W.A.
Abutilon Avicennae, Gerard, gen. hist. of pl. 790 (1597).
Urena lobata ... ... N.A.
Hibiscus panduriformis ... ... W.A.,Q.
Hibiscus Normani ... ... M.fr.V,44.
Hibiscus Sturtii ... ... W.A.
Gossypium australe ... . W.A. M.fr.V,44.
Gossypium flaviflorum ... ... M.fr.V,44.
Waltheria Indica ... ... W.A.
Commerçonia Fraseri ... ... Q.
Euphorbia erythrantha ... ... W.A.
Euphorbia pilulifera ... ... N.A.
Monotaxis luteiflora ... ... S.A.
Poranthera microphylla ... ... M.fr.XII,11.
Pseudanthus micranthus... ... ... M.fr.XII,11.
Beyeria viscosa ... ... M.fr.XII,10.
Ricinocarpus pinifolius ... ... S.A.
Bertya rosmarinifolia ... ... Q. M.fr.XII,10.
Bertya rotundifolia ... ... M.fr.XII,11.
Micrantheum hexandrum ... ... M.fr.XII,11.
Dissiliaria tricornis ... ... M.fr.XII,11.
Phyllanthus rhytidospermus ... ... S.A. M.fr.XII,11.
Phyllanthus rigens ... ... M.fr.XII,11.
Phyllanthus Gastroemii ... ... M.fr.XII,11.
Phyllanthus australis ... ... M.fr.XII,11.
Securinega Leucopyrus ... ... S.A., N.A.
Adriana tomentosa ... ... M.fr.XII,11.
Adriana quadripartita ... ... M.fr.XII,11.
Tragia Novae Hollandiae ... ... M.fr.XII,11.
Fontainea Pancheri ... ... M.fr.XII,11.
Omalanthus populifolius... ... . M.fr.XII,11.
Ficus platypoda ... ... W.A.
Ficus scabra ... ... N.A.
Ficus stenocarpa=Ficus subglabra, F. v. M., fragm. IX, 152 (1875) ... ... N.S.W.,Q.
Pouzolzia quinquenervis ... ... N.A.
Casuarina suberosa ... ... S.A.
Casuarina inophloia ... ... N.S.W.
Peperomia Urvilleana ... ... N.S.W.
Leea aculeata=L. Brunoniana, Clarke in Britten's Journ. of Bot. X, 105 (1881).
Castanospora Alphandi ... ... N.S.W.
Atalaya variifolia ... ... Q.
Dodonaea lanceolata ... ... S.A.
Dodonaea truncatiales=D. calycina, Cunningham in A. Gray, Bot. of Wilk. Expl. Exped. I, 262 (1854).
Dodonaea platyptera ... ... W.A.
Dodonaea microzyga ... ... S.A.
Bueseraceae=Burseraceae.
Celastrus bilocularis ... ... N.S.W.
Stackhousia muricata ... ... S.A.
Plumbago Zeilanica ... ... S.A.
Aegialitis annulata ... ... N.A.
Portulaca from W. Turner (1538).
Sagina apetala ... ... N.S.W.
Saponaria from Bock (1539).
Ptilotus incanus ... ... S.A.
Ptilotus psilotrichoides ... ... N.A.
Ptilotus helipteroides ... ... W.A.
Ptilotus rotundifolius ... ... W.A.
Ptilotus latifolius... ... ... W.A.
Rhagodia Billardieri ... ... N.A. M.fr.XII,16.
Rhagodia parabolica ... ... W.A.,V. M.fr.XII,15.
Rhagodia crassifolia ... ... Q. M.fr.XII,15.
Rhagodia Preissii ... ... S.A. M.fr.XII,15.
Rhagodia Gaudichaudii ... ... M.fr.XII,15.
Rhagodia spinescens ... ... M.fr.XII,15.
Rhagodia nutans ... ... W.A.,N.A. M.fr.XII,15.
Rhagodia linifolia ... ... M.fr.XII,15.
Chenopodium nitrariaceum ... ... M.fr.XII,16.
Chenopodium microphyllum ... ... M.fr.XII,16.
Chenopodium rhadinostachyum (1882).
Chenopodium cristatum ... ... M.fr.XII,17.
Chenopodium atriplicinum ... ... M.fr.XII,17.
Chenopodium carinatum ... ... N.A. M.fr.XII,17.
Dysphania plantaginella... ... ... W.A.
Dysphania litoralis ... ... M.fr.XII,17.
Dysphania myriocephala ... ... M.fr.XII,17.
Atriplex from W. Turner (1538).
Atriplex prostratum includes A. microcarpum and A. Pumilio ... ... M.fr.XII,17.
Atriplex fissivalve ... ... M.fr.XII,16.
Atriplex paludosum ... ... M.fr.XII,16.
Atriplex halimoides ... ... W.A.,N.A. M.fr.XII,17.
Atriplex vesicarium ... ... W.A. M.fr.XII,17.
Atriplex holocarpum ... ... M.fr.XII,17.
Atriplex nummularium ... ... M.fr.XII,17.
Atriplex leptocarpum ... ... M.fr.XII,17.
Atriplex crystallinum ... ... M.fr.XII,17.
Kochia sedifolia ... ... W.A. M.fr.XII,15.
Kochia lobiflora ... ... M.fr.XII,15.
Kochia lanosa ... ... M.fr.XII,15.
Kochia triptera ... ... M.fr.XII,15.
Kochia oppositifolia ... ... M.fr.XII,15.
Kochia brevifolia ... ... M.fr.XII,15.

Kochia humillima ... ... M.fr.XII,15.
Kochia planifolia ... ... M.fr.XII,15.
Kochia microphylla ... ... M.fr.XII,15.
Kochia brachyptera ... . M.fr.XII,15.
Kochia ciliata ... ... Q. M.fr.XII,15.
Kochia stelligera ... ... Q. M.fr.XII,15.
Babbagia dipterocarpa ... ... M.fr.XII,15.
Bassia tricornis ... ... S.A.
Bassia carnosa .. .. M.fr.XII,13.
Bassia sclerolaenoides ... ... M.fr.XII,13.
Bassia bicornis ... ... M.fr.XII,13.
Bassia paradoxa ... ... M.fr.XII,13.
Bassia salsuginosa ... . M.fr.XII,13.
Bassia lanicuspis ... ... W.A.,N.A. M.fr.XII,13.
Bassia biflora ... ... Q. M.fr.XII,13.
Bassia echinopsila ... ... W.A. M.fr.XII,13.
Bassia enchylaenoides ... ... Q. M.fr.XII,13.
Enchylaena tomentosa ... ... M.fr.XII,15.
Salicornia from Gerard (1597).
Salicornia robusta ... ... W.A.,N.A. M.fr.XII,18.
Salicornia arbuscula ... ... M.fr.XII,18.
Salicornia cinerea ... ... M.fr.XII,18.
Salicornia australis ... ... M.fr.XII,18.
Salsola Kali ... ... M.fr.XII,18.
Tetragonia expansa ... ... W.A.
Zaleya decandra ... ... W.A.
Trianthema turgidifolia ... . W.A.
Trianthema crystallina ... ... W.A.
Trianthema pilosa ... ... W.A.
Macarthuria australis ... ... M.fr.II,11.
Mollugo Ginus = M. Glinus.
Mollugo Cerviana ... ... W.A.,N.A.
Polygonum barbatum ... ... N.A.
Didymotheca pleiococca ... ... N.A.
Gastrolobium grandiflorum ... ... S.A.
Mirbelia speciosa ... ... Q.
Mirbelia daviesioides = M. aphylla, F. v. M., fragm. IV, 11 (1863).
Gompholobium latifolium ... ... V.
Aotus villosa .. .. S.A.
Crotalaria medicaginea ... ... S.A.
Psoralea cephalantha ... ... N.A.
Psoralea balsamica ... ... S.A.
Psoralea Schultzii ... ... N.A.
Indigofera monophylla ... ... W.A.
Tephrosia flammea ... ... N.A.
Tephrosia juncea ... ... N.A.
Sesbania aculeata ... ... W.A.
Swainsona coronillifolia ... ... S.A.
Swainsona colutoides ... ... W.A.
Aeschynomene Indica ... ... S.A.,N.S.W. M.fr.XII,18.
Aeschynomene falcata ... ... M.fr.XII,18.
Aeschynomene Americana ... ... M.fr.XII,18.
Uraria lagopoides ... ... N.A.
Lespedeza lanata ... ... N.A.
Erythrina vespertilio ... ... W.A.
Canavalia obtusifolia ... ... W.A.
Vigna lanceolata ... ... W.A.
Rhynchosia minima ... ... W.A.,S.A.
Caesalpinia Bonducella (1810).
Labichea Buettneriana ... ... M.fr.XII,18.
Labichea punctata ... ... M.fr.XII,19.
Labichea rupestris ... ... M.fr.XII,19.
Adenanthera abrosperma ... ... N.A.
Acacia iteaphylla ... ... S.A.
Acacia dissoneura ... ... N.A.
Acacia Doratoxylon ... ... N.A.
Acacia Farnesiana ... ... W.A.
Acacia triptera, A. rhigiophylla, A. Oxycedrus, A. verticillata, A. Riceana typographically transposed from Juliferae to Plurinerves.
Albizzia Hendersoni ... ... M.fr.V,191.
Alchemilla from Bock (1539).
Ammannia crinipes = Koehnea crinipes, F. v. M. (inedit.) = Nesaca, sect. Crinipedium, Koehne in Engler, bot. Jahrb. III, 337 (1882).
Lagerstroemia Flos Reginae ... ... M.fr.XII,20.
Ceratophyllum demersum ... ... N.A.
Bruguiera caryophylloides ... ... Q.
Thryptomene cricaea ... . V.
Baeckea densifolia ... ... Q.
Kunzea peduncularis ... ... Q.
Melaleuca armillaris ... ... Q.
Eucalyptus amygdalina ... ... S.A
Eucalyptus macrorrhyncha ... ... S.A.
Eucalyptus capitellata ... ... Q.
Eucalyptus clavigera ... ... Q.
Eucalyptus tessellaris ... ... S.A.
Eucalyptus populifolia ... ... N.A.
Eucalyptus microtheca ... ... N.A.
Pomaderris prunifolia ... ... Q.
Pomaderris myrtilloides ... ... S.A.
Astrotricha Hamptoni ... ... N.A.
Didiscus hemicarpus = D. setulosus, F. v. M. in Proceed. R. S. Tasm III, 238 (1857).
Trachymene cirrosa (not in S.A. and V.) ... W.A.
Trachymene heterophylla (not in W.A.) ... V.
Actinotus Gibbonsii ... ... Q.
Oenanthe from l'Obel (1576).
Choretrum lateriflorum ... ... Q.
Viscum from W. Turner (1538).
Viscum angulatum ... ... N.A.
Viscum articulatum ... ... S.A.
Isopogon ceratophyllus ... ... W.A.
Isopogon asper (not in S.A., T., V.) ... ... W.A.
Persoonia lanceolata ... ... Q.
Grevillea agrifolia ... ... S.A.
Grevillea ilicifolia ... ... Q.
Grevillea juncifolia ... ... N.A.
Grevillea ramosissima ... ... V.
Grevillea Chrysodendron ... ... W.A.
Grevillea stenobotrya ... . W.A.
Grevillea striata ... ... W.A.
Hakea Cunninghamii ... ... W.A.
Hakea saligna ... ... V.
Hakea microcarpa ... ... Q.
Hakea multilineata ... ... N.A.
Banksia serrata ... ... Q.
Pimelea sanguinea ... ... N.A.
Pimelea trichostachya ... ... S.A.
Pimelea elachantha ... ... S.A.
Pimelea ammocharis ... ... S.A.
Ixora Dallachyana ... ... N.A.
Canthium coprosmoides ... ... N.A.
Canthium latifolium ... ... W.A.
Oldenlandia (Anotis).
Oldenlandia scleranthoides, Bentham & J. Hooker, gen. pl. II, 88 (1873).
Oldenlandia galioides ... ... W.A.
Opercularia varia ... ... Q.
Asperula from Dalechamps (1554).
Galium umbrosum ... ... W.A.
Sambucus from Plinius.
Trichosanthes cucumerina ... ... Q.
Cucumis trigonus .. .. W.A.
Eupatorium from Gerard (1597).
Lagenophora Emphysopus ... ... S.A.
Brachycome Stuartii ... ... Q.
Brachycome graminea, F. v. M., first gen. Rep. 14 (1853) ... ... Q.
Brachycome ciliaris ... ... Q.
Calotis microcephala ... ... Q.
Aster ramulosus ... ... Q.
Vittadinia macrorrhiza ... ... Q.
Podocoma cuneifolia ... ... W.A.
Pterigeron liatroides ... ... N.A.
Pterigeron microglossus ... ... S.A.
Epaltes (Gynaphanes).
Pterocaulon sphacelatus ... ... W.A.
Podotheca fuscescens = Helipterum, Turczaninow in Bull. de Mosc. 80 (1851) ... ... M.fr.XII,22.
Podosperma gnaphaloides ... ... M.fr.XII,21.
Podosperma chrysanthum ... ... M.fr.XII,21.
Podosperma angustifolium ... ... M.fr.XII,21.
Leptorrhynchos squamatus ... ... Q.
Helipterum Kendallii = Podolepis Kendallii, F. v. M. (inedit.)
Helipterum Tietkensii ... ... W.A.,N.S.W
Helichrysum dealbatum ... ... V.
Helichrysum baccharoides ... ... N.S.W.
Helichrysum pholidotum = Humea squamata.
Eclipta alba ... ... N.A.
Wedelia verbesinoides ... ... S.A.
Bidens bipinnatus ... ... S.A.
Flaveria Australasica ... ... W.A.,S.A.

Erechtites mixa ... ... S.A.,V.
Erechtites prenanthoides ... ... S.A.
Candollea pilosa, Labillardière in Annal. du Mus. VI., 453, t. 63 (1805).
Candollea glauca, Labillardière in Annal. du Mus. VI., 454, t. 64 (1805).
Wahlenbergia (Strzeleckia).
Lobelia Benthamii ... ... S.A.
Isotoma petraea ... ... N.A.
Dampiera Brownii ... ... Q.
Dampiera diversifolia ... ... M.fr.I,120.
Dampiera incana ... ... N.A.
Leschenaultia hirsuta ... ... M.fr.XII,23.
Leschenaultia longiloba ... ... M.fr.XII,23.
Catosperma Muelleri ... ... S.A.
Scaevola depauperata ... ... N.A.
Scaevola globuliflora=S. globulifera.
Goodenia Chambersii ... ... N.A.
Goodenia Hassallii=G. Hassellii.
Mitrasacme pilosa ... ... S.A.
Logania floribunda ... ... Q.
Plantago from W. Turner (1538).
Ligustrum from W. Turner (1538).
Notelaea linearis ... ... Q.
Cynanchum floribundum ... ... W.A.
Gymnanthera nitida ... ... W.A,
Tylophora erecta ... ... N.A.
Ipomoea Turpethum ... ... N.A.
Ipomoea Muelleri... ... ... S.A.
Convolvulus from W. Turner (1538).
Polymeria angusta ... ... S.A.
Cuscuta from Bock (1539).
Cuscuta australis ... ... W.A.,S.A.
Solanum viride, Solander in G. Forster, prodr. 89 (1786).
Datura Leichhardtii ... ... W.A.
Lindernia, Allioni in misc. Taurin. III, 178 (1766), from Gagnebin (Pyxidaria).
Peplidium Muelleri ... ... W.A.,S.A.
Buechnera parviflora ... ... W.A.
Coldenia procumbens ... ... S.A.
Pollichia latisepalea ... ... Q.,N.A.
Clerodendrum lanceolatum ... ... W.A.
Spartothamnus teucriiflorus ... . W.A.
Eremophila oppositifolia ... ... W.A.
Eremophila Mitchelli ... ... S.A.
Eremophila exilifolia ... ... W.A.
Eremophila Freelingii ... ... W.A.
Eremophila Macdonnelli... ... ... N.A.
Eremophila Goodwinii ... ... N.A.
Eremophila platycalyx ... ... S.A.
Eremophila alternifolia ... ... W.A.
Eremophila scoparia ... ... W.A.
Prasophyllum Fimbria=P. Reichenbachii, F. v. M. (inedit.)
Chiloglottis Gunnii ... ... N.S.W.
Astelia (Funkia 1805).
Typhonium Brownii ... ... N.A.

Wolffia Michelii ... ... S.A.
Cyperus Eragrostis ... ... M.fr.XII,25.
Cyperus pulchellus ... ... M.fr.XII,25.
Cyperus gracilis ... ... M.fr.XII,25.
Cyperus tenellus ... ... M.fr.XII,25.
Cyperus squarrosus ... ... M.fr.XII,25.
Cyperus fulvus ... ... W.A.
Cyperus rotundus ... ... M.fr.XII,25.
Cyperus polystachys ... ... M.fr.XII,25.
Cyperus Iria ... ... M.fr.XII,25.
Cyperus decompositus ... ... M.fr.XII,25.
Scirpus grossus ... ... M.fr.XII,24.
Scirpus supinus ... ... M.fr.XII,25.
Scirpus litoralis ... ... M.fr.XII,25.
Cyathochaete clandestina ... ... M.fr.XII,25.
Cyathochaete diandra ... ... M.fr.XII,25.
Schoenus aphyllus ... ... M.fr.XII,26.
Schoenus ericetorum ... ... M.fr.XII,26.
Schoenus sphaerocephalus ... ... S.A.
Lepidospora tenuissima ... ... M.fr.XII,26.
Lepidosperma gladiatum ... ... M.fr.XII,26.
Lepidosperma filiforme ... ... M.fr.XII,26.
Cladium junceum... ... ... M.fr.XII,25.
Cladium schoenoides ... ... M.fr.XII,26.
Cladium trifidum ... ... M.fr.XII,26.
Cladium filum ... ... M.fr.XII,26.
Cladium melanospermum ... ... M.fr.XII,26.
Cladium Mariscus ... . M.fr.XII,25.
Cladium teretifolium ... ... M.fr.XII,25.
Cladium psittacorum ... ... M.fr.XII,25.
Cladium Radula ... ... S.A.
Caustis pentandra ... ... M.fr.XII,26.
Arthrostylis aphylla ... ... M.fr.XII,25.
Carex Brownii ... ... Q. M.fr.XII,26.
Carex tereticaulis... ... ... M.fr.XII,26.
Carex declinata ... ... M.fr.XII,26.
Carex breviculmis ... ... M.fr.XII,26.
Carex inversa ... ... M.fr.XII,26.
Carex acuta ... ... M.fr.XII,26.
Carex pumila ... ... M.fr.XII,26.
Tragus racemosus... ... ... W.A.
Andropogon Gryllus ... ... W.A.
Androphogon procerus ... ... W.A.
Anthistiria membranacea ... ... W.A.
Chionachne cyathopoda ... ... Q.
Sporobolus Lindleyi ... ... N.A.
Eriachne obtusa, W.A., S.A.
Eleusine Aegyptia=E. cruciata, Lamarck, tabl. encycl. et meth. I, 203 t. 48 (1791) W.A.
Setaria glauca ... ... W.A.
Setaria macrostachya ... ... W.A.
Eragrostis pilosa ... ... W.A.
Eragrostis tenella ... ... W·A.
Trirhaphis pungens ... ... Q.
Arundo Phragmites ... ... W.A.
Botrychium ternatum ... ... S.A.

SEQUENCE OF ORDERS ACCORDING TO PREDOMINANCE,

With Indications of their Numbers of Species.

Total number of Vasculares=8646. Dicotyledoneae=6897. Monocotyledoneae=1522. Acotyledoneae=227.

Leguminosae ... 1058	Tiliaceae 52	Rosaceae 17	Ericaceae 5
Myrtaceae 651	Loganiaceae ... 51	Anonaceae 16	Nymphaeaceae ... 4
Proteaceae 586	Apocyneae 48	Salicarieae 16	Magnoliaceae ... 4
Compositae ... 529	Droseraceae ... 46	Junceae 16	Samydaceae ... 4
Cyperaceae ... 372	Santalaceae ... 43	Monimieae 15	Elatineae 4
Gramineae 346	Pittosporeae ... 38	Celastrinae ... 15	Lineae 4
Epacrideae 273	Lauraceae 37	Olacinae 15	Cupuliferae ... 4
Orchideae 255	Meliaceae 36	Ebenaceae 15	Plumbagineae ... 4
Euphorbiaceae ... 224	Saxifrageae ... 35	Menispermeae ... 14	Plantagineae ... 4
Goodeniaceae ... 212	Campanulaceae ... 34	Stackhousieae ... 13	Dioscorideae ... 4
Filices 200	Polygaleae 32	Cycadeae 13	Gesneriaceae ... 3
Rutaceae 185	Portulaceae ... 32	Violaceae 12	Pedalinae 3
Liliaceae 161	Acanthaceae ... 30	Myrsinaceae ... 12	Philydreae 3
Rubiaceae 124	Coniferae 29	Phytolacceae 11	Nepenthaceae ... 2
Labiatae 124	Fluviales 29	Scitamineae ... 11	Guttiferae 2
Sterculiaceae ... 123	Combretaceae ... 27	Piperaceae 10	Malpighiaceae .. 2
Salsolaceae 112	Ficoideae 27	Aroideae 10	Burseraceae ... 2
Malvaceae 105	Palmae 26	Pandaneae 10	Connaraceae ... 2
Umbelliferae ... 103	Polygonaceae ... 25	Anacardiaceae ... 9	Caprifoliaceae 2
Sapindaceae ... 100	Loranthaceae . . 25	Hydrocharideae ... 9	Aquifoliaceae ... 2
Candolleaceae ... 95	Irideae 25	Xyrideae 9	Styraceae 2
Dilleniaceae ... 95	Caryophylleae ... 24	Rhizospermeae ... 9	Hydrophylleae ... 2
Amarantaceae ... 94	Capparideae ... 23	Flacourtieae ... 8	Burmanniaceae ... 2
Restiaceae 93	Casuarineae ... 23	Simarubeae ... 7	Typhaceae 2
Rhamnaceae ... 89	Cucurbitaceae ... 23	Geraniaceae ... 7	Myristicaceae ... 1
Amaryllideae ... 86	Gentianeae 23	Frankeniaceae ... 7	Papaveraceae ... 1
Solanaceae 80	Lentibularinae ... 23	Bignoniaceae ... 7	Hypericinae ... 1
Scrophularinae ... 78	Araliaceae 21	Crassulaceae ... 6	Ochnaceae 1
Myoporinae ... 78	Zygophylleae ... 20	Rhizophoreae ... 6	Podostemoneae ... 1
Verbenaceae ... 77	Jasmineae ... 20	Melastomaceae ... 6	Balonophoreae ... 1
Thymeleae 74	Sapotaceae 19	Primulaceae ... 6	Hamamelidae ... 1
Convolvulaceae ... 67	Commelineae ... 19	Lemnaceae 6	Elaeagneae ... 1
Urticaceae 63	Viniferae 18	Alismaceae ... 6	Cornaceae 1
Halorageae ... 60	Eriocauleae ... 18	Aristolochieae ... 5	Orobancheae ... 1
Asclepiadeae ... 60	Lycopodinae ... 18	Nyctagineae ... 5	Taccaceae 1
Cruciferae 54	Ranunculaceae ... 17	Onagreae 5	Roxburghiaceae ... 1
Asperifoliae .. 53	Tremandreae ... 17	Passifloreae ... 5	Pontederiaceae ... 1

Total number of Genera of Vasculares=1355.

INDEX OF ORDERS AND GENERA.

Abbottia 74
Abelmoschus 14
Abildgaardia 125
Abroma 15
Abrophyllum 48
Abrotanella 84
Abrus 41
Abutilon 14, 141
Acacia 43, 142
Acaena 47
Acalypha 20
Acanthaceae 99
Acanthocarpus 119
Acanthocladium 81
Acanthus 99
Achilleopsis 16
Achras 91
Achromolaena 82
Achrysum 83
Achyranthes 28
Acianthus 113
Acicalyptus 59
Aciphylla 53
Ackama 48
Acmene 59
Acomis 82
Acrachne 134
Acradenia 12
Acroclinium 80
Acronychia 12
Acrophyllum 48
Acrostichum 139
Acrotriche 105
Actephila 19
Actinocarpus 121
Actinodium 51
Actinopappus 82
Actinostachys 136
Actinostigma 10
Actinostrobus 109
Actinotus 63, 142
Adansonia 15
Adeliopsis 4
Adenanthera 43, 142
Adenanthos 66, 140
Adenochilus 113
Adenosma 97
Adenostemma 77
Adenostephanus 68
Adiantum 137
Adrastaea 1
Adriana 21, 141
Aegialinites 26
Aegialitis 26, 141
Aegiceras 91
Aegopogon 133
Aeschynomene 40, 142
Afzelia 43
Agastachys 67
Agathis 109
Agati 39
Ageratum 77
Aglaia 9
Aglaiopsis 9
Agnostus 71
Agonis 54
Anigozanthos 116
Agrimonia 47
Agropyron 135
Agrostis 133
Agrostocrinum 118
Ailantus 12
Aillya 88
Aira 133
Aizoon 31
Ajuga 102
Akania 24
Albizzia 47, 142
Aldrovanda 7
Alchemilla 47, 142
Alchornea 21
Alectryon 24
Alepyrum 123
Aleurites 20
Alisma 121
Alismaceae 121
Allania 118
Allantodea 138

Allophylus 24
Alopecurus 132
Alphitonia 60
Alpinia 114
Alsomitra 76
Alsophila 137
Althenia 121
Altingia 109
Alstonia 93
Alternanthera 28
Alysicarpus 40
Alyssum 5
Alyxia 93
Amanoa 20
Amarantaceae 28
Amarantus 29
Amaryllideae 115
Amblysperma 85
Ambrina 29
Ammannia 49, 142
Ammobium 82
Amomum 114
Amoora 9
Amorphophallus 120
Amorphospermum 91
Amperea 19
Amphibolis 121
Amphibromus 134
Amphicosmia 137
Amphipogon 133
Amphodus 41
Anacardiaceae 25
Anadenia 68
Anagallis 91
Anarthria 123
Ancana 3
Ancistrostigma 31
Ancistrum 47
Andersonia 108
Andrachne 19
Andrewsia 104
Andropogon 132,143
Androstemma 116
Androstoma 105
Aneilema 122
Anemone 1
Angianthus 82
Angiopteris 136
Angophora 57
Anguillaria 117
Anisacantha 30
Anisandra 102
Aniseia 94
Anisogonium 138
Anisolepis 80
Anisomeles 101
Anisopogon 134
Anisotome 63
Anonaceae 3
Anodopetalum 48
Anopterus 48
Anosporum 124
Antennaria 80
Antheidosorus 82
Anthericum 118
Anthistiria 132,143
Anthobolus 64
Anthocerastes 82
Anthocercis 96
Anthotium 87
Anthotroche 96
Anthosachne 135
Antiaris 22
Anticoryne 53
Antidesma 19
Antirrhoea 75
Antrophyum 139
Aotus 35,142
Aphananthe 21
Aphelia 123
Apium 63
Aplotaxis 85
Apluda 132
Apocyneae 92
Apodytes 64
Apolochlamys 82
Aponogeton 121
Apophyllum 5
Aporosa 20
Apostasia 114
Aquifoliaceae 63
Arabis 5
Araliaceae 61
Araucaria 109
Archeria 108
Archidendron 47
Archontophoenix 120
Ardisia 91, 105
Areca 120
Arenaria 27
Argophyllum 47
Argyrocome 80
Argyrodendron 15
Argyroglottis 81
Argyrophanes 81
Aristida 131
Aristolochia, 23
Aristolochiaceae 23
Aristotelia 17
Arnocrinum 119
Aroideae 120
Artanema 97
Arthratherum 132
Arthraxon 131
Arthrochilus 113
Arthrocnemum 30
Arthropodium 118
Arthropteris 139
Arthrostylis 129,143
Arthrotrichum 28
Arum 120
Arundinella 130
Arundo 135,143
Aryteria 24
Asclepiadeae 93
Asparagus 117
Asparagopsis 117
Asperella 132
Asperifoliae 99
Asperula 76,142
Aspidium 138
Asplenium 138
Asprella 132
Astartea 54
Astelia 117,143
Aster 78,142
Asteridea 80
Asterochiton 16
Asterolasia 10
Asteromyrtus 55
Asterotrichon 13
Astraea 52
Astrebla 134
Astroloma 105
Astrotricha 61,142
Atalantia 12
Atalaya 24,141
Atelandra 101
Atherocephala 108
Atherosperma 3
Athrixia 80
Athrodactylis 120
Athrotaxis 109
Athyrium 138
Atkinsonia 65
Atriplex 29,140,141
Atylosia 41
Australina 22
Avicennia 103
Aylmeyera 27
Azolla 136
Azorella 63
Babbagia 30,141
Babingtonia 53
Backhousia 59
Bacularia 120
Baea 98
Baeckea 53,142
Baeobotrys 91
Balanops 22
Balantium 137
Balaustion 54
Balfouria 93
Baloghia 21
Balonophora 23
Balonophoreae 23
Bambusa 135
Banksia 71,142
Barbarea 5,141
Barklya 42
Barringtonia 39
Bartlingia 36,118
Basilicum 100
Bassia 30,140,141
Batatas 94
Batratherum 131
Bauera 48
Bauhinia 42
Baumea 128
Baxteria 119
Beaufortia 56
Bedfordia 84
Beilschmiedia 3
Bellis 77
Bellendena 67
Benincasa 76
Berchemia 60
Bergia 8,141
Berrya 17
Bertolonia 104
Bertya 19,141
Beyeria 18,141
Beyeriopsis 18
Bidaria 94
Bidens 84,142
Bignoniaceae 99
Billardiera 7,141
Billiotia 54
Billotia 56
Bischoffia 20
Blackburnia 12
Blackwellia 6
Blancoa 116
Blandfordia 117
Bleasdalea 68
Blechnum 138
Blennodia 5
Blennospora 83
Blepharanthemum 13
Blepharocarya 25
Blitum 29
Blumea 79
Blyxa 115
Boehmeria 22
Boerhaavia 32
Bombax 15
Bonnaya 97
Boronia 10
Borya 119
Bosistoa 11
Bossiaea 37,140
Bothryodendron 61
Botrychium 136,143
Bouchardatia 12
Bowenia 110
Brachychiton 15
Brachycome 77,142
Brachyloma 107
Brachynema 48
Brachypterum 42
Brachysema 32
Brachyspatha 120
Brachystephium 77
Brackenridgea 10
Bradleia 19
Bramia 97
Brasenia 1
Brassaia 61
Breweria 95
Breynia 20
Bridelia 20
Brizopyrum 134
Brizula 123
Brochosiphon 99
Brombya 12
Bromidium 133
Bromus 135
Brucea 12
Bruguiera 50,142
Brunella 101
Brunonia 87
Bryonia 76
Bryonopsis 76
Buchanania 25
Buckinghamia 71
Buechnera 98,143
Bulbine 118
Bulbophyllum 110
Bulliardia 48
Bulweria 99
Bunnya 103
Burchardia 117
Burmannia 115
Burmanniaceae 115
Burnettia 113
Burgesia 32
Bursaria 7
Burseraceae 25,151
Burtonia 34
Busbeckea 5
Butomopsis 121
Byblis 8
Byronia 63

Cabomba 1
Cadaba 5
Cadellia 13
Caesalpina 42,142
Caesia 118
Cajanus 41
Cakile 6,141
Caladenia 113
Caladium 120
Calamus 119
Calandrinia 27
Calanthe 111
Caldesia 121
Caleana 113
Calectasia 119
Caletia 18
Caleya 113
Calliandra 47
Callicarpa 103
Callicoma 48
Callicysthus 41
Callipteris 138
Callistachys 32
Callistemon 55,140
Callitriche 50
Callitris 109
Calocephalus 83
Calochilus 112
Calogyne 88
Calomeria 82
Calonyction 94
Calopetalon 7
Calophyllum 8
Calosciadium 63
Calothamnus 56
Calorophus 124
Calostemma 116
Calostrophus 124
Calotis 78,142
Caltha 1

Calycopeplus 18
Calycothrix 52
Calyptranthes 59
Calyptrostegia 73
Calyptrostigma 18
Calystegia 95
Campanulaceae 85
Camphoromyrtus 53
Camptostemon 15
Campylanthera 7
Campynema 115
Cananga 3
Canarium 25
Canarvonia 71
Canavalia 41,142
Candollea 1,85,143
Candolleaceae 85
Canscora 90
Cansjera 63
Canthium 75,142
Capparideae 4
Capparis 5,141
Caprifoliaceae 76
Capsella 6,141
Carallia 50
Carapa 9
Carara 141
Cardamine 5
Cardiospermum 24
Cardwellia 71
Carex 129,140,143
Careya 60
Cargillia 92
Carissa 93
Carmichaelia 39
Carpha 127
Carpodontos 48
Carronia 4
Cartonema 122
Carumbium 21
Caryodaphne 3
Caryophylleae 27
Caryospermum 26
Caryota 120
Casearia 6
Cassia 42
Cassinia 82
Cassiniola 80
Cassytha 4
Castanospermum 42
Castanospora 24,141
Casuarina 22,141
Casuarineae 22
Catakidozamia 110
Cathartocarpus 42
Cathormion 47
Catosperma 87,142
Caulinia 41, 121
Caustis 129
Cedrela 9
Celastrineae 26
Celastrus 26,141
Celmisia 78
Celosia 29
Celtis 21
Cenarrhenes 67
Cenchrus 130
Centaurea 85
Centella 63
Centipeda 84
Centranthera 98
Centratherum 77
Centrolepis 123
Centropappus 84
Centrotheca 134
Centunculus 91
Cephaelis 75
Cephalipterum 83
Cephalosorus 82
Cephalotus 48
Ceranthes 92
Ceratogyne 84
Ceratopetalum 48
Ceratophyllum 50,142
Ceratopteris 137
Cerbera 93
Cercodia 49
Ceriops 50
Ceropegia 94
Cesatia 62
Chaetanthus 124
Chaetaria 132
Chaetospora 127
Chalcas 12
Chamaeraphis 130
Chamaescilla 118
Chamaesphaerion 83
Chamaexeros 119
Chamaelaucium 51
Chapelliera 127
Charlwoodia 117
Cheilanthes 138
Cheilococca 37
Cheiloplecton 138
Cheiranthera 7,141
Cheiroloma 78
Chenolea 30
Chenopodina 30
Chenopodium 29,141
Cheynia 54
Chilocarpus 92
Chilodia 101
Chiloglottis 114,143
Chionachne 132,143
Chionanthus 92
Chithonanthus 43
Chlamydia 117
Chlamyspermum 118
Chloanthes 103,140
Chloris 134
Chlorophytum 118
Chondrachne 127
Choretrum 64,142
Choriceras 19
Chorilaena 11
Choripetalum 91
Chorizandra 127
Chorizema 32,33
Chromochiton 82
Chrysobactron 118
Chrysocephalum 81
Chrysocoryne 82
Chrysodium 139
Chrysodiscus 80
Chrysogonum 83
Chrysophyllum 91
Chrysopogon 132
Chrysorrhoea 51
Chtonocephalus 83
Chuncoa 50
Cibotium 137
Cinna 133
Cinnamomum 4
Cissodendron 61
Citriobatus 7,141
Citrus 12
Cladium 128,143
Cladodes 21
Claoxylon 20
Clausena 12
Claytonia 27
Clematis 1
Cleistanthus 20
Cleome 4,141
Clerodendrum 103,143
Cleisostoma 111
Clianthus 39
Clidanthera 40
Clinostigma 120
Clitoria 40
Cloezia 59
Coatesia 12
Cocculus 4
Cochlospermum 6
Cocos 120
Codiaeum 21
Codonocarpus 32
Coelachne 133
Coelandria 110
Coelebogyne 21
Coelorrhachis 131
Coelospermum 75
Coenopteris
Coffea 75
Coilocarpus 30
Coldenia 100,143
Coleanthera 107
Coleocoma 79
Coleostylis 86
Coleus 100
Colobandra 101
Colobanthus 27
Colletia 61
Colmeiroa 48
Colocasia 120
Colubrina 60
Columbea 109
Combretaceae 50
Comesperma 8,141
Commelina 122
Commelineae 122
Commerçonia 16,141
Compositae 77
Conanthodium 81
Conchium 70
Coniferae 109
Connaraceae 32
Conospermum 66
Conostephiopsis 107
Conostephium 107
Conostylis 116
Conothamnus 56
Convolulaceae 94
Convolvulus 95,143
Conyza 79
Coprosma 75
Corchorus 17
Cordia 99
Cordyline 117
Corethrostylis 16
Coridochloa 130
Cornaceae 74
Coronopus 141
Correa 11
Corybas 112
Corymbis 111
Corymborchis 111
Corypha 120
Corynotheca 118
Corysanthes 112
Cosmelia 108
Costus 114
Cotula 84
Covellia 21
Craspedia 83
Crassulaceae 48
Crantzia 63
Crantziola 63.
Crepis 85
Cressa 95
Crinum 116
Crossolepis 83
Crossotoma 87
Crotalaria 38,140,142
Croton 20
Crowea 10
Cruciferae 5
Cryphia 101
Cryptandra 60,140
Cryptocarya 3
Cryptosema 32
Cryptostemon 51
Cryptostylis 112
Cucumis 76,142
Cucurbitaceae 76
Cudrania 22
Cupania 24
Cupuliferae 22
Curculigo 116
Curcuma 114
Cuscuta 95,143
Cuttsia 48
Cyanostegia 103
Cyanothamnus 10
Cyanotis 121
Cyathea 137
Cyathochaete 127,143
Cyathodes 105
Cyathopappus 83
Cyathopsis 105
Cybele 71
Cycadeae 109
Cycas 109
Cyclogyne 39
Cyclotheca 32
Cycnogeton 121
Cylicodaphne 4
Cylindrosorus 82
Cymbidium 111
Cymbonotus 85
Cyminosma 12
Cymodocea 121
Cynanchum 93,143
Cynoctomum 93
Cynodon 134
Cynoglossum 100,143
Cynometra 43
Cyperaceae 124
Cyperus 124,143
Cyphanthera 96
Cyphokentia 120
Cypselocarpus 32
Cyrtostylis 113
Cystanthe 108
Cystopteris 138

Dacrydium 109
Dactyloctenium 134
Daemia 94
Dalbergia 42
Dallachya 60
Damasonium 121
Dammara 109
Dampiera 87,142
Danthonia 134
Daphnandra 3
Daphne 74
Daphnobryon 74
Darea 138
Darlingia 71
Dasymalla 103
Dasypogon 119
Datura 96,143
Daucus 63
Davallia 137
Davidsonia 48
Daviesia 35,140
Darwinia 51
Decalophium 51
Decaspermum 59
Decaspora 107
Decazesia 83
Deeringia 29
Delabechea 15
Dendrobium 110
Denhamia 26
Denisonia 103
Dennstaedtia 137
Dentella 74
Deparia 137
Deplanchea 99
Depremesnilia 101
Derris 42
Deschampsia 133

Desmocladus 124
Desmodium 40
Desvauxia 123
Deyeuxia 133
Diaspasis 87
Dianella 117
Dicarpidium 15
Dicera 17
Dicerma 40
Dichelachne 133
Dichondra 95
Dichopetalum 63
Dichopogon 118
Dichosema 33
Dichrostachys 43
Dicksonia 137
Dicladanthera 140
Diclidopteris 138
Dicliptera 99
Dicrastylis 103
Dictyopteris 139
Didiscus 62,142
Didymanthus 30
Didymeria 11
Didymochaete 133
Didymonema 128
Didymotheca 32,142
Digitaria 130
Dilivaria 99
Dillenia 2
Dilleniaceae 1
Dillwynia 37
Dimeria 130
Dimetopia 62
Dimorpholepis 80
Diodontium 84
Dioscorea 115
Dioscorideae 115
Diospyros 92
Diotosperma 84
Diplachne 134
Diplacrum 129
Diplanthera 99,121
Diplarrhena 114
Diplaspis 63
Diplax 132
Diplazium 138
Diploglottis 24
Diplolaena 11
Diplolobium 39
Diplopeltis 25
Diplopogon 133
Diplora 138
Diplospora 74
Dipodium 111
Dipogonia 133
Dipteracanthus 99
Disarrhenum 132
Discaria 61
Dischidia 94
Discopodium 127
Discospermum 74
Diselma 109
Disemma 76
Disoon 104
Dissiliaria 19,141
Dissocarpus 30
Distichlis 135
Distichostemon 25
Distylis 88
Dithyrocarpus 122
Dithyrostegia 82
Diuris 112
Dodonaea 25,141
Dolichandra 99
Dolichandrone 99
Dolichos 41
Donatia 87
Donia 39
Doodia 138
Dopatrium 97
Doryanthes 116
Doryphora 3,141
Dracaena 117
Dracophyllum 109
Drakaea 113
Drapetes 74
Drosera 7
Droseraceae 7
Drummondita 11
Dryandra 72
Drymaria 27
Drymophila 117
Drymis 2
Drynaria 139
Duboisia 96
Dunbaria 41
Duperreya 95
Duttonia 80,104
Dysophylla 100
Dysoxylum 9
Dysphania 29,141

Eadesia 96
Earlia 99
Ebelingia 13
Ebenaceae 92
Ebermayera 99
Ecdeiocolea 123
Echinocarpus 17
Echinochloa 130
Echinocroton 21
Echinodorus 121
Echinopogon 133
Echinopsilon 30
Echinospermum 100
Echinus 21
Eclipta 83,142
Ectrosia 135
Ehretia 99
Ehrharta 132
Elachanthus 84
Elachocroton 21
Elachopappus 82
Elachothamnus 77
Elaeagneae 64
Elaeagnus 64
Elaeocarpus 17
Elaeodendron 26
Elaphoglossum 139
Elatine 8
Elatineae 8
Elatostemma 22
Elattostachys 24
Electrosperma 123
Eleochiton 125
Elephantopus 77
Elettaria 114
Eleusine 134,143
Eleutranthes 76
Elionurus 131
Elynanthus 127
Elythrophorus 135
Emblingia 5
Embothrium 71
Emex 31
Emilia 84
Emmenospermum 60
Emphysopus 77
Empleurosma 25
Encephalartos 110
Enchylaena 30,142
Enchysia 85
Endiandra 4,141
Enhalus 115
Enhydra 83
Entada 43
Enterolobium 47
Epacrideae 105
Epacris 108
Epaltes 79,140,142
Epiandra 128
Epiblema 112
Epicharis 9
Epilobium 49
Epipogum 111
Epipremnum 120
Epithinia 75
Epitriche 82
Eragrostis 135,143
Eranthemum 99
Erechtites 85,143
Eremaea 57
Eremodendron 104
Eremophila 104,140
Eremopyxis 52
Eremosyne 48
Eria 140
Eriachne 133,143
Erianthus 131
Ericaceae 105
Ericomyrtus 53
Erigeron 79
Eriocalia 63
Eriocauleae 123
Eriocaulon 123
Eriochilus 113
Eriochiton 30
Eriochlamys 82
Eriochloa 130
Eriocladium 82
Eriosema 41
Eriostemon 10
Eritrichum 100
Erodium 13
Erodiophyllum 77
Erycibe 94
Eryngium 63
Erysimum 5
Erythraea 90
Erythrina 41,142
Erythrorchis 111
Erythrophlaeum 43
Erythroxylum 13
Etaeria 111
Ethuliopsis 79
Euandra 129
Eucalyptus 57,141,142
Eucarya 64
Euchilopsis 37
Euchilus 36
Euchiton 79
Eucryphia 48
Eudesmia 57
Eugenia 59
Euneapogon 133
Euodia 12
Euonymus 26
Euosma 91
Eulalia 131
Eulophia 111
Eupatorium 77,141
Euphorbia 18,140,141
Euphorbiaceae 18
Euphoria 24
Eupomatia 3
Euphrasia 98
Euroschinus 25
Eurostorrhiza 129
Eurybia 78
Eurybiopsis 79
Eurycles 116
Euryomyrtus 53
Eustrephus 117
Eutacta 109
Eutassa 109
Eutaxia 32
Euthales 89
Euxolus 29
Exarrhena 100
Excaecaria 21
Exocarpos 64
Exocarya 127
Evolvulus 95
Eyrea 79

Fabricia 54
Fagus 22
Fagraea 90
Faradaya 103
Fatoua 22
Faustula 81
Fawcettia 4
Fenzlia 59
Festuca 134
Fieldia 98
Ficoideae 31
Ficus 21,140,141
Filices 136
Fimbristylis 125
Fischera 62
Fisquetia 120
Fitzalania 3
Fitzroya 109
Flacourtieae 6
Flagellaria 117
Flaveria 84,142
Flemingia 41
Fleurya 22
Flindersia 9
Floriscopa 122
Floscopa 122
Flueggea 20
Fluviales 121
Fontainea 21,141
Forstera 86
Forsteropsis 85
Fragosa 63
Francisia 51
Frankenia 26
Frankeniaceae 26
Franklandia 67
Freirea 22
Fremya 59
Frenela 109
Freycinetia 120
Friesia 17
Froebelia 105
Fugosia 15
Fuirena 126
Fusanus 64

Gahnia 128
Galactia 41
Galeola 111
Galium 76,142
Gamelythrum 133
Gamozygis 82
Ganophyllum 24
Gardenia 74
Garuga 25
Gastrodia 111
Gastrolobium 33,141
Gaultheria 105
Gaultiera 105
Geijera 12
Geitonoplesium 117
Geleznovia 10
Genetyllis 51
Geniostoma 90
Genoplesium 112
Genosiris 114
Gentiana 90
Gentianeae 90
Geococcus 5
Geodorum 111
Geophila 75
Geraniaceae 13
Geranium 13
Geum 47
Gilberta 82
Gilesia 15
Gillbeea 48
Gingidium 63

Glabaria 4
Gleichenia 137
Glinus 31
Glischocaryon 49
Glochidion 19
Glossodia 114
Glossogyne 84
Glossostigma 97
Glyceria 134
Glycine 41
Glycosmis 12
Glycyrrhiza 40
Gmelina 103
Gnaphalium 79
Gnaphalodes 83
Gnephosis 83
Gomphandra 64
Gompholobium34,142
Gomphrena 28
Gongronema 94
Goniocarpus 49
Goniophlebium 139
Goniopogon 78
Goniopteris 139
Goniotriche 28
Goodenia 88,143
Goodeniaceae 87
Goodenoughia 88
Goodia 38
Goodyera 111
Gossypium 15,141
Gouania 61
Gramineae 131
Grammatotheca 85
Grammitis 139
Graptophyllum 99
Gratiola 97
Graumuellera 121
Greevesia 14
Grevillea 68,140,142
Grewia 17
Grisebachia 120
Grumilia 75
Guettarda 75
Guettardella 75
Guichenotia 16
Guilandina 42
Gunnera 50
Gunnia 31,111
Guioia 24
Guttiferae 8
Gymnagathis 55
Gymnanthera 93,143
Gymnema 94
Gymnochaete 127
Gymnoschoenus 127
Gymnococca 73
Gymnogramme 139
Gymnogyne 84
Gymnopteris 139
Gymnopogon 93
Gymnosporia 26
Gymnostachys 120
Gymnostylis 84
Gymnothrix 131
Gynaphanes 142
Gynura 84
Gypsophila 27
Gyrocarpus 51
Gyrostachys 111
Gyrostemon 32
Gyrostephium 83

Habenaria 114
Haekeria 82
Haemodorum 115
Hakea 70,142
Halgania 100
Halfordia 12
Halocnemum 30
Halodule 121
Halophila 115
Halorageae 49
Haloragis 49
Halothamnus 13
Hamamelideae 48
Hamelinia 117
Hannafordia 16
Haplopappus 79
Hardenbergia 41
Harmogia 53
Harpulia 24
Harrisonia 13
Hartigshea 9
Haussmannia 99
Hearnia 9
Hedaroma 51
Hedera 61
Hedraianthera 9,141
Hedycarya 3
Hedyotis 74
Hedyscepe 120
Heleocharis 125
Helichrysum 81,142
Helicia 68
Helicteres 15
Heliophytum 100
Heliotropium 100
Helipterum80.140.142
Hellenia 114
Helmholtzia 122
Helmintostachys 136
Helophyllum 87
Helopus 130
Helostigma 85
Helothrix 127
Hemarthria 131
Hemarrhena 97
Hemiandra 101
Hemicarpus 62
Hemichroa 29
Hemiclidia 72
Hemicyclia 20
Hemigenia 101
Hemiphora 103
Hemiphues 63
Hemisteirus 28
Hemistemma 1
Hemistephus 1
Hemitelia 137
Heptopleurum 61
Heritiera 15
Hermannia 15
Hernandia 4
Herpestis 97
Herpolirion 118
Hetaeria 122
Heterachne 135
Heterodendron 24
Heterolaena 73
Heteropogon 132
Hewardia 117
Hexatheca 121
Hibbertia 1,141
Hibiscus 14,141
Hierochloe 132
Hippocratea 26
Hippocrepandra 18
Hodgkinsonia 75
Hodgsonia 118
Hodgsoniola 118
Holcus 132
Hologamium 131
Holotome 63
Homalium 6
Homalocalyx 52
Homalospermum 54
Homalostoma 108
Hormogyne 92
Hovea 38
Howittia 14
Hoya 94
Huanaca 63
Huegelia 62
Huenefeldia 78
Hugonia 13
Humata 137
Humea 82,142
Hutchimsia 6
Huttia 1
Hyalochlamys 82
Hyalolepis 82
Hyalosperma 80
Hybanthus 6,141
Hydriastele 120
Hydrilla 115
Hydrocharideae 115
Hydrocharis 115
Hydrocotyle 62
Hydroglossum 136
Hydrolea 99
Hydropeltis 1
Hydrophila 99
Hydnophytum 75
Hydrophylleae 99
Hylococcus 19
Hymenanthera 6,141
Hymenolepis 139
Hymenophyllum 137
Hymenosporum 7,141
Hymenotheca 32
Hypaelyptum 126
Hypelytrum 127
Hypericinae 8
Hypericum 8
Hypocalymma 54
Hypoestes 99
Hypolepis 139
Hypoporum 129
Hypolaena 124
Hypoxis 116
Hypserpa 4
Hyptiandra 12
Hyrtandra 22

Iberis 6
Ichnocarpus 93
Ilex 63
Ilysanthes 97
Imbricaria 53
Imperata 131
Indigofera 39,140,142
Ionidium 6
Iphigenia 117
Ipomoea 94,143
Irideae 114
Iris 114
Irwingia 61
Isachne 130
Isandra 140
Ischaemum 131
Iseilema 132
Isoetes 136
Isoetopsis 84
Isolepis 125
Isoloma 137
Isopogon 65,142
Isoschoenus 127
Isotoma 85,142
Isotropis 33,140
Ixauchenus 77
Ixiochlamys 79
Ixiolaena 80
Ixiosporum 7
Ixodia 82
Ixora 74,142

Jacksonia 34
Jambosa 59
Jansonia 32
Jasmineae 92
Jasminum 92
Johnsonia 119
Josephia 72
Josephinia 105
Junceae 122
Juncella 123
Juncus 122
Jungia 53
Jussieua 49
Justicia 99

Kaleniczenkia 32
Kamphusia 87
Kamptzia 59
Kelleria 74
Kennedya 41
Kentia 120
Kentiopsis 120
Kentropsis 30
Keraudrenia 16
Kernera 121
Kibara 3
Kingia 119
Kippistia 77
Kissodendron 61
Klanderia 101
Knoxia 76
Kochia 30.140,142
Koehnea 142
Kreyssigia 117
Kunzea 54,142
Kyllingia 124

Labiatae 100
Labichea 42,142
Laboucheria 43
Laccospadix 120
Lachnagrostis 133
Lachnocephalus 103
Lachnostachys 102
Lachnothalamus 85
Lactaria 93
Lagenaria 76
Lagenophora 77,142
Lagerstroemia 49,142
Lagrezia 29
Lagunaria 15
Lalage 37
Lamarchea 57
Lambertia 68
Lampocarya 128
Lamprochlaena 82
Lamprolobium 39
Laportea 22
Lappago 131
Lappula 100
Lasianthus 75
Lasiopetalum 16
Lastraea 138
Latourea 87
Latrobea 37
Lauraceae 3
Laurencia 85
Lavatera 13
Lawrencia 13
Lawrencella 81
Lawsonia 49
Laxmannia 118
Learosa 3
Lebediera 20
Ledgeria 111
Lcea 24,141
Leersia 132
Leewenhoekia 86
Legnophora 4
Leguminosae 32
Leichhardtia 4,94,109
Lemna 121
Lemnaceae 121
Lentibularinae 98
Leontopodium 80
Lepia 6
Lepidium 6,141

Lepidobolus 124
Lepidocoma 82
Lepidosperma 128
Lepidospora 128,143
Lepidozamia 110
Lepigonum 27
Lepironia 127
Lepistemon 94
Leptaspis 131
Leptatherum 131
Leptinella 84
Leptocarpus 124
Leptoceras 113
Leptochloe 134
Leptocyamnus 41
Leptocytisus 37
Leptolobium 41
Leptomeria 64
Leptopteris 137
Leptorrhynchos 80,142
Leptosema 32
Leptospermum 54
Leptotriche 83
Lepturus 131
Lepyrodia 123
Leschenaultia 87,142
Lespedeza 40,142
Lestibudesia 29
Leucas 101
Leucocarpum 26
Leucolaena 62
Leucophyta 83
Leucopogon 105
Leucothamnus 16
Leuzea 85
Lhotzkia 52
Libertia 114
Licuala 120
Ligusticum 63
Ligustrum 92,143
Liliaceae 117
Limnanthemum 90
Limnophila 97
Limnostachys 121
Limosella 97
Lindernia 97,143
Lindsaea 137
Lindsaya 137
Lineae 13
Linkia 67
Linociera 92
Linospadix 120
Linschotenia 87
Linum 13
Liparis 110
Liparophyllum 90
Lipocarpha 126
Lippaya 74
Lippia 102
Lissanthe 105
Litobrochia 138
Litsea 4
Livistona 120
Lobelia 85,142
Lobophyllum 100
Lopopogon 107
Logania 91,143
Loganiaceae 90
Lomandra 119
Lomaria 138
Lomariopsis 139
Lomatia 71
Lonchocarpus 42
Lopadocalyx 63
Lophoclinium 80
Lophostemon 58
Lopidium 136
Loranthaceae 64
Loranthus 65
Lotus 38
Loudonia 49
Lourea 40
Loxocarya 124
Lucuma 91
Ludwigia 49
Luffa 76
Luisia 140
Lumnitzera 50
Luzula 122
Luzuriago 117
Lycium 96
Lycopodineae 136
Lycopodium 136
Lycopus 101
Lyginia 123
Lygodictyon 136
Lygodium 136
Lyonsia 93
Lyperanthus 113
Lysanthe 68
Lysicarpus 59
Lysimachia 91
Lysinema 108
Lysiosepalum 17
Lythrum 49

Maba 92
Macadamia 68
Macaranga 21
Macarthuria 31,142
Macdonaldia 111
Macgregoria 26
Mackinlaya 61
Macropidia 116
Macropiper 23
Macropodia 116
Macropteranthes 51
Macrostegia 73
Macrostigma 47
Macrozamia 110
Maesa 91
Magnoliaceae 2
Mahernia 15
Maireana 30
Malacochaete 126
Malaisia 22
Malcolmia 5
Mallophora 103
Mallotus 21
Malpighiaceae 25
Malva 13
Malvaceae 13
Malvastrum 13
Manglesia 68
Manisuris 131
Mapania 127
Mappa 21
Marattia 136
Marianthus 7,141
Mariscus 124
Marlea 74
Marsdenia 94
Marsilea 136
Marquisia 75
Mayepea 92
Mazus 97
Megalotheca 124
Meionectes 50
Medicosma 12
Melachne 128
Melaleuca 55,140,142
Melanthesa 20
Melanthesiopsis 20
Melanthium 117
Melastoma 60
Melastomaceae 60
Melhania 15
Melia 9
Meliaceae 9
Melichrus 105
Melicope 12
Melochia 15
Melodinus 92
Melodorum 3
Melothria 77
Memecylon 60
Memorialis 22
Menkea 5
Meniocus 5
Meniscium 139
Menispermeae 4
Mentha 101
Merkusia 87
Mertya 61
Mesembrianthemum 31
Mesomelaena 127
Mesotriche 105
Methorium 15
Metrosideros 59
Mezoneurum 42
Michiea 107
Micraira 133
Micrantheum 18,141
Microcachrys 109
Microcarpaea 97
Microchloa 134
Microcladium 63
Microclisia 4
Microcorys 102
Microcybe 10
Microgyne 79
Microlaena 132
Microlepidium 6
Microlepia 137
Micromelum 12
Micromyrtus 52
Micropyxis 91
Microseris 85
Microstachys 21
Microstemma 94
Microtis 112
Microstylis 111
Miliusia 3
Milletia 39
Milligania 50,117
Millotia 82
Milnea 9
Mimosa 43
Mimulus 97
Mimusops 92
Minuria 77
Mirbelia 33,142
Mischocarpus 24
Mitragyne 90
Mitrasacme 90,142
Mitreola 90
Mniarum 27
Modecca 76
Molkenboeria 87
Mollinedia 3
Molloya 68
Mollugo 31,142
Momordica 76
Monacather 134
Monencyanthes 80
Monenteles 79
Monimieae 3
Monogramma 138
Monochoria 121
Monococcus 32
Monoploca 6
Monotaxis 18,141
Monotoca 105
Montia 27
Moonia 83
Moraea 114
Morelotia 128
Morgania 97
Morinda 75
Morisia 127
Morna 80
Moschosma 100
Motherwellia 61
Mucuna 41
Muehlenbeckia 31
Muehlenbergia 133
Mukia 77
Murraya 12
Musa 114
Myoporinae 104
Myoporum 104,140
Myosotis 100
Myosurus 1,141
Myriocephalus 82
Myriogyne 84
Myriophyllum 50
Myristica 3
Myristiceae 3
Myrmecodia 75
Myrsinaceae 91
Myrsine 91
Myrtaceae 51
Myrtus 59

Nablonium 83
Nageia 109
Najas 121
Nania 59
Nasturtium 5
Needhamia 107
Negria 98
Nelitris 59
Nelsonia 99
Nelumbium 1
Nelumbo 1
Nematoceras 112
Nematolepis 11
Nematophyllum 38
Nematopus 83
Nematostigma 114
Nemedra 9
Neoroepera 20
Neottopteris 138
Nepenthaceae 23
Nepenthes 23,140
Nephelium 24
Nephrodium 138
Nephrolepis 138
Neptunia 43
Nertera 75
Nesaea 49,142
Nesodaphne 3
Nesogenes 102
Nettoa 17
Neurachne 131
Newcastlia 102
Nicotiana 96
Niemeyera 91,114
Nipa 120
Niphobolus 139
Nitraria 13
Notelaea 92,143
Notholaena 138
Nothopanax 61
Notonerium 93
Notothixos 65
Nuytsia 65
Nyctagineae 32
Nymphaea 1
Nymphaeaceae 1
Nyssanthes 28

Oberonia 110
Obione 29
Ochnaceae 10
Ochrolasia 9
Ochrosia 93
Ocimum 100
Octoclinis 109
Oenanthe 63,142
Oenothera 49
Ogcerostylus 82
Olacinae 63
Olax 63

Oldenlandia 74,142
Olea 92
Oleandra 138
Olearia 78
Oliganthemum 79
Oligarrhena 107
Omalanthus 21,141
Omphacomeria 64
Onagreae 49
Oncosporum 7
Oncostylis 125
Onychosepalum 124
Opercularia 75,142
Ophioderma 136
Ophioglossum 136
Ophiorrhiza 74
Ophiuros 131
Opilia 63
Oplismenus 130
Orchideae 110
Oreobolus 127
Oreomyrrhis 63
Oreostylidium 87
Orites 68
Oritina 68
Ormocarpum 40
Ornithochilus 111
Orobanche 98
Orobancheae 98
Orthoceras 112
Orthopogon 130
Orthostemon 90
Orthosiphon 100
Orthotropis 33
Orthrosanthus 114
Oryza 132
Osbeckia 60
Osbornia 59
Oschatzia 63
Osmunda 137
Osteocarpum 30
Otanthera 60
Ottelia 114
Ourisia 97
Owenia 9
Oxalis 13
Oxleya 9
Oxycladium 33
Oxylobium 32
Oxymyrrhine 53
Oxystelma 93
Ozothamnus 81

Pachycornia 30
Pachygone 4
Pachynema 2,140
Pachysurus 83
Pacquerina 77
Pagetia 12
Palmae 119
Palmeria 3
Panax 61
Pancovia 24
Panaetia 80
Pandaneae 120
Pandanophyllum 127
Pandanus 120,140
Panicum 130
Panopsis 68
Papaver 4,141
Papaveraceae 4
Pappophorum 133
Paratropia 61
Parkeria 137
Parietaria 22
Parinarium 47
Paritium 14
Parsonsia 93
Paryphantha 52
Paspalum 130
Passiflora 76
Passifloreae 76
Patania 137
Patersonia 114
Pavetta 74
Pavonia 14
Pedalinae 105
Pedilonum 110
Pelargonium 13
Pellaea 138
Pelonastes 50
Peltophorum 42
Pemphis 49
Pennantia 64
Pennisetum 131
Pentaceras 12
Pentachondra 107
Pentacraspedion 133
Pentadactylon 67
Pentadynamis 38
Pentagonaster 54
Pentalepis 83
Pentapeltis 62
Pentataphrus 105
Pentapogon 133
Pentaptelion 105
Pentatropis 94
Peperomia 23,140,141
Peplidium 97,143
Pericalymma 54
Pericampylus 4
Pernettya 105
Peroa 105
Perojoa 105
Perotis 131
Persoonia 67,142
Petalolepis 81
Petalostigma 19
Petalostylis 42
Petermannia 115
Petrophila 65
Phacelothrix 82
Phaenopoda 80
Phajus 111
Phalangium 118
Phaleria 74
Phanerandra 105
Pharbitis 94
Phaseolus 41
Phebalium 10
Phegopteris 139
Pherosphaera 109
Philotheca 11,140
Philoxerus 28
Philydreae 122
Philydrum 122
Phlebocalymma 63
Phlebocarya 115
Phlebodium 138
Phoberos 6
Pholidia 104
Pholidiopsis 104
Pholidota 110
Phormium 117
Phragmites 135
Phreatia 110
Phyllachne 87
Phyllanthus 19,140,141
Phyllocalymma 82
Phyllocladus 109
Phyllodium 40
Phylloglossum 136
Phyllopappus 85
Phyllota 36
Phymatocarpus 56
Phymatodes 139
Physalis 95
Physolobium 41
Physopsis 103
Phytolaccaeae 32
Picrophyta 88
Pigea 6
Pileanthus 52
Pilitis 108
Pilularia 136
Pimelea 73,142
Piper 23
Piperaceae 23
Piptandra 53
Piptomeris 34
Piptostemma 82
Pisonia 32
Pithecolobium 47
Pithocarpa 82
Pithyrodia 103
Piptocalyx 3
Pipturus 22
Pittosporeae 6
Pittosporum 6,141
Plagianthus 13
Plagiogyria 138
Plagiolobium 38
Plagiosetum 131
Plantagineae 91
Plantago 91,143
Platycarpidium 62
Platycerium 139
Platychilum 38
Platylobium 37
Platyloma 138
Platyptelea 48
Platysace 62
Platytheca 9
Platyzoma 137
Plectranthus 100
Pleea 117
Pleiococca 12
Pleiogyne 84
Pleogyne 4
Pleopeltis 139
Pleurandra 1
Pleurandropsis 10
Pleurocarpaea 77
Pleurogramme 138
Pleuropappus 82
Plexaure 110
Plinia 59
Plintacanthus 134
Plokiostigma 26
Pluchea 79
Plumbagineae 26
Plumbago 26,141
Poa 134
Podocarpus 109
Podocoma 79,142
Podolepis 80,142
Podolobium 32
Podopetalum 42
Podosperma 80,140,142
Podostemoneae 23
Podotheca 80,140,142
Poecilodermis 15
Pogonia 104,111
Pogonolepis 82
Pogonolobus 75
Pogonatherum 131
Pogonetes 87
Pogostemon 100
Poiretia 38,108
Polanisia 4
Pollia 122
Pollichia 100,143
Pollinia 131
Polyalthia 3,140
Polycalymma 82
Polycarpaea 27
Polycarpon 27
Polycnemon 29
Polygala 8,141
Polygaleae 8
Polygonaceae 31
Polygonum 31,142
Polymeria 95,143
Polyosma 48
Polyphragmon 75
Polypodium 139
Polypogon 131
Polypompholyx 98
Polystichum 138
Polyzone 51
Pomaderris 60,142
Pomatotheca 31
Pomax 76
Ponceletia 108
Pongamia 42
Pontederiaceae 121
Popowia 3
Porana 95
Poranthera 18,141
Porospermum 61
Portulaca 27,141
Portulaceae 27
Posidonia 121
Potamogeton 121
Potamophila 132
Potentilla 47
Pothos 120
Pouzolzia 22,141
Pozoa 63
Pozoopsis 63
Prasophyllum 112,140,143
Pratia 85
Premna 103
Primulaceae 91
Prionosepalum 124
Prionotes 108
Pritzelia 62,122
Procris 22
Pronaya 7
Prostanthera 101,140
Proteaceae 65
Prunella 101
Pseudalangium 74
Pseudanthus 18,141
Pseudatalaya 24
Pseudomorus 22
Pseudopholidia 104
Psilotrichum 28
Psilotum 136
Psoralea 38,142
Psychotria 75
Psydrax 75
Pterigeron 79,142
Pteris 138
Pterocaulon 79
Pterochaete 80
Pterolobium 42
Pteroloma 40
Pterostigma 97
Pterostylis 113
Pterygopappus 80
Ptilotus 28,140,141
Ptychosema 39
Ptychosperma 120
Pultenaea 36
Pumilio 82
Punicella 54
Pycnolachne 102
Pycnosorus 83
Pycnospora 40
Pycreus 124
Pygmaea 97
Pyxidaria 143

Quinetia 82
Quintinia 48
Quoya 103

Ramphidia 111
Randia 74
Ranunculaceae 1
Ranunculus 1,141
Raoulia 80
Rapuntium 85

Ratonia 24
Reedia 129
Regelia 56
Remirea 127
Renealmia 114
Restiaceae 123
Restio 124
Rhabdothamnus 98
Rhagodia 29,141
Rhamnaceae 60
Rhamnus 60
Rhamphicarpa 98
Rhamphidia 111
Rhaphidophora 120
Rhipogonum 117
Rhizophora 50
Rhizophoreae 50
Rhizospermeae 136
Rhodamnia 59
Rhodanthe 80
Rhodomyrtus 59
Rhodosphaera 25
Rhopala 68
Rhus 25
Rhyncharrhena 94
Rhynchosia 41,142
Rhynchospora 127
Rhychostemon 16
Rhytidandra 74
Rhytidanthe 80
Rhytidosporum 7
Richea 83,108
Ricinocarpus 19,141
Riedleya 15
Rinzia 53
Rochelia 100
Roea 35
Roeperia 5,19
Rosaceae 47
Rostellaria 99
Rostellularia 99
Rotala 49
Rottboellia 131
Rottlera 21
Roupala 68
Rourea 32
Roxburghia 117
Roxburghiaceae 117
Rubiaceae 74
Rubus 47
Ruelingia 16
Ruellia 99
Rumex 31
Ruppia 121
Rutaceae 10
Rutidochlamys 80
Rutidosis 82
Ryssopteris 25

Saccharum 131
Saccolabium 111
Saccopetalum 3
Sagenia 138
Sagina 27,141
Salacia 26
Salicarieae 49
Salicornia 38,142
Salisia 54
Salomonia 8
Salsola 31,142
Salsolaceae 29
Salvia 101
Samara 91
Sambucus 76,142
Samolus 91
Samydaceae 6
Sandfordia 10
Santalaceae 64
Santalum 64
Sapindaceae 24
Sapindus 24
Saponaria 27,141
Sapota 91
Sapotaceae 91
Sarcocephalus 74
Sarcochilus 111
Sarcopetalum 4
Sarcostemma 94
Sarcozygium 13
Sarotes 16
Saussurea 85
Saxifrageae 47
Scaevola 87,142
Scalia 80
Scaliopsis 80
Schelhammera 117
Schenodorus 134
Schidiomyrtus 53
Schizaea 136
Schizachyrium 132
Schizeilema 63
Schizolema 137
Schizomeria 48
Schizopleura 56
Schmidelia 24
Schoberia 30
Schoenia 81
Schoenodum 123
Schoenolaena 62
Schoenus 127,143
Scholtzia 53
Schuermannia 51
Schwaegrichenia 116
Scirpidium 125
Scirpodendron 127
Scirpus 126,143
Scitamineae 114
Scleranthus 27
Scleria 129
Sclerochlamys 30
Sclerolaena 30
Scleroleima 84
Sclerothamnus 37
Scolopendrium 138
Scolopia 6
Scoparia 97
Scorzonera 85
Scottia 37
Scrophularinae 97
Scutellaria 101
Scyphiphora 75
Scyphocoronis 82
Seaforthia 120
Sebaea 90
Sebastiania 21
Secamone 93
Securinega 20,141
Selaginella 136
Selliera 88
Selliguea 139
Semecarpus 26
Senebiera 6,141
Senecio 84
Sentis 104
Sequoia 109
Serianthes 47
Seringea 16
Sersalisia 91
Sesbania 39,142
Seseli 63
Sesuvium 31
Setaria 130,143
Selwynia 4
Sheffieldia 91
Sicyos 77
Sida 14,141
Sideroxylon 91
Siebera 62
Siegesbeckia 83
Siemssenia 80
Sieversia 47
Siloxerus 82
Silphiospermum 77
Simarubeae 12
Simsia 66
Siphonodon 26
Sisymbrium 5
Sisyrinchium 114
Sium 63
Skinnera 94
Skirrhophorus 82
Sloanea 17
Smilax 117,140
Smithia 40
Solanaceae 95
Solanum 95,143
Soleniscia 105
Solenogyne 77
Solenostigma 21
Soliva 84
Sollya 7
Sonneratia 60
Sonzaya 25
Sophora 42
Sopubia 98
Sorghum 132
Sorostachys 124
Sowerbaea 118
Spadostyles 36
Spanoghea 24
Sparganium 120
Spartothamnus 102,143
Spathoglottis 111
Spathodea 99
Spergularia 27
Spermacoce 76
Spermalepis 59
Spermaxyrum 63
Sphaeranthos 79
Sphaerolobium 35
Sphaeromorphaea 79
Sphaeropus 129
Sphenotoma 109
Sphincterostoma 108
Spiculaea 113
Spinifex 131
Spilanthes 84
Spiranthes 111
Spirodela 121
Spiropodium 79
Spodiopogon 131
Spondias 26
Sponia 21
Sporobolus 133,143
Sprengelia 108
Spyridium 60
Stachystemon 18
Stackhousia 26,141
Stackhousieae 26
Statice 26
Stawellia 119
Steetzia 78
Stegania 138
Steiroglossa 77
Stekovia 88
Stellaria 27
Stemodia 97
Stenanthemum 60
Stenanthera 105
Stenocarpus 71
Stenochilus 104
Stenochlaena 138
Stenodiscus 60
Stenopetalum 5
Stenosperma 84
Stephania 4
Sterculia 15
Sterculiaceae 15
Stipa 132
Stirlingia 66
Stomarrhena 105
Strangea 68
Stravadium 59
Streblorrhiza 39
Streptachne 132
Streptothamnus 6
Streptocarpus 98
Streptoglossa 79
Strobilanthes 99
Stromatopteris 137
Strongylodon 41
Strongylospermum 84
Strychnos 90
Strzeleckia 9,143
Stuartina 79
Sturmia 110
Sturtia 15
Stylidium 74, 85
Stylobasium 47
Stylocoryne 74
Stylolepis 80
Styloncerus 82
Stylurus 68
Stypandra 118
Styphelia 105
Styraceae 92
Suaeda 30
Suriana 13
Symphyomera 84
Symphyomyrtus 57
Symphyonema 67
Symphyopetalum 11
Symplocos 92
Synaphea 66
Synaptantha 74
Syncarpia 59
Synophlebium 137
Synostemon 19
Synoum 9
Syzygium 59
Swainsona 39,141,142
Swammerdamia 81

Tabernaemontana 93
Tacca 115
Taccaceae 115
Taeniophyllum 111
Talinum 27
Tamarindus 42
Tapeinocheilos 114
Tarrietia 15
Tasmania 2
Tecoma 99
Telanthera 28
Telmatophace 121
Telopea 71
Temminkia 87
Templetonia 38
Tenagocharis 121
Tephrosia 39,142
Terminalia 50
Tersonia 32
Tetracarpaea 48
Tetracera 2
Tetracheilus 43
Tetragonella 31
Tetragonia 30,142
Tetralepis 127
Tetralobus 98
Tetranthera 4
Tetrapasma 61
Tetraphora 53
Tetraphylax 88
Tetrarrhena 132
Tetratelia 5
Tetratheca 9
Teucrium 102
Thalamia 109
Thamnopteris 138
Thecanthes 73
Theleophyton 29
Thelychiton 110
Thelymitra 111
Therogeron 77

Thesium 64
Thespesia 15
Thespidium 79
Thlaspi 6
Thomasia 16
Thouarsea 131
Thozetia 94
Threlkeldia 30
Thrixpermum 111
Thryptomene 52,142
Thuarea 131
Thunbergia 99,140
Thymeleae 73
Thyrsacanthus 99
Thysanotus 118
Tiaridium 100
Tiliaceae 17
Tillaea 48
Tinonius 75
Tinospora 4
Titania 110
Tittmannia 97
Tmesipteris 136
Todea 137
Torenia 97
Tournefortia 100
Toxanthus 82
Trachycaryon 21
Trachymene 62,142
Trachypogon 132
Tragia 21,141
Tragus 131,143
Trema 21
Tremandra 9
Tremandreae 9
Trianthema 31,142
Tribonanthes 116
Tribulopsis 13
Tribulus 13,140,141
Trichanthodium 83
Trichelostylis 125
Trichinium 28
Trichobasis 56
Trichobolus 32
Trichocline 85
Trichodesma 100
Trichodium 133
Tricholoma 97
Trichomanes 137
Trichosanthes 76,142
Trichosiphon 15
Trichostegia 80
Tricondylus 71
Tricoryne 118
Tricostularia 127
Trigastrotheca 31
Triglochin 121
Trigonella 38
Trineuron 84
Triodia 135
Triphelia 51
Tripladenia 117
Tripetelus 76
Tripterococcus 26
Triptilodiscus 80
Trirhaphis 135,143
Trisetum 133
Tristania 58
Tristaniopsis 58
Tristellateia 25
Tristichocalyx 4,141
Trithuria 123
Triticum 135
Triumfetta 17
Trochocarpa 107
Trollius 1
Trymalium 60
Turraea 9
Turritis 5
Tylophora 94,143
Typha 120
Typhaceae 120
Typhonium 120,143

Ulmus 21
Umbelliferae 62
Uncinia 129
Ungeria 15
Urachne 132
Uragoga 75
Uralepis 134
Uraria 40,142
Urena 14,141
Urocarpus 10
Urodon 36
Urostigma 21
Urtica 22
Urticaceae 21
Utricularia 98
Uvaria 3
Uvedallia 97

Vachellia 43
Vaginularia 138
Vallisneria 115
Vandellia 97
Veitchia 120
Velleya 89,140
Ventenata 85
Ventenatia 105
Ventilago 60
Verbena 102
Verbenaceae 102
Vernonia 77
Veronica 97
Verreauxia 87
Verticordia 51
Vigna 41,142
Vilfa 133
Villarsia 90
Villaresia 64
Viminaria 35
Vinca 93
Vincetoxicum 93
Viniferae 23
Viola 6
Violaceae 6
Viraya 80
Viscum 64,142
Vitex 103
Vitis 23
Vittaria 137
Vittadinia 79,142
Vlamingia 6
Vulpia 134

Wahlenbergia 85,143
Waitzia 80
Walcottia 102
Waltheria 15,141
Wartmannia 21
Webera 74
Wedelia 83,142
Wehlia 53
Weinmannia 48
Westonia 38
Westringia 102
Wichurea 60
Wickstroemia 74
Wilkia 5
Wilkiea 3
Wilsonia 95
Wistaria 39
Wittsteinia 105
Wolffia 121,143
Wollastonia 83
Woodwardia 138
Woollsia 108
Wormia 1
Wrightia 93
Wrixonia 101
Wurmbea 117

Xanthochrysum 81
Xanthophyllum 8
Xanthorrhoea 119
Xanthosia 62
Xanthostemon 59
Xanthoxylum 12
Xerochloa 131
Xerosollya 7
Xerotes 119
Ximenia 63
Xiphopteris 139
Xylocarpus 9
Xylomelum 68
Xylosma 6
Xyridanthe 80
Xyrideae 122
Xyris 122

Youngia 85

Zaleya 31,142
Zamia 110
Zannichellia 121
Zanonia 76
Zapania 102
Zehneria 77
Zichya 41
Zieria 10
Zizyphus 60
Zornia 40
Zostera 121
Zosterostylis 112
Zoysia 131
Zygia 47
Zygomenes 122
Zygophylleae 13
Zygophyllum 13

M'Carron, Bird and Co., Printers, 37 Flinders Lane West, Melbourne.